

# Financial Mathematics 32000

## Lecture 6

Roger Lee

2025 April 28

# Variance reduction techniques

Standard error = number of stdev to cover confidence interval times the stdev  
 number of stdev to cover confidence interval is 1.96 (~2) for 95% confidence interval

Recall the radius of a  $100(1 - p)\%$  confidence interval is

$$\Phi^{-1}\left(1 - \frac{p}{2}\right) \sqrt{\text{Var}(\hat{C}_M)}$$

Variance reduction techniques  $\rightarrow$  estimators with lower Var than  $\hat{C}_M$

- ▶ Introduce dependence across simulations: we want to have a smaller variance than  $\text{Var}(\hat{C}_M)$ 
  - ▶ Antithetic variates  $\hat{C}_M^{\text{av}}$
- ▶ Simulate something other than the contract's payoff:
  - ▶ Control variates  $\hat{C}_M^{\text{cv}}$
  - ▶ Conditional Monte Carlo  $\hat{\sigma}_M^{\text{cmc}}$
- ▶ Modify the probability distribution:
  - ▶ Importance sampling  $\hat{C}_M^{\text{ris}}$

(Increasing  $M$  reduces Var, but not described as “variance reduction”)

Variance reduction: Antithetic variates

Variance reduction: Control variates

Variance reduction: Importance sampling

Variance reduction: Conditional MC

## Antithetic variates

Let  $Y$  be discounted payoff. So

$$C = \mathbb{E}Y.$$

generate  $M$  simulations and take the average of the  $M$  simulations, where each simulation is IID as  $Y$

Ordinary MC:  $\hat{C}_M = \frac{1}{M}(Y_1 + \dots + Y_M)$ , where each  $Y_m$  is IID as  $Y$ .

Example: Call on stock under GBM:

$$Y_m = e^{-rT}(S_0 e^{(r-\sigma^2/2)T + \sigma\sqrt{T}Z_m} - K)^+$$

where we generate the  $Z_m$  to be IID  $N(0, 1)$  for  $m = 1, \dots, M$ .

**Antithetic Variates:** If you draw a realization from the  $q$ th percentile, then you should *also draw one from the  $(100 - q)$ th percentile*.

Symmetry of normal  $\Rightarrow$  for each realization of  $Z$ , rerun also with  $-Z$ .

## Antithetic variates

In the same example, for each  $m = 1, \dots, M$ , let

$$\tilde{Z}_m := -Z_m \quad \text{if } Z_m = 93\text{th percentile, } \tilde{Z}_m \text{ is the 7th percentile}$$

$$\tilde{Y}_m := e^{-rT} (S_0 e^{(r-\sigma^2/2)T + \sigma\sqrt{T}\tilde{Z}_m} - K)^+$$

$$Y_m^{\text{av}} := \frac{Y_m + \tilde{Y}_m}{2} \quad \text{take the pair average of the original and antithetic simulation}$$

$$\hat{C}_M^{\text{av}} := \frac{1}{M} (Y_1^{\text{av}} + Y_2^{\text{av}} + \dots + Y_M^{\text{av}}) \quad \text{MC estimate is the average of all } M \text{ pair averages}$$

This is the antithetic-variate Monte Carlo estimate.

Its expectation is

$$\mathbb{E}\hat{C}_M^{\text{av}} = \frac{1}{M} \sum_m \mathbb{E}Y_m^{\text{av}} = \mathbb{E}\left(\frac{Y + \tilde{Y}}{2}\right) = \frac{\mathbb{E}Y + \mathbb{E}\tilde{Y}}{2} = \mathbb{E}Y = C$$

so  $\hat{C}_M^{\text{av}}$  is an unbiased estimate of  $C$ .

## Antithetic variates: variance analysis

Variance of AV estimate is **Mon**

$$\begin{aligned}\text{Var}(\hat{C}_M^{\text{av}}) &= \text{Var}\left(\frac{1}{M} \sum_m Y_m^{\text{av}}\right) = \frac{1}{M^2} M \text{Var}(Y^{\text{av}}) \\ &= \frac{1}{M} \text{Var}\left(\frac{Y + \tilde{Y}}{2}\right) = \frac{1}{M} \frac{\text{Var}Y + 2\text{Cov}(Y, \tilde{Y}) + \text{Var}(\tilde{Y})}{4} \\ &= \frac{1}{M} \left(\frac{1}{2} \text{Var}Y + \frac{1}{2} \text{Cov}(Y, \tilde{Y})\right)\end{aligned}$$

**Y and Y\_tilde is iid  
therefore their var is the  
same**

Comparing to ordinary MC,

$$\text{Var}(\hat{C}_M) = \frac{1}{M} \text{Var}(Y).$$

**MC uses M = 1,000 data  
Antithetic uses 2M = 2,000 (pairs)  
Not fair**

and  $\text{Cov}(Y, \tilde{Y}) \leq \sqrt{\text{Var}(Y)\text{Var}(\tilde{Y})} = \text{Var}(Y)$  so  **$\text{Var}(\hat{C}_M^{\text{av}}) \leq \text{Var}(\hat{C}_M)$ .**

But maybe this overstates the benefit of AV.

## Antithetic variates: variance analysis

Maybe better to compare **Mon**

$$\text{Var}(\hat{C}_M^{\text{av}}) = \frac{1}{M} \left( \frac{1}{2} \text{Var}Y + \frac{1}{2} \text{Cov}(Y, \tilde{Y}) \right)$$

against

$$\text{Var}(\hat{C}_{2M}) = \frac{1}{2M} \text{Var}(Y).$$

So  $\text{Var}(\hat{C}_M^{\text{av}})$  is smaller iff  $\text{Cov}(Y, \tilde{Y}) < 0$ .

This often holds, but for some payoffs, it doesn't.

**For straddle, the antithetic doesn't work**

## Antithetic variates: variance analysis

Because

$$\text{Var}(\hat{C}_M^{\text{av}}) = \frac{1}{M} \text{Var}(Y^{\text{av}}),$$

an asymptotic  $100(1 - p)\%$  confidence interval has endpoints

$$\hat{C}_M^{\text{av}} \pm \Phi^{-1}\left(1 - \frac{p}{2}\right) \frac{\hat{\sigma}_M^{\text{av}}}{\sqrt{M}}$$

where

$$\hat{\sigma}_M^{\text{av}} := \sqrt{\frac{1}{M-1} \sum_m (Y_m^{\text{av}} - \hat{C}_M^{\text{av}})^2}.$$

Sum of squared difference between the pair average and the true value o the pair average

Note that  $\hat{\sigma}_M^{\text{av}}$  is the sample standard deviation of the *pair averages*, not of the individual realizations.

Dividing it by  $\sqrt{M}$  gives the standard error  $\hat{\sigma}_M^{\text{av}}/\sqrt{M}$ , an estimate of the standard deviation of  $\hat{C}_M^{\text{av}} = \frac{1}{M} \sum_m Y_m^{\text{av}}$ .



Variance reduction: Antithetic variates

Variance reduction: Control variates

Variance reduction: Importance sampling

Variance reduction: Conditional MC

## Control variates

A control variate  $Y^*$  is a random variable, related to  $Y$ , such that

$C^* := \mathbb{E}Y^*$  is known (e.g. via explicit formula). Examples:

- ▶ Let  $dS_t = \sigma_t S_t dW_t$ , where  $\sigma_t$  is stochastic. If  $Y$  is the discounted payoff of a call on  $S$ , we can let  $Y^*$  be the discounted payoff of a call on  $S^*$  where  $dS_t^* = \sigma S_t^* dW_t$ . So  $C^*$  is known (B-S formula). Here  $\sigma$  is some constant that approximates the  $\sigma_t$  process.
- ▶ If  $Y$  is discounted payoff of Asian-style call on GBM:

$$Y = e^{-rT} \left( \frac{1}{T} \int_0^T S_t dt - K \right)^+$$

Can let  $Y^*$  be the discounted *European* call payoff on the GBM.

Or let  $Y^*$  be the discounted payoff of a *geometric Asian* call:

$Y^* = e^{-rT} \left( e^{\frac{1}{T} \int_0^T \log S_t dt} - K \right)^+$ . In either case,  $C^*$  has a formula.

## Control variates

We want to price asian-style option, but we cannot, so use a European Option (using BS)

$Y$  is asian, European is  $Y^*$ . No need to hedge one-to-one. That's why there's beta.

The control variate technique simulates the **difference**  $Y - \beta Y^*$ ,

where  $\beta$  is a constant. The difference has expectation

$$\mathbb{E}(Y - \beta Y^*) = C - \beta C^*$$

and variance  $\text{Var}(Y - \beta Y^*)$  which is, we hope, small. Therefore

$$C = \mathbb{E}Y + \beta(C^* - \mathbb{E}Y^*).$$

Replacing expectations by sample averages, we define the control variate Monte Carlo estimate of  $C$  to be

$$\hat{C}_M^{\text{cv},\beta} := \bar{Y}_M + \beta(C^* - \bar{Y}_M^*) \quad \text{where} \quad \bar{Y}_M := \frac{1}{M}(Y_1 + \cdots + Y_M)$$

$$\bar{Y}_M^* := \frac{1}{M}(Y_1^* + \cdots + Y_M^*)$$

expectation of  $Y_1^*, Y_2, \dots$  is  $C^*$

second term is 0 -->  $C^* C^*$

MC Estimate of  $C$  becomes only the first term

and we use the *same* pseudo-random numbers generate  $Y_m$  and  $Y_m^*$ .

## Control variates: variance analysis

Expectation:

$$\mathbb{E}\hat{C}_M^{\text{cv},\beta} := \mathbb{E}\hat{C}_M + \beta(C^* - C^*) = C$$

so  $\hat{C}_M^{\text{cv},\beta}$  is an unbiased estimate of  $C$ .

Variance:

$$\begin{aligned}\text{Var}(\hat{C}_M^{\text{cv},\beta}) &= \text{Var}(\bar{Y}_M - \beta\bar{Y}_M^*) = \frac{1}{M} \text{Var}(Y - \beta Y^*) \\ &= \frac{1}{M} [\text{Var}(Y) - 2\beta \text{Cov}(Y, Y^*) + \beta^2 \text{Var}(Y^*)]\end{aligned}$$

Now choose  $\beta$  to minimize this. Obtain

$$\beta_{\text{optimal}} = \text{Cov}(Y, Y^*) / \text{Var}(Y^*)$$

$$\text{and } \text{Var}(\hat{C}_M^{\text{cv},\beta_{\text{optimal}}}) = \text{Var}(\hat{C}_M) \times (1 - \text{Corr}^2(Y, Y^*)) \leq \text{Var}(\hat{C}_M)$$

## Control variates: optimizing the coefficient

Unfortunately the Cov and Var are usually not both known.

But we can use the *sample* covariance and variance. Let

$$\hat{\beta} := \frac{\sum_m (Y_m - \bar{Y}_M)(Y_m^* - \bar{Y}_M^*)}{\sum_m (Y_m^* - \bar{Y}_M^*)^2}$$

This is the OLS coefficient on the control variate,  $Y^*$  in a regression of  $Y$  on  $Y^*$  and an intercept

And define

$$\hat{C}_M^{\text{cv}, \hat{\beta}} := \bar{Y}_M + \hat{\beta}(C^* - \bar{Y}_M^*).$$

Note  $\mathbb{E}(C^* - \bar{Y}_M^*) = 0$  does not imply  $\mathbb{E}[\hat{\beta}(C^* - \bar{Y}_M^*)] = 0$ , so this creates **bias** in the estimate:

$$\mathbb{E}\hat{C}_M^{\text{cv}, \hat{\beta}} = C + \mathbb{E}[\hat{\beta}(C^* - \bar{Y}_M^*)].$$

As  $M \rightarrow \infty$  the bias  $\rightarrow 0$ .

## Control variates: confidence intervals

To compute confidence intervals for fixed  $\beta$ , note that

$$\hat{C}_M^{\text{cv},\beta} = \frac{1}{M} \sum_m Y_m^{\text{cv},\beta}$$

where the independent simulations

$$Y_m^{\text{cv},\beta} := Y_m + \beta(C^* - Y_m^*)$$

have an estimated variance

$$\frac{1}{M-1} \sum_m (Y_m^{\text{cv},\beta} - \hat{C}_M^{\text{cv},\beta})^2.$$

Let  $\hat{\sigma}_M^{\text{cv},\beta}$  be the square root of this estimated variance.

## Control variates: confidence intervals

Note that

$\hat{\sigma}_M^{\text{cv},\beta}$  = estimated standard deviation of *one* of the  $Y_m^{\text{cv},\beta}$

$\hat{\sigma}_M^{\text{cv},\beta} / \sqrt{M}$  = estimated st. dev. of  $\hat{C}_M^{\text{cv},\beta}$ , the *average* of the  $Y_m^{\text{cv},\beta}$

Then an asymptotic  $100(1 - p)\%$  confidence interval has endpoints

$$\hat{C}_M^{\text{cv},\beta} \pm \Phi^{-1}\left(1 - \frac{p}{2}\right) \frac{\hat{\sigma}_M^{\text{cv},\beta}}{\sqrt{M}}.$$

If instead of  $\beta$  we use  $\hat{\beta}$  estimated from those  $M$  simulations, the variance estimate is biased for finite  $M$ , but as  $M \rightarrow \infty$  its bias and variance  $\rightarrow 0$ , so the same confidence interval calculation (with  $\hat{\beta}$  in place of  $\beta$ ) is asymptotically valid.

Variance reduction: Antithetic variates

Variance reduction: Control variates

Variance reduction: Importance sampling

Variance reduction: Conditional MC



## Motivating example

Using Monte Carlo, a rare event with a big payoff is harder to price than a more frequent event with a smaller payoff.

- ▶ Consider an asset that pays 100 dollars with probability 2% (under some pricing measure), and pays zero otherwise.

Its price is 2 and the payoff variance is  $10000 \times 0.02 - 2^2 = 196$

- ▶ Now consider an asset that pays 10 dollars with probability 20%, zero otherwise.

Its price is 2. But its payoff variance is only  $100 \times 0.20 - 2^2 = 16$

So maybe if we *sample from a different distribution* and *modify the payoff* we can preserve the expectation, while decreasing the variance.

# Importance sampling

Let  $X$  be a random vector in  $\mathbb{R}^N$ , with density function  $f : \mathbb{R}^N \rightarrow \mathbb{R}$ .

Let  $h : \mathbb{R}^N \rightarrow \mathbb{R}$ . Think of  $h$  as a discounted payoff function.

We want to estimate

$$C := \mathbb{E}h(X) = \int h(x)f(x)dx$$

Ordinary Monte Carlo:

Instead of  $f$ , maybe draw from  $g$  distribution that can provide us with a smaller variance

$$\hat{C} := \frac{1}{M} \sum_{m=1}^M h(X_{[m]})$$

where  $X_{[1]}, \dots, X_{[M]}$  are IID random draws from density  $f$ .

Can apply to  $X$  = some financial variable (such as an asset price), or  
can apply to  $X$  = as the randomness that drives the dynamics (such as a vector of Brownian increments).

only take the integral in the domain where the payoff times density ( $f$ ) is non-zero

## Change of measure

Let  $g$  be any density on  $\mathbb{R}^N$  that vanishes only where  $fh$  does, meaning  $f(x)h(x) = 0$  if  $g(x) = 0$ . Let  $A := \{x : f(x)h(x) \neq 0\}$ . Then

$$C = \int_A h(x)f(x)dx = \int_A h(x)\frac{f(x)}{g(x)}g(x)dx = \mathbb{E}^*\left[h(X)\frac{f(X)}{g(X)}\right]$$

where  $\mathbb{E}^*$  denotes expectation with respect to a new measure  $\mathbb{P}^*$  under which  $X$  has density  $g$ .

The *importance sampling* Monte Carlo estimate of  $C$  is

$$\hat{C}^{\text{is}} := \frac{1}{M} \sum_{m=1}^M h(X_{[m]}) \frac{f(X_{[m]})}{g(X_{[m]})}$$

where  $X_{[1]}, \dots, X_{[M]}$  are IID random draws from density  $g$ , which we call the *importance sampling* density.

## Change of measure

So we have made two changes to the ordinary MC estimate

- ▶ Change the distribution from which  $X$  is sampled. Draw from density  $g$  instead of  $f$ .
- ▶ Change the payoff. Instead of  $h(X)$ , it becomes  $h(X)$  times the *Radon-Nikodym derivative* or *likelihood ratio*  $f(X)/g(X)$ .

What about bias and variance?

- ▶ By construction,  $\hat{C}^{\text{is}}$  is an unbiased estimator of  $C$  because

$$\mathbb{E}^* \hat{C}^{\text{is}} = \frac{1}{M} \sum_{m=1}^M \mathbb{E}^* \left[ h(X_{[m]}) \frac{f(X_{[m]})}{g(X_{[m]})} \right] = C$$

- ▶ To minimize variance, try to choose  $g$  such that  $hf/g$  is constant on the set where  $g \neq 0$ . Try to “flatten the curve” (the payout  $h$ ).

## Variance

The importance-sampling MC estimate  $\hat{C}^{\text{is}}$  has (wrt  $\mathbb{P}^*$ ) variance

$$\frac{1}{M} \text{Var}^* \frac{h(X)f(X)}{g(X)} = \mathbb{E}^* \frac{h(X)^2 f(X)^2}{g(X)^2} - C^2 = \mathbb{E} \left[ h(X)^2 \frac{f(X)}{g(X)} \right] - C^2$$

which can be bigger or smaller than the ordinary MC variance:

$$\frac{1}{M} \text{Var} h(X) = \mathbb{E}[h(X)^2] - C^2.$$

How to choose  $g$  to minimize  $\text{Var}^* \hat{C}^{\text{is}}$ ? Answer: if  $h \geq 0$  and

$$g(x) = \text{constant} \times h(x)f(x)$$

then

$$\text{Var}^* \hat{C}^{\text{is}} = \frac{1}{M} \text{Var}^* \frac{h(X)f(X)}{g(X)} = 0.$$

## Variance reduction

- ▶ However, note that the unique constant such that

$$g(x) = \text{constant} \times h(x)f(x)$$

defines a legitimate density is  $1 / \int h(x)f(x)dx = 1/C$ .

If we knew the value of  $C$ , we wouldn't need Monte Carlo. So in practice, this recipe for choosing  $g$  cannot be followed exactly.

- ▶ What we *can* do is choose  $g$  to be a known density *approximately* proportional to the original density  $f$  times the payoff  $h$ .


Irrespective of whether this approximation is bad or good, we still have an unbiased estimate  $\hat{C}^{\text{is}}$ .

A good approximation can make the variance of  $\hat{C}^{\text{is}}$  small.

## Variance reduction

If  $X = (X_1, \dots, X_N)$  are Brownian increments driving stock  $S(X)$ , where  $S : \mathbb{R}^N \rightarrow \mathbb{R}$ , then can apply this idea to  $S(X)$  instead of  $X$ .

In other words, if  $h(x) = H(S(x))$ , then variance is zero if  $S(X)$  has IS-density = constant  $\times$   $H$   $\times$  original density of  $S(X)$ .



## Likelihood ratio computations

To compute  $f(X)/g(X)$ , usually you won't be specifying  $g$ . Rather you'd specify, for instance, a desired mean or drift, then compute  $g$ . Compute  $g$  by starting in  $X$  space.

Suppose, in particular, that the components of  $X \in \mathbb{R}^N$  under the original measure are independent with densities  $f_1, f_2, \dots, f_N$ , and that under importance sampling measure (not necessarily optimal), they are independent with densities  $g_1, g_2, \dots, g_N$ . Then

$$\frac{f(x)}{g(x)} = \frac{f(x_1, \dots, x_N)}{g(x_1, \dots, x_N)} = \prod_{n=1}^N \frac{f_n(x_n)}{g_n(x_n)}$$

(Actually, independence is unnecessary, if you let  $f_n$  be the *conditional* density of  $X_n$  given  $X_1, \dots, X_{n-1}$ , and similarly for  $g_n$ .)



## Example: Changing of drift of BM

Let  $X_1, \dots, X_N$  be successive increments of BM  $W$  sampled at times  $0, t_1, \dots, t_N = T$ . Then density  $f_n$  is Normal( $0, \Delta t$ ), where  $\Delta t = T/N$ .

$$f_n(x_n) = \frac{1}{\sqrt{2\pi\Delta t}} e^{-x_n^2/(2\Delta t)}$$

The  $g_n$  can be the density of any other distribution supported on  $\mathbb{R}$ .

In particular, if we take  $g_n$  to be the Normal( $\lambda\Delta t, \Delta t$ ) density, then

$$g_n(x_n) = \frac{1}{\sqrt{2\pi\Delta t}} e^{-(x_n - \lambda\Delta t)^2/(2\Delta t)} \quad \text{Drift of } \lambda \text{ * } \Delta t$$

So for all  $x = (x_1, \dots, x_n)$ ,

$$\frac{f_n(x_n)}{g_n(x_n)} = e^{\frac{1}{2\Delta t}(-x_n^2 + x_n^2 - 2x_n\lambda\Delta t + \lambda^2(\Delta t)^2)} = e^{-\lambda x_n + \frac{1}{2}\lambda^2\Delta t}$$

$$\frac{f(x)}{g(x)} = \prod_{n=1}^N \frac{f_n(x_n)}{g_n(x_n)} \implies \frac{f(X)}{g(X)} = e^{-\lambda \sum X_n + \frac{1}{2}\lambda^2 T} = e^{-\lambda W_T + \frac{1}{2}\lambda^2 T}$$

## Example: Changing the drift of BM

### Implementation

- ▶ If you want to change the drift of  $W$  to  $\lambda$ , then work under  $\mathbb{P}^*$  such that  $W_t^* := W_t - \lambda t$  is  $\mathbb{P}^*$ -Brownian Motion. Then

$$f(X)/g(X) = e^{-\lambda \sum X_n + \frac{1}{2}\lambda^2 T} = e^{-\lambda W_T + \frac{1}{2}\lambda^2 T} = e^{-\lambda W_T^* - \frac{1}{2}\lambda^2 T}$$

Multiply the payoff  $h(X)$  by this.

- ▶ When simulating under the measure  $\mathbb{P}^*$ , the zero-mean normals that you generate are simulations of  $W^*$ , not  $W$ .
- ▶ The  $f(X)/g(X)$  is also known as the *Radon-Nikodym derivative* of  $\mathbb{P}$  with respect to  $\mathbb{P}^*$ .

## Example: Changing the drift of BM

Choice of  $\lambda$

- ▶ How to choose  $\lambda$ ? Maybe too difficult to make  $S_T$ 's distribution agree entirely with the optimal distribution. So let's just make  $S_T$ 's *mean*  $\approx$  the optimal distribution's *mean*. If  $S$  is GBM,

$$dS_t = rS_t dt + \sigma S_t dW_t = (r + \sigma\lambda)S_t dt + \sigma S_t dW_t^*$$

So  $\mathbb{E}^* S_T = S_0 e^{(r+\sigma\lambda)T}$ . You could, for instance, try to choose  $\lambda$  such that  $S_0 e^{(r+\sigma\lambda)T} \approx$  the optimal distribution's mean.

- ▶ Even if  $\lambda$  is not chosen optimally, the importance sampling MC estimate is still unbiased.

Variance reduction: Antithetic variates

Variance reduction: Control variates

Variance reduction: Importance sampling

Variance reduction: Conditional MC

Look at annotated version

## Conditional Monte Carlo

No need to simulate green and blue lines

Just analytically compute  $X$  and use a simulation for lines for  $X$

For any random variable  $X$ , by the law of iterated expectations,

$$C = \mathbb{E}Y = \mathbb{E}[\mathbb{E}(Y|X)] = \mathbb{E}f(X)$$

Assuming that the function  $f(X) := \mathbb{E}(Y|X)$  can be computed analytically, Conditional Monte Carlo does the following:

- ▶ Simulate not  $Y$  but rather  $X$ , generating  $X_1, \dots, X_M$  IID as  $X$ .
- ▶ Estimate  $\mathbb{E}[\mathbb{E}(Y|X)]$  by taking the average

$$\hat{C}_M^{\text{cmc}} := \frac{1}{M} \sum_m f(X_m).$$

The standard error is  $\hat{\sigma}_M^{\text{cmc}}/\sqrt{M}$  where

$$\hat{\sigma}_M^{\text{cmc}} := \sqrt{\frac{1}{M-1} \sum_m (f(X_m) - \hat{C}_M^{\text{cmc}})^2}$$

# Conditional Monte Carlo: variance analysis

Expectation: The CMC estimate is unbiased, because

$$\mathbb{E}\hat{C}_M^{\text{cmc}} = \mathbb{E}f(X) = C.$$

Variance: CMC has smaller variance than ordinary MC because

$$\text{Var}(\hat{C}_M^{\text{cmc}}) = \frac{1}{M} \text{Var}f(X) = \frac{1}{M} \text{Var}[\mathbb{E}(Y|X)] \leq \frac{1}{M} \text{Var}(Y) = \text{Var}(\hat{C})$$

where the last step is because  $\text{Var}(Y) = \text{Var}[\mathbb{E}(Y|X)] + \mathbb{E}\text{Var}(Y|X)$ .

Intuition: suppose  $Y = F(\mathbf{X})$  where  $\mathbf{X} \in \mathbb{R}^D$ . Then ordinary MC computes a  $D$ -dimensional integral  $\int F(\mathbf{x})p(\mathbf{x})d\mathbf{x}$  by drawing random  $\mathbf{X}_m \in \mathbb{R}^D$  with density  $p$ . But if, along  $d$  of those dimensions, the integration can be performed *analytically*, then we only need to draw random vectors  $X_m \in \mathbb{R}^{D-d}$ . This reduces the noise in the estimate.

# Conditional Monte Carlo: example 1

Each  $S_t$  is log normal, but their sum is not log normal

Let  $0 < t_1 < t_2 < t_3$ . Consider at time 0 an Asian-style call on a GBM  $S$ , with expiry  $t_3$  and discounted payoff

$$Y := e^{-rt_3} \left( \frac{S_{t_1} + S_{t_2} + S_{t_3}}{3} - k \right)^+$$

Let  $X = (S_{t_1}, S_{t_2})$ . Then  $\mathbb{E}(Y|X)$  is analytically known, because

$$Y = \frac{1}{3} e^{-rt_3} \left( S_{t_3} - (3k - S_{t_1} - S_{t_2}) \right)^+$$

We will simulate for time1 and time2, but not for time3

So  $\mathbb{E}(Y|X)$  can be analytically evaluated using the B-S formula.

CMC: Generate  $M$  realizations of  $(S_{t_1}, S_{t_2})$  and take the average of

Take 1,000 realizations of time1 Stock and time2 Stock, then compute  $f(S_{t_1}, S_{t_2})$

$$\mathbb{E}(Y|X) = \frac{1}{3} e^{-rt_2} C^{BS}(S = S_{t_2}, t = t_2, K = 3k - (S_{t_1} + S_{t_2}), T = t_3).$$

Use time3 as expiry

This random variable has *smaller variance* than  $Y$ .

## Conditional Monte Carlo: example 2a

Consider a call with discounted payoff  $Y = e^{-rT}(S_T - K)^+$  where

$$dS_t = rS_t dt + \sqrt{V_t}S_t dW_{1t}$$

$$dV_t = \alpha(V_t)dt + \beta(V_t)dW_{2t}$$

and  $W_1$  and  $W_2$  are (in this particular example) *independent* BM.

Note that *conditional on the entire path of  $V_t$* , the dynamics of  $S$  are

$dS_t = rS_t dt + \sqrt{V_t}S_t dW_{1t}$  where the  $V_t$  path is conditionally *fixed*,

and  $W_1$  is still conditionally a BM. So  $Y$  has conditional expectation

$$\mathbb{E}(Y|X) = C^{BS}(\bar{\sigma}) \quad \text{where } \bar{\sigma}^2 := \frac{1}{T} \int_0^T V_t dt$$

instantaneous variance

and so  $Y$  has unconditional expectation given by the “mixing formula”

$$C = \mathbb{E}C^{BS}(\bar{\sigma}).$$



## Conditional Monte Carlo: example 2a

Two approaches:

► Ordinary MC:

Generate  $M$  realizations of the two-dimensional path of  $(S, V)$ .

On each path calculate the discounted call payoff. Average.

► Conditional MC:

Generate  $M$  realizations of the one-dimensional path of  $V$ .

On each path calculate  $C^{BS}(\bar{\sigma})$ . Average.

CMC reduces variance by removing from the simulations the noise due to the randomness of  $S$ .

## Conditional Monte Carlo: example 2b

Stochastic volatility model with nonzero correlation: Let

$$W_t = \sqrt{1 - \rho^2} dW_{1t} + \rho dW_{2t}$$

where  $W_1$  and  $W_2$  are independent BMs, so  $W$  and  $W_2$  have correlation  $\rho$ . Let

$$dS_t = rS_t dt + \sigma_t S_t dW_t$$

$$\sigma_t = \sqrt{V_t}$$

$$dV_t = \alpha(V_t)dt + \beta(V_t)dW_{2t},$$

## Conditional Monte Carlo: example 2b

Then  $L := \log S$  satisfies

$$\begin{aligned} dL_t &= rdt - \frac{1}{2}\sigma_t^2 dt + \sigma_t \sqrt{1 - \rho^2} dW_{1t} + \sigma_t \rho dW_{2t} \\ &= rdt - \frac{1 - \rho^2}{2} \sigma_t^2 dt + \sigma_t \sqrt{1 - \rho^2} dW_{1t} - \frac{\rho^2}{2} \sigma_t^2 dt + \sigma_t \rho dW_{2t} \end{aligned}$$

So conditional on  $\mathcal{F}_T^{W_2}$ ,

$$L_T \sim \text{Normal} \left( L_0 + rT + \log M_T(\rho) - \frac{1 - \rho^2}{2} \bar{\sigma}^2 T, (1 - \rho^2) \bar{\sigma}^2 T \right)$$

where

$$M_T(\rho) := \exp \left( -\frac{\rho^2}{2} \int_0^T \sigma_t^2 dt + \rho \int_0^T \sigma_t dW_{2t} \right)$$

So even with correlation,  $S$  is still  $W_2$ -conditionally lognormal.

SKIP

## Conditional Monte Carlo: example 2b

“Mixing formula” for price of option paying  $C_T = C(S_T)$

$$C_0 = \mathbb{E}(e^{-rT} C_T) = \mathbb{E}(\mathbb{E}(e^{-rT} C_T | \mathcal{F}_T^{W_2})) = \mathbb{E} C^{BS}(S_0 M_T(\rho), \bar{\sigma} \sqrt{1 - \rho^2})$$

where

$$M_T(\rho) := \exp \left( -\frac{\rho^2}{2} \int_0^T \sigma_t^2 dt + \rho \int_0^T \sigma_t dW_{2t} \right)$$

Option price is expectation of the B-S formula for that option, evaluated at random volatility *and* a randomized spot.

Therefore:

Simulate  $W_2$ . Given  $W_2$ , evaluate  $C^{BS}(S_0 M_T, \bar{\sigma} \sqrt{1 - \rho^2})$ .

Average across all simulations of  $W_2$ . No need to simulate  $W_1$ .