

A new conceptual typology for the population impacts of sea level rise *

Mathew E. Hauer^{1*} *University of Georgia*
R. Dean Hardy² *University of Maryland*

This paper provides a conceptual framework for assessments of the impacts of sea level rise on populations. Population impact assessments of sea level rise are common in the literature and such assessments directly influence our understanding of the societal effects of climate change and sea level rise. However, these assessments typically limit population exposure to inundation areas only. Drawing on increasingly sophisticated flood modeling, we propose a new typology for understanding population exposure risk to sea level rise based on a spatial envelope probability of exposure to tidal flooding approach. The typology identifies three types of exposure risk: direct, indirect, and tertiary. Implications and effects of sea level rise on each exposure risk typology is then discussed. Using Chatham County, Georgia as an example, we find that current assessments could grossly underestimate both the spatial and temporal dimensions of the societal impacts of sea level rise. The typology can be used to guide new research, assist with impact assessments, evaluate policy options, and provide more robust and holistic scenarios of sea level rise.

- Corresponding author. hauer@uga.edu. p: 706-542-9369.

¹ Carl Vinson Institute of Government, University of Georgia. 201 N. Milledge Ave. Athens, GA USA 30602.

² National Socio-Environmental Synthesis Center, University of Maryland College Park. 1 Park Place, Annapolis MD USA 21401.

Introduction

The societal impacts of climate change and sea level rise (SLR) are of utmost importance in the 21st century and climate assessments are an integral part of adaptation and risk management ([Howden and Jacobs, 2016](#)). Research has long sought to identify the impact of sea level rise on society ([Curtis and Schneider, 2011](#); [Wu, Yarnal and Fisher, 2002](#)), with an increasing focus on estimation of at-risk populations associated with sea level rise ([Curtis and](#)

*All data and code that supports these conclusions are available as supplementary materials.

[Schneider, 2011](#); [Strauss, Kulp and Levermann, 2015](#)). With 40% of the US population living in a NOAA designated coastal community ([Ache et al., 2015](#)) and over one billion people living in coastal areas around the world ([Neumann, Vafeidis, Zimmermann and Nicholls, 2015](#)), questions on the societal impacts of sea level rise continue to be asked.

One of the primary components of a climate assessments of the impacts of sea level rise involves calculating at-risk populations. A wide variety of definitions of ‘at-risk’ populations have been utilized including the populations in lower-elevation coastal zones ([Neumann, Vafeidis, Zimmermann and Nicholls, 2015](#); [McGranahan, Balk and Anderson, 2007](#)), populations under the mean higher high water (MHHW) mark ([Hauer, Evans and Mishra, 2016](#); [Strauss, Kulp and Levermann, 2015](#)), and populations in the 100-year flood plain ([Nicholls and Cazenave, 2010](#); [Brown et al., N.d.](#); [Hallegatte et al., 2011](#); [Heberger et al., 2011](#)). All three approaches conceptualize at-risk populations as binary outcomes, ie populations are either at-risk or not at-risk, located within specific geographies defined through each approaches’ characterization schema and share shortcomings related to an equality of risk, an equality of temporal exposure, and an equality of exposure. First, these approaches presume an equality of risk within each geography assuming a homogenous risk profile across heterogeneous terrain. For instance, in a flood plain based approach, all populations in the 100-year flood plain are implicitly assumed to have equal flood risk, despite certain populations living in the 10-year or 20-year flood plain. Second, these approaches presume an equality of temporal exposure. For populations living within 0.9m of the current MHHW mark, those living within 0.1-0.2m of the MHHW mark will surely experience effects of sea level rise sooner than those living in the 0.8-0.9m mark. And lastly, these approaches presume an equality of exposure. For populations living in a lower-elevation coastal zone, some will be exposed to sea level rise and others won’t be exposed to sea level rise but could be exposed to other associated hazards such as storm surges and other livelihood impacts. These shortcomings combine to create a gross oversimplification of the extent, timing, exposure, and ultimately adaptation options for coastal communities.

In this paper, we compare and contrast these three theoretical, binary approaches and suggest a unified typology for investigating exposure to sea level rise using a probability spatial envelope typology. We characterize exposed populations into three main categories: directly impacted, indirectly impacted, and tertiary impacted. These categories combine the strengths of all three previous approaches into a single unified framework. We then examine the implications for this novel framework within the context of Chatham County, Georgia.

Conceptual Models in the

The way populations exposed or at-risk to sea level rise are conceptualized drives the societal impacts from SLR. Assessments using lower-elevation coastal zones, for instance, use generalized risk allocations that can frame climate adaptation discussions across the widest possible area. It is not just simply SLR, but rather a combination of SLR, recurrent tidal flooding, and elevated risk of disastrous storm surges that comprise the full range of effects associated with sea level rise. However, more explicit spatiotemporal linkages to localized SLR hazards is clearly warranted. Adaptation strategies for those currently living under 1.8m of elevation, the “high” scenario of sea level rise expected in the 21st century ([Sweet et al., 2017](#)), are inherently different from those living above 1.8m of elevation but still within the LECZ. Appropriate accounting of type of exposure to the hazards of sea level rise would improve upon our current frameworks.

Recent tidal events in the Atlantic Coast of the United States and elsewhere ([Carbognin et al., 2010](#); [Spanger-Siegfried, Fitzpatrick and Dahl, 2014](#); [Dahl, Fitzpatrick and Spanger-Siegfried, 2017](#)) have demonstrated that the impacts of rising sea levels are advancing further and faster than those indicated by inundation models that assess impacts at mean higher high water (MHHW). The increased frequency and magnitude of tidal flooding in coastal communities can be considered as the immediate impact of sea level rise (SLR). Perigean spring tide events cause regularly recurring water levels well above MHHW, and in many parts of the coast this causes significant flooding, sometimes referred to as nuisance flooding

or recurring tidal flooding ([Fennessey et al., 2001](#)), particularly when these tidal events are supported by significant onshore winds from tropical cyclones or storms. Mean higher high water does not mark the highest extent of regularly occurring tide waters as the tide range will naturally exceed this level approximately half of the time, as its name suggests. Assessments using impacts from changes in MHHW inadvertently turn a blind eye to the potential impacts of sea level rise on areas above the MHHW mark. However, property just above the MHHW mark will be regularly flooded long before it transitions below the MHHW mark, thus individuals and properties would not be counted as impacted, despite being regularly flooded by tides that exceed MHHW with added SLR. This is especially important in areas with large variations in their tidal datums.

A more broad approach involves utilizing either lower-elevation coastal zones or the 100-year flood plain to characterize the populations exposed to hazards from sea level rise ([Hallegatte et al., 2011](#); [Neumann, Vafeidis, Zimmermann and Nicholls, 2015](#)). In this way, populations within these geographies are framed as having equal risk regardless of their location within the LECZ or the 100-year flood plain. This approach has a side effect of equalizing the temporal horizon of hazards associated with sea level rise. By characterizing all populations in the LECZ or the 100-year flood plain as being at-risk to the hazards associated with sea level rise, regardless of either the extent of sea level rise or the timing of the rise. In other words, both lower elevation coastal zones and 100-year flood plain approaches render any coastal population as exposed to sea level rise in any time period. This has an unintended consequence of inflating sea level rise hazards as all coastal populations are effectively “exposed” to sea level rise and makes it difficult to decompose the effect of sea level rise on typical coastal hazards. Shoreline populations in areas with tropical cyclone activity are already exposed to storm surges, for instance, regardless of sea level rise. Which additional populations will be exposed to storm surges due to a higher base water level becomes increasingly difficult with a LECZ or 100-year flood plain approach.




Figure 1: **SLR typology outline.** The vertical dashed line represents 0.5m of SLR, the solid vertical line represents the exposed population using a floodplain approach.

Figure 1 summarize the current conceptual models for examining the impacts of sea level rise. Panels a, b, and c in Figure 1 demonstrate conceptually who is at risk of SLR assuming 0.5m of SLR. Using a LECZ conceptualization (a), all persons within the LECZ are assumed to be exposed to SLR. Using the 1 in 100 year floodplain conceptualization (b), all persons below a given elevation are assumed to be at risk despite reduced flood risks at higher elevations. Using a MHHW conceptualization (c), only those persons who have exactly 100% probability of flooding are considered at risk.

A Unified Conceptual Model

When taken together, these three models capture all possible hazards associated with sea level rise but are plagued by issues on an individual conceptual basis. We propose a new model, unifying the MHHW, LECZ, and 100-year floodplain approaches into a more conceptually sound framework based on a spatial envelope, probability of exposure model. In a departure from the other three models' characterizations of at-risk and not at-risk, we characterize exposure to sea level rise as **inundated**, **direct**, and **indirect** based on a location's probability of flooding within spatial envelopes. A three tiered exposure approach allows for a pliable examination of both hazard exposure and adaptive options.

We characterize those who are **inundated** as those people living below the future MHHW mark. These are the populations who will be the most adversely affected, will experience impacts from sea level rise the soonest, and who are directly threatened by inundation if adaptive measures are not undertaken. Inundation exposure is analogous to the population's risk understood through the MHHW approach discussed above. Eventually, these populations will experience water levels above their elevation 50%+ of the time. These areas are typically discussed in conversations concerning managed retreats ([Huntington, Goodstein and Euskirchen, 2012](#); [Hino, Field and Mach, 2017](#)). Such areas will be directly affected as soon as 2045 under some of the most aggressive sea level rise curves ([DeConto and Pollard, 2016](#); [@ Sweet et al., 2017](#)).

We characterize those who are **directly** exposed to sea level rise as those people who are living above the future MHHW mark, but below the spatial extent of future flooding from sea level extremes or tidal flooding. These are the populations exposed to recurrent or nuisance flooding (Chang et al., 2010; Dahl, Fitzpatrick and Spanger-Siegfried, 2017) and will experience water levels above their elevation less than 50% of the time. These areas are typically not discussed in the context of managed retreats, but are frequently discussed regarding adaptive measures such as elevating roads (Titus et al., 2009), coastal armoring (Jin et al., 2015), or other near-term adaptations.

Finally, we characterize those who are **indirectly** exposed to sea level rise as living in areas above both the MHHW mark and the extent of future tidal flooding, but in neighborhoods that are directly or semi-directly at risk to sea level rise. Here, we conceptualize the rest of the areas located in a LECZ or 100-year floodplain or other coastal geography. While these areas are not likely to see their properties exposed to elevated sea levels, the people in these areas will likely drive on flooded roads, go to workplaces in flooded areas, and could see their property values depressed (Neumann, Emanuel, Ravela, Ludwig, Kirshen, Bosma and Martinich, 2015).

Our framework bases SLR exposure on both flood exposure, expressed here as probability of flooding, and geographic location. We can also see the presence of indirectly impacted populations under current conditions (\autoref{figure1}). Sea level rise is typically imagined as a hazard typified by long range time horizons up to 2,000 years into the future (Strauss, Kulp and Levermann, 2015), but related impacts from sea level rise, specifically manifested as coastal flooding, are occurring today in parts of the Atlantic Coast of the United States and Venice, Italy, for instance.

To demonstrate the differences in impacts under our proposed model compared to previous frameworks, we use an empirical example of Chatham County, Georgia (Figure 2). Chatham County is an ideal platform for demonstrating the various populations exposed to sea level rise in our conceptual model. It is a coastal county with a Census 2010 population

Table 1: My caption

| | |
|------------|--|
| Inundated | $PR_j^s = P_j \cdot \frac{A_j^s}{A_j^0}$ |
| Directly | $PR_j^s = P_j \cdot \frac{A_j^{s+}}{A_j^0}$ |
| Indirectly | $\frac{A_j^{s+}}{A_j^0} > 0, \text{ Indirectly} = P_j$ |

of approximately 266,000, making it a medium-sized area, and it is typified by its coastal marsh ecosystem and very large astronomical tides. It has also been identified as having approximately 10% of its population at risk to sea level rise using a MHHW approach ([Hauer, Evans and Mishra, 2016](#)).

We use the National Oceanographic and Atmospheric Administration’s (NOAA) sea level rise database that simulate expected changes in the mean higher high water (MHHW) mark on areas that are hydrologically connected to coastal areas for the 0ft through 5ft datasets. These datasets does not take into account additional land loss caused by other natural factors such as erosion, subsidence, or future construction and NOAA provides these data “as is” without warranty to their performance. Land lost due to sea level rise is calculated with a spatial overlay workflow in ArcGIS 10.1 as one minus the percentage of land lost under the 0ft base layer of sea level rise, ie 1ft divided by 0ft, 2ft divided by 0ft, etc. The first step in the analysis was to utilize a base, 0ft Mean Higher High Water (MHHW) layer, which was derived from NOAA’s 0ft scenario, and used as the initial condition to calculate a base of dry land area contained within the geographies of 2010 Census Block Groups. The resulting calculation is therefore a total area of dry land currently available for human habitation within each Census Block Group geography.

We used an area based approach ([Hauer, Evans and Mishra, 2016](#)) to classify populations as inundated, directly, or indirectly impacted using Census Block Group geographies through the following sets of equations in Table X. All populations reported come from Census 2010.

Where the population at risk of being impacted (PR_j^s) under scenario s in census block group j is equal to the population (P) in census block group j multiplied by the ratio of dry




Figure 2: **Proposed SLR typology effect in Chatham County, GA** The vertical dashed line represents 0.5m of SLR, the solid vertical line represents the exposed population using a floodplain approach.

land area (A_j^s) in census block group j under sea level rise scenario s to the dry land area under 0ft of sea level rise (A_j^0). For indirectly and tertiary impacted areas, we use $s+$ to denote the dry land area under sea level rise scenario s plus the highest astronomical tide. For Chatham County, this is roughly equivalent to approximately 2ft above the MHHW mark.

Figure 2 shows the results for all three at-risk classifications under current conditions (0ft) through 3ft of SLR. We can see that the narrowest interpretation of exposure to sea level rise (**inundated**), roughly corresponds to those who will be inundated by MHHW, represents the smallest exposed population. Approximately 12,000 people could be directly exposed with 1ft of SLR, growing to approximately 27,000 with 3ft of SLR. Those who could

be exposed to coastal flooding, or **directly exposed**, currently sits at approximately 21,000 people under a 0ft, baseline sea level rise scenario representing current conditions. This nearly doubles to approximately 43,000 people with 3ft of SLR. Lastly, we see that those who live in neighborhoods that will experience flooding is currently approximately 63,000 but grows to nearly 140,000 with just 3ft of SLR.

References

- Ache, Brent W, Kristen M Crossett, Percy A Pacheco, Jeffery E Adkins and Peter C Wiley. 2015. "The coast" is complicated: a model to consistently describe the nation's coastal population." *Estuaries and coasts* 38(1):151–155.
- Brown, Sally, RJ Nicholls, Philip Goodwin, ID Haigh, Daniel Lincke, AT Vafeidis and Jochen Hinkel. N.d. "Quantifying Land and People Exposed to Sea-Level Rise with No Mitigation and 1.5 and 2.0 C Rise in Global Temperatures to Year 2300." *Earth's Future*. Forthcoming.
- Carbognin, Laura, Pietro Teatini, Alberto Tomasin and Luigi Tosi. 2010. "Global change and relative sea level rise at Venice: what impact in term of flooding." *Climate Dynamics* 35(6):1039–1047.
- Chang, Heejun, Martin Lafrenz, Il-Won Jung, Miguel Figliozzi, Deena Platman and Cindy Pederson. 2010. "Potential impacts of climate change on flood-induced travel disruptions: a case study of Portland, Oregon, USA." *Annals of the Association of American Geographers* 100(4):938–952.
- Curtis, Katherine J and Annemarie Schneider. 2011. "Understanding the demographic implications of climate change: estimates of localized population predictions under future scenarios of sea-level rise." *Population and Environment* 33(1):28–54.
- Dahl, Kristina A, Melanie F Fitzpatrick and Erika Spanger-Siegfried. 2017. "Sea level rise drives increased tidal flooding frequency at tide gauges along the US East and Gulf Coasts: Projections for 2030 and 2045." *PloS one* 12(2):e0170949.
- DeConto, Robert M and David Pollard. 2016. "Contribution of Antarctica to past and future sea-level rise." *Nature* 531(7596):591.
- Fennessey, Lawrence AJ, James M Hamlett, Gert Aron and David LaSota. 2001. "Changes in runoff due to stormwater management pond regulations." *Journal of Hydrologic Engineering* 6(4):317–327.
- Hallegatte, Stephane, Nicola Ranger, Olivier Mestre, Patrice Dumas, Jan Corfee-Morlot, Celine Herweijer and Robert Muir Wood. 2011. "Assessing climate change impacts, sea level rise and storm surge risk in port cities: a case study on Copenhagen." *Climatic change* 104(1):113–137.
- Hauer, Mathew E, Jason M Evans and Deepak R Mishra. 2016. "Millions projected to be at risk from sea-level rise in the continental United States." *Nature Climate Change* 6(7):691.
- Heberger, Matthew, Heather Cooley, Pablo Herrera, Peter H Gleick and Eli Moore. 2011. "Potential impacts of increased coastal flooding in California due to sea-level rise." *Climatic Change* 109(1):229–249.
- Hino, Miyuki, Christopher B Field and Katharine J Mach. 2017. "Managed retreat as a response to natural hazard risk." *Nature Climate Change* 7(5):364.

- Howden, Mark and Katharine L Jacobs. 2016. "Innovations in assessment and adaptation: building on the US National Climate Assessment." *Climatic change* 135(1):157–171.
- Huntington, Henry P, Eban Goodstein and Eugénie Euskirchen. 2012. "Towards a tipping point in responding to change: rising costs, fewer options for Arctic and global societies." *Ambio* 41(1):66–74.
- Jin, Di, Porter Hoagland, Donna K Au and Jun Qiu. 2015. "Shoreline change, seawalls, and coastal property values." *Ocean & Coastal Management* 114:185–193.
- McGranahan, Gordon, Deborah Balk and Bridget Anderson. 2007. "The rising tide: assessing the risks of climate change and human settlements in low elevation coastal zones." *Environment and urbanization* 19(1):17–37.
- Neumann, Barbara, Athanasios T Vafeidis, Juliane Zimmermann and Robert J Nicholls. 2015. "Future coastal population growth and exposure to sea-level rise and coastal flooding-a global assessment." *PloS one* 10(3):e0118571.
- Neumann, James E, Kerry Emanuel, Sai Ravela, Lindsay Ludwig, Paul Kirshen, Kirk Bosma and Jeremy Martinich. 2015. "Joint effects of storm surge and sea-level rise on US Coasts: new economic estimates of impacts, adaptation, and benefits of mitigation policy." *Climatic Change* 129(1-2):337–349.
- Nicholls, Robert J and Anny Cazenave. 2010. "Sea-level rise and its impact on coastal zones." *science* 328(5985):1517–1520.
- Spanger-Siegfried, Erika, Melanie Fitzpatrick and Kristina Dahl. 2014. "Encroaching tides: How sea level rise and tidal flooding threaten US East and Gulf Coast communities over the next 30 years."
- Strauss, Benjamin H, Scott Kulp and Anders Levermann. 2015. "Carbon choices determine US cities committed to futures below sea level." *Proceedings of the National Academy of Sciences* 112(44):13508–13513.
- Sweet, William V, Robert E Kopp, Christopher P Weaver, Jayantha Obeysekera, Radley M Horton, E Robert Thieler and Chris Zervas. 2017. "Global and regional sea level rise scenarios for the United States."
- Titus, James G, Daniel E Hudgens, Daniel L Trescott, Michael Craghan, William H Nuckols, Carl H Hershner, JM Kassakian, Chris J Linn, Peter G Merritt, TM McCue et al. 2009. "State and local governments plan for development of most land vulnerable to rising sea level along the US Atlantic coast." *Environmental Research Letters* 4(4):044008.
- Wu, Shuang-Ye, Brent Yarnal and Ann Fisher. 2002. "Vulnerability of coastal communities to sea-level rise: a case study of Cape May County, New Jersey, USA." *Climate Research* 22(3):255–270.