

Asymptotic Flatness and BMS

Mathew Calkins

August 28, 2026

Abstract

Notes on asymptotic flatness and the BMS group.

Contents

1 Geometric preliminaries	1
2 Bondi Gauge	2
3 Asymptotic Flatness from Fall-Off	4

1 Geometric preliminaries

Let M be a four-dimensional spacetime, *i.e.*, a real smooth 4-manifold endowed with a metric g metric signature $(-, +, +, +)$, plus a smoothly varying choice made at every $T_p M$ as to which time-like and null vectors are future-versus past-directed. Any future-directed null geodesic which doesn't eventually get permanently trapped in an isolated system¹ is said to run off to *future null infinity* $\mathcal{I}^+$. *Past null infinity* $\mathcal{I}^-$ is defined analogously. We will see later that conformal compactifications let us more literally interpret $\mathcal{I}^\pm$ as boundary components.

Let $N \subset M$ be a smooth codimension-1 submanifold. For $p \in N$, a vector $n \in T_p N$ is said to be *normal* to N at p if it is orthogonal to all vectors tangent to N at p , *i.e.*, if $\langle n, m \rangle = 0$ for all $m \in T_p N$. Conversely, given such an n the tangent space $T_p N$ is exactly the set of all $m \in T_p M$ such that $\langle m, n \rangle = 0$. Just as in the Euclidean case, at fixed p and N such vectors form a 1-dimensional vector space. By choosing a normalization convention and a local orientation for N , we may therefore speak of *the* normal vector $n = (n^a)$ or the corresponding normal covector $(n_a) \equiv (g_{ab} n^b) = g(n, \cdot)$.

N is called a *null hypersurface* if, at every point $p \in N$, the² normal vector (n^a) is null, or equivalently if the normal covector (n_a) . Then from our observations above we see that $n \in T_p N$, so that n is simultaneously tangent and normal to N at p . Now we abuse notation slightly and extend n to a vector field defined everywhere on N . Then the integral curves of n are all trajectories within N , and cover the latter. We call these null curves the *null generators* of N . If everything is analytically nice enough, then the set of integral curves (equivalently, the set of all equivalence classes $[p \in N]$ where $p \sim q$ iff some integral curve passes through both points) inherits the structure of a real smooth two-manifold. If this occurs, a smooth family of null generators is called a *null congruence*. This smooth structure may hold in some regions but not in others. A *caustic* is a point where neighboring generators intersect or focus so strongly that the inherited smooth structure breaks down.

Suppose we have covered N by such a null congruence of null generators. A *cut* of N is a smooth space-like³ 2-submanifold of N which intersects each null generator exactly once in some region under consideration.

¹This is deliberately loose phrasing which we'll make more precise later

²Here 'the' is a mild abuse of vocabulary since it relies on choices of orientation and normalization, but all such choices agree on whether N is null.

³That is, having exclusively spacelike or vanishing tangent vectors.

Choosing a cut of a null hypersurface should be thought of as the null-geometry analogue of choosing a spatial slice of a Lorentzian manifold.

Why should cuts exist at all? In more algebraic terms and rephrased slightly, why should the induced metric on N have signature $(0, +, +)$? Recall that, algebraically, the signature of a quadratic form $F : V \times V$ over a real vector space of finite dimensions is defined procedurally:

- Pick any basis for F w.r.t. which the component expression for F is diagonal.
- Read off the number of positive, negative, and vanishing diagonal components; this triple is the signature of F .
- Step 1 is provably possible, and the results of Step 2 are provably independent of choices made during Step 1.

So the claim that N has induced metric signature $(0, +, +)$ (or more precisely, that every tangent space $T_p N$ has inner product with this signature) says that there exists a basis of three vectors in $T_p N$ which are pairwise orthogonal and which are respectively null, spacelike, and spacelike. To show this, we take the first of these vectors to be n itself. By definition $\langle n, m \rangle = 0$ for all $m \in T_p N$, so n is null and any completion to a basis (n, k, k') automatically obeys $\langle n, k \rangle = \langle n, k' \rangle = 0$; the only non-automatic constraints remaining are that (n, k, k') be linearly independent (which excludes only a measure-0 set of sets) and that $\langle k, k' \rangle = 0$ (which can be accomplished by subsequently applying a Gram-Schmidt procedure). What, then, can we say for sure about k, k' ? We know that no null A and timelike B obey $\langle A, B \rangle = 0$, so k, k' must both be null or spacelike. We also know that two nonzero null vectors A, B satisfy $\langle A, B \rangle = 0$ iff one is a rescaling of the other, but k, k' are both assumed linearly independent of n . So by elimination k, k' are both spacelike, and the induced metric on N must have signature $(0, +, +)$.

To build a field theory on M , we define a collection of fields to be fibered over M plus prescribed boundary conditions⁴ The *(on- / off-)shell asymptotic phase space* of our theory is the set of all histories obeying the prescribed boundary conditions (and / but not necessarily) the equations of motion.

2 Bondi Gauge

When we must work in coordinates, it will be convenient to locally work in *Bondi gauge*, which is well-adapted to studying null geodesics. Here we develop the Bondi gauge in slightly different language from the standard references [?, ?, ?]. First, take a null hypersurface N and extend it locally to a 1-parameter family of such surfaces $\{N_u\}_u$ parametrized by a real variable u . This can be done in some region about a point of interest, but attempting to extend it to a smooth foliation $M = \cup_u N_u$ may run into geometric obstructions (in the form of, *e.g.*, caustics) or global topological obstructions, so already our gauge choice is at most guaranteed to work only locally. This gives us our first local coordinate u , telling which copy of N a point lives on.

Let n be a null vector field defined near our point of interest, such that at every point $p \in N_u \subset M$ the local value $n(p) \in T_p M$ is normal and tangent to N_u . From n we can construct null generators as described above, so that by construction every point in our region lies on exactly one generator. Now along each N_u pick a cut $\Sigma_u \subset N_u$, chosen such that N_u varies smoothly with u . Pick coordinates $(x^A)_{A=1,2}$ along each cut Σ_u , then extend these to coordinates on N_u by requiring that they be constant along each null generator; we can locally do so in such a way that x^A varies smoothly with u . This gives us coordinates (u, x^A) .

At this point we have by construction that u and x^A are constant along generators, so we take the remaining coordinate r to grow along generators, and to vary smoothly more generally. Since moving along a particular generator changes r while leaving (u, x^A) fixed, we have automatically that ∂_r points along the null generators, so ∂_r is some local rescaling of n .

⁴More precisely [?], in defining a variation problem we should think of spatial and null boundary conditions as defining a theory, but leave future/past boundary conditions undefined as these define states in the theory.

What do these geometric choices tell us about the metric in components (u, r, x^A) ? First, since ∂_r points along the null generators, ∂_r must be null, hence

$$g_{rr} = g(\partial_r, \partial_r) = 0 \quad (1)$$

Next, varying x^A at fixed (u, r) keeps us in the same N_u , so ∂_A is tangent to N_u . But $n = \partial_r$ is normal to N_u , hence by definition orthogonal to every vector tangent to N_u . So we have

$$g_{rA} = g(\partial_r, \partial_A) = 0. \quad (2)$$

Our metric thus takes the form

$$ds^2 = g_{uu}du^2 + 2g_{ur}dudr + 2g_{uA}dudx^A + g_{AB}dx^A dx^B. \quad (3)$$

We write this coordinate system as $(x^a) = (u, r, x^A)$. For convenience we now rename our coefficients. First, from our earlier discussion of cuts, metric signature considerations ensure that (g_{AB}) is strictly positive-definite, hence invertible as an array. We therefore define

$$U^A \equiv g^{AB}g_{uB} \implies g_{uA} = g_{AB}U^B. \quad (4)$$

Next we complete the square and further rename as

$$\begin{aligned} ds^2 &= g_{uu}du^2 + 2g_{ur}dudr + 2g_{AB}U^Bdudx^A + g_{AB}dx^A dx^B \\ &= (g_{uu} - g_{AB}U^A U^B) du^2 + 2g_{ur}dudr + g_{AB}(U^A du + dx^A)(U^B du + dx^B) \\ &\equiv -Fdu^2 - 2e^\beta dudr + g_{AB}(U^A du + dx^A)(U^B du + dx^B). \end{aligned} \quad (5)$$

This is pure notation, except for the choice $g_{ur} \equiv e^\beta$ which implies $\langle \partial_u, \partial_r \rangle < 0$, which in turn can be enforced within remnant coordinate freedom by appropriate choice of the direction in which u or r increases.

If we further choose⁵ the determinant of the induced metric on the cuts to scale like r^4

$$\partial_r \det\left(\frac{g_{AB}}{r^2}\right) = 0 \quad (6)$$

we say that r has been taken to be *areal* or *luminosity* distance. Integrating at fixed (u, r) , this implies that each fixed- r surface (which by similar arguments to those given above, must have purely spatial induced metric) has surface area proportionate to r^2 , if indeed it has finite surface area at all. Then we write $g_{AB} = r^2 h_{AB}$ and

$$ds^2 = -Fdu^2 - 2e^\beta dudr + r^2 h_{AB}(U^A du + dx^A)(U^B du + dx^B). \quad (7)$$

Motivated by (6), we sometimes write the induced spherical metric on the celestial sphere at fixed (u, r) as

$$g_{AB}dx^A dx^B \equiv r^2 q_{AB}dx^A dx^B \implies \partial_r \det(q_{AB}) = 0. \quad (8)$$

Any coordinate system obeying $g_{rA} = g_{rr} = 0$ plus the luminosity distance condition is said to be in *Bondi gauge*; it is left to the reader's geometric imagination that every metric in this gauge can be cooked up by the geometric story we have told here. The archetypal example of a spacetime described in Bondi gauge in $3+1$ Minkowski space where, starting from some inertial coordinates $(t, \vec{x})$, we take $r = |\vec{x}|$ to be spatial distance from the original, $u = t - r$ to be retarded time, and (x^A) to be any convenient set of coordinates on the celestial sphere. For example, taking $(x^A) = (\theta, \phi)$ we have the Minkowski metric in Bondi coordinates

$$\begin{aligned} ds^2 &= -du^2 - 2dudr + r^2(d\theta^2 + \sin^2\theta d\phi^2) \\ &= -ds^2 - 2dudr + 2r^2\gamma_{z\bar{z}}dzd\bar{z}, \quad \gamma_{z\bar{z}} \equiv \frac{2}{(1+z\bar{z})^2} \\ &\equiv g_{ab}^{\text{MB}}dx^a dx^b. \end{aligned} \quad (9)$$

In studying asymptotic symmetries and celestial holography it is frequently useful to replace the angular coordinates (θ, ϕ) with complex coordinates $(z, \bar{z})$ induced by, *e.g.*, stereographic projection as we have just done.

⁵This cannot in general be accomplished by a redefinition of the coordinate r ; it constrains the possibilities for choosing $\{N_u\}$ at the start of this construction. For example, it is possible in flat Minkowski space if we take N_u to be the forward lightcone of the point $(t, x, y, z) = (u, 0, 0, 0)$ or equivalently the locus $t - r = u$ but not if we take N_u to be the locus $t - x = u$.

3 Asymptotic Flatness from Fall-Off

We saw above that we can always fix our coordinates locally so that the conditions (5) and (6) are satisfied. This is completely generic, and doesn't require that our spacetime obey certain geometric properties or be on-shell. One of several inequivalent (but not totally independent) definitions of asymptotic flatness is that, near future null infinity⁶ $\mathcal{I}^+$, there exists some coordinate chart (u, r, x^A) in Bondi gauge such that approaching $\mathcal{I}^+$ is given exactly by taking $r \rightarrow \infty$ at fixed (u, x^A) , and where the components of the metric approach (9) in that limit. Schematically, we have

$$g_{ab}(u, r, z, \bar{z}) = g_{ab}^{\text{MB}}(u, r, z, \bar{z}) + (\text{corrections strictly subleading in } r). \quad (10)$$

There are various inequivalent conventions for the rate at which we demand the corrections decay. In one reasonably well-motivated convention, we have

$$\begin{aligned} g_{rr} &= 0, \\ \partial_r \det(g_{AB}/r^2) &= 0, \\ g_{rA} &= 0, \\ g_{uu} &= -1 + \mathcal{O}(r^{-1}), \\ g_{ur} &= -1 + \mathcal{O}(r^{-2}), \\ g_{uA} &= \mathcal{O}(1), \\ g_{AB} &= r^2 q_{AB}^{\text{MB}} + \mathcal{O}(r). \end{aligned} \quad (11)$$

⁶We haven't carefully defined this, and won't until a later section on conformal compactification. Roughly speaking, $\mathcal{I}^+$ this is the asymptotic spacetime boundary to which future-directed null rays all tend as long as they don't hit a boundary at finite affine parameter or get trapped in some compact spatial region. We assume our spacetime's geometry is such that this characterizes a single connected region. In the present construction we are demanding that there exists a certain nice chart covering the whole of this region.