Multipole Moments

Mathew Calkins

November 2, 2026

Abstract

Extremely rough notes on multipole moments. Maybe these will turn into something coherent, maybe not!

Contents

1 Spherical Harmonics **1**

2 $SO(3)$ Irreps **3**

3 Harmonic Polynomials **4**

4 STF tensors **5**

5 Multipole Moments **6**

6 Shifting the Origin **8**

7 Multipole Moments as Conformal Killing Tensors **9**

8 Conformal Killing Tensors and Trace Subtraction **10**

1 Spherical Harmonics

Spherical harmonics most naturally arise when solving the 3D Euclidean Laplace equation

$$\nabla\phi = 0, \quad \phi : \mathbb{R}^3 \to \mathbb{R} \tag{1}$$

by the ansatz of separation of variables in spherical coordinates

$$\phi(r,\theta,\phi) = R(r)Y(\theta,\phi) \tag{2}$$

whereupon the angular function equivalently obeys

$$-\lambda = \frac{1}{Y}\frac{1}{\sin\theta}\partial_\theta\left(\sin\theta\partial_\theta Y\right) + \frac{1}{Y}\frac{1}{\sin^2\theta}\partial_\phi^2 Y \tag{3}$$

for λ a constant also appearing in the ODE for R. Further separating

$$Y(\theta,\phi) = \Theta(\theta)\Phi(\phi) \tag{4}$$

gives the system

$$\begin{aligned} \Phi'' &= -m^2\Phi, \\ \lambda\sin^2\theta + \frac{\sin\theta}{\Theta}\frac{d}{d\theta}\left(\sin\theta\frac{d\Theta}{d\theta}\right) &= m^2 \end{aligned} \tag{5}$$

for $m \in \mathbb{R}$ an additional constant. The first equation forces $m \in \mathbb{Z}$. The second, plus demanding regularity at the poles $\theta = 0, \pi$, forces

$$\lambda = \ell(\ell+1) \text{ for } \ell \in \mathbb{Z}, \ell \geq |m|. \tag{6}$$

So we have a basis of solutions

$$Y_\ell^m(\theta, \phi) \propto e^{im\phi} P_\ell^m(\cos\theta), \quad m = -\ell, -\ell+1, \ldots, \ell \tag{7}$$

where P_ℓ^m are the associated Legendre polynomials. In that case, for summations over basis solutions converging at large r, the radial function is then

$$R(r) \propto \frac{1}{r^{\ell+1}}. \tag{8}$$

We can say more about the functions Y, namely that they make up exactly the complete set of eigenfunctions of the spherical Laplacian

$$r^2 \nabla^2 Y_\ell^m = -\ell(\ell+1) Y_\ell^m. \tag{9}$$

This notation is given in a slightly ambiguous way in some presentation, leaving it unclear where we mean the 3D Euclidean Laplacian applied to a function which just happens to be r-independent, or the Laplacian intrinsic to S^2 applied to a map $S^2 \to \mathbb{R}$? As it turns out, these agree even in higher dimensions. To see this, take the usual curvilinear expression for the Laplacian of a scalar

$$\nabla^2 f = \frac{1}{\sqrt{g}} \partial_a \left(\sqrt{g} g^{ab} \partial_b f \right) \tag{10}$$

and apply this to the Euclidean metric for $\mathbb{R}^N$ written in generalized spherical coordinates

$$ds^2 = dr^2 + r^2 d\Omega^2 \implies \sqrt{g} = r^{N-1} \tag{11}$$

Considering the cases where a, b are both angles, both radial, or a mixture, we find that

$$\nabla^2_{\mathbb{R}^N} f = \frac{1}{r^{N-1}} \frac{\partial}{\partial r} \left(r^{N-1} \frac{\partial f}{\partial r} \right) + \frac{1}{r^2} \Delta^2_{S^{N-1}} f. \tag{12}$$

So a function $f : S^2 \to \mathbb{R}$ is annihilated by the sphere Laplacian iff, when interpreted as a radius-independent function on $\mathbb{R}^N$, it is annihilated by the Euclidean Laplacian.

We have just diagonalized the linear operator[1]

$$\nabla^2_{\mathbb{R}^3} : C^\infty(S^2, \mathbb{C}) \to C^\infty(S^2, \mathbb{C}). \tag{13}$$

If we interpret its domain and codomain as linear spaces endowed with the unique-up-to-scaling rotation-invariant inner product

$$\langle f, g \rangle \propto \int_{S^2} d\Omega f(\bar{\Omega}) g(\Omega) \tag{14}$$

then $\nabla^2_{\mathbb{R}^3}$ becomes a Hermitian operator, so its eigenfunctions form a complete orthonormal basis. In fact, since we can take infinite sums over this space of differentiable functions and get non-differentiable limits, they span the larger space

$$L^2(S^2, \mathbb{C}) = \text{Span}_{\mathbb{C}} \left(Y_\ell^m \right)_{|m| \leq \ell}. \tag{15}$$

Again, refer to a real math text to get the careful version.

[1] As usual in theoretical physics, here we are assuming as much smoothness from our test functions as we could ever plausibly need. The domains and codomains here are kind of cartoon placeholders, and a real treatment of representation theory will give the right ones.

2 $SO(3)$ Irreps

Recall that $SO(3)$ has a distinct[2] complex irrep V_ℓ of every odd dimension[3] $2\ell+1$ for $\ell \in \mathbb{Z}_{\geq 0}$. These are generated by the Lie algebra basis elements which, in one standard choice, have matrix elements

$$\begin{aligned}
[L_x^{(\ell)}]_{ij} &= \frac{1}{2}(\delta_{i,j+1} + \delta_{i+1,j}), \sqrt{(\ell+1)(i+j-1) - ij} \\
[L_y^{(\ell)}]_{ij} &= \frac{1}{2i}(\delta_{i,j+1} - \delta_{i+1,j})\sqrt{(\ell+1)(i+j-1) - ij}, \\
[L_z^{(\ell)}]_{ij} &= (\ell+1-j)\delta_{ij}.
\end{aligned} \tag{16}$$

Each irrep is labeled by the eigenvalues of the unique casimir

$$\vec{L}^2 \equiv L_x^2 + L_y^2 + L_z^2 = -\ell(\ell+1)\mathbf{1}_{2\ell+1} \tag{17}$$

with degeneracy within each irrep broken by (in the usual choice) L_z, having eigenvalues $m \in \mathbb{Z}$, $|m| \leq \ell$.

$SO(3)$ has a natural representation on the space of square-integrable maps $S^2 \to \mathbb{C}$, acting by pullback as

$$f \xmapsto{R \in SO(3)} f \circ R. \tag{18}$$

Under this choice of representation, $L^2(S^2, \mathbb{C})$ should decompose as the direct sum of irreps

$$L^2(S^2, \mathbb{C}) \simeq \bigoplus_{\ell=0}^{\infty} n_\ell V_\ell \tag{19}$$

for multiplicities $n_\ell \in \mathbb{Z}_{\geq 0}$. On the other hand, we see geometrically that each eigenspace

$$\mathrm{Span}_{\mathbb{C}} \, (Y_\ell^m)_{|m|\leq \ell}^{\text{fixed } \ell} = \{f : S^2 \to \mathbb{C} \text{ with } \nabla_{S^2}^2 f = -\ell(\ell+1)\} \subset L^2(S^2, \mathbb{C}) \tag{20}$$

is closed under rotation, so we already have a decomposition into closed subspaces

$$L^2(S^2, \mathbb{C}) = \bigoplus_{\ell=0}^{\infty} \mathrm{Span}_{\mathbb{C}} \, (Y_\ell^m)_{|m|\leq \ell}^{\text{fixed } \ell}. \tag{21}$$

The irrep decomposition must be either this decomposition or some strict refinement of the same. Since the dimensions here at each ℓ are $2\ell+1$, which matches the dimension of V_ℓ, it is tempting to guess that

$$\mathrm{Span}_{\mathbb{C}} \, (Y_\ell^m)_{|m|\leq \ell}^{\text{fixed } \ell} \stackrel{?}{\simeq} V_\ell. \tag{22}$$

This turns out to be true! We can see this by an exact calculation. In this pullback representation, in the physics convention

$$(\text{group element}) = e^{i(\text{algebra element})} \tag{23}$$

the standard basis for the $so(3)$ is represented by the differential operators

$$L_i = -i\epsilon_{ijk} x_j \partial_k. \tag{24}$$

A concrete calculation gives the unique Casimir here as

$$\vec{L}^2 = L_x^2 + L_y^2 + L_z^2 = \nabla_{\mathbb{R}^3}^2 \tag{25}$$

so our unique Casimir acts on level-ℓ harmonics as

$$\vec{L}^2 Y_\ell^m = -\ell(\ell+1) Y_\ell^m \tag{26}$$

and up to isomorphism our level-ℓ harmonics all live in V_ℓ. Next, we can compute exactly

$$L_z = -i(x\partial_y - y\partial_x) = \partial_\phi \implies L_z Y_\ell^m = m Y_\ell^m. \tag{27}$$

so every value of m in V_ℓ is accounted for. So we have exactly

$$\mathrm{Span}_{\mathbb{C}} \, (Y_\ell^m)_{|m|\leq \ell}^{\text{fixed } \ell} \simeq V_\ell. \tag{28}$$

[2] Up to the usual conjugation equivalence.

[3] If we loosen to representations of $so(3)$ that don't exponentiate to single-valued reps of the group but generally represent the double cover $SU(2)$, we get further irreps at half-integer $\ell \in \frac{1}{2} + \mathbb{Z}_{\geq 0}$.

3 Harmonic Polynomials

Denote by $\mathrm{Hom}_\ell(\mathbb{R}^3)$ the space of all degree-ℓ homogeneous polynomial functions from $\mathbb{R}^3$ to $\mathbb{C}$, and by $\mathcal{H}_\ell(\mathbb{R}^3)$ the space of all harmonic polynomial functions from $\mathbb{R}^3$ to $\mathbb{C}$ which are homogeneous of degree ℓ.

$$\mathcal{H}_\ell(\mathbb{R}^3) = \{p \in \mathrm{Hom}_\ell(\mathbb{R}^3) \text{ with } \Delta p = 0\}. \tag{29}$$

For example, at $\ell = 0$ we get the space of all linear functions $\mathbb{R}^3 \to \mathbb{C}$.

$$\mathcal{H}_1(\mathbb{R}^3) = \mathrm{span}_{\mathbb{C}}(x_1, x_2, x_3). \tag{30}$$

At $\ell = 2$ we get the slightly more involved space

$$\begin{aligned}\mathcal{H}_2(\mathbb{R}^3) &= \{a_{11}x_1^2 + a_{22}x_2^2 + a_{33}x_3^2 a_{12}x_1x_2 + a_{13}x_1x_3 + a_{23}x_2x_3 \text{ such that } x_1 + x_2 + x_3 = 0\} \\ &= \mathrm{span}_{\mathbb{C}}\left(x_1x_2, x_1x_3, x_2x_3, x_1^2 - x_2^2, x_1^2 - x_3^2\right).\end{aligned} \tag{31}$$

How large is $\mathcal{H}_\ell(\mathbb{R}^3)$ in general? The space of degree-ℓ polynomials finds a natural basis in the space of monomials

$$\mathrm{Hom}_\ell(\mathbb{R}^3) = \mathrm{Span}_{\mathbb{C}}\left(\{x_1^{a_1}x_2^{a_2}x_3^{a_3} \text{ with } a_1 + a_2 + a_3 = \ell\}\right). \tag{32}$$

Specifying such a monomial means specifying $a_1 \in \{0, 1, \ldots, \ell\}$, then specifying $a_2 \in \{\ell - a_1, \ell - a_1 + 1, \ldots, \ell\}$, after which a_3 is fixed. So the number of such choices is exactly

$$\begin{aligned}\dim_{\mathbb{C}}\left(\mathrm{Hom}_\ell(\mathbb{R}^3)\right) &= \sum_{i=0}^{\ell}\sum_{j=\ell-i}^{\ell} 1 \\ &= \sum_{i=0}^{\ell}(i+1) \\ &= \frac{(\ell+2)(\ell+1)}{2}.\end{aligned} \tag{33}$$

We can compute the dimension of $\mathcal{H}_\ell(\mathbb{R}^3)$ by observing that the restriction of the Laplacian to arguments of degree ℓ

$$\Delta_\ell : \mathrm{Hom}_\ell(\mathbb{R}^3) \to \mathrm{Hom}_{\ell-2}(\mathbb{R}^3), \quad \ell \geq 2 \tag{34}$$

is surjective. As a low-order non-trivial example, x^2y^2 can be written non-uniquely as the Laplacian

$$\begin{aligned}xy &= \Delta\left(ax^4 + bx^3y + cx^2y^2 + dxy^3 + ey^4\right) \\ &= 12ax^3 + 6bxy + 2c(x^2 + y^2) + 6dxy + 12y^2\end{aligned} \tag{35}$$

which has solutions $6(b + d) = 1$ with all other coefficients vanishing. So rank-nullity gives the dimension as

$$\begin{aligned}\dim_{\mathbb{C}}\left(\mathcal{H}_\ell(\mathbb{R}^3)\right) &= \dim_{\mathbb{C}}\left(\ker\,\Delta_\ell\right) \\ &= \dim_{\mathbb{C}}\left(\mathrm{Hom}_\ell(\mathbb{R}^3)\right) - \dim_{\mathbb{C}}\left(\mathrm{Hom}_{\ell-2}(\mathbb{R}^3)\right) \\ &= \frac{(\ell+2)(\ell+1)}{2} - \frac{\ell(\ell-1)}{2} \\ &= 2\ell + 1.\end{aligned} \tag{36}$$

We introduced these homogeneous harmonic polynomials just now because there is a cute trick for turning them into a spherical harmonic. First, observe that a degree-ℓ homogeneous polynomial $p : \mathbb{R}^3 \to \mathbb{C}$ gives rise in a canonical way to $P : S^2 \to \mathbb{R}$ as

$$p(\vec{x}) = r^\ell p(\hat{x}) \equiv r^\ell p(\Omega) \equiv r^\ell P(\Omega), \quad \vec{x} = r\hat{x}. \tag{37}$$

In one of several equivalent phrasings, P is the restriction of p to S^2. Now if p is furthermore harmonic w.r.t. the Euclidean Laplacian, we can compute explicitly

$$\begin{aligned} 0 &= \nabla^2_{\mathbb{R}^3} p(\vec{x}) \\ &= \frac{1}{r^2}\frac{\partial}{\partial r}\left(r^2 \frac{\partial (r^\ell P(\Omega))}{\partial r}\right) + \frac{1}{r^2}\Delta^2_{S^2}(r^\ell P(\Omega)) \\ &= \frac{P(\Omega)}{r^2}\frac{\partial}{\partial r}\left(r^2\frac{\partial (r^\ell)}{\partial r}\right) + r^{\ell-2}\Delta^2_{S^2}P(\Omega) \\ &= \ell(\ell+1)r^{\ell-2}P(\Omega) + r^{\ell-2}\Delta^2_{S^2}P(\Omega) \\ \implies \Delta^2_{S^2}P(\Omega) &= -\ell(\ell+1)P(\Omega). \end{aligned} \tag{38}$$

So P is an eigenfunction of the sphere Laplacian, hence is a linear combination of the level-ℓ spherical harmonics. So we have cooked up a linear map

$$\mathcal{H}_\ell(\mathbb{R}^3) \to \mathrm{Span}_{\mathbb{C}} \left(Y_\ell^m\right)_{|m|\leq \ell}, \quad p \mapsto p|_{S^2} \equiv P. \tag{39}$$

Some elementary algebraic geometry tells us this map is injective: if two polynomials $p, q : \mathbb{R}^3 \to \mathbb{C}$ of the same degree ℓ give rise to identical restrictions $P, Q : S^2 \to \mathbb{C}$, then their difference $p - q : \mathbb{R}^3 \to \mathbb{C}$ vanishes on the locus $x^2 + y^2 + z^2 - 1 = 0$.

$$(p-q)(\vec{x}) = 0 \text{ whenever } |\vec{x}| = 1. \tag{40}$$

But p, q have the same degree-ℓ scaling behavior by assumption, hence so does their difference, and for general $\vec{x} = r\hat{x} \in \mathbb{R}^3$ we have

$$(p-q)(\vec{x}) = r^\ell (p-q). \tag{41}$$

So $p = q$, and as claimed our restriction map is injective. On the other hand, we have already computed the dimension of the domain and codomain of this map, finding $2\ell + 1$ in each case. So our injective map must be an isomorphism, and we conclude that we have furnished a linear equivalence between the space of degree-ℓ harmonic polynomials and the degree-ℓ spherical harmonics when both are taken as complex vector spaces.

$$\therefore \mathcal{H}_\ell(\mathbb{R}^3) \simeq \mathrm{Span}_{\mathbb{C}} \left(Y_\ell^m\right)_{|m|\leq \ell} \tag{42}$$

4 STF tensors

We find a further equivalence with the space of symmetric trace-free (STF) tensors over $\mathbb{C}^3$. To be clear, here we mean literally tensors over the linear space $\mathbb{C}^3$, not sections of tensor bundles fibered over $\mathbb{C}^3$ taken as a manifold. We also mean specifically contravariant tensors, which in the standard basis have exclusively lowered indices. To furnish this equivalence, observe that to any rank-ℓ contravariant tensor T over $\mathbb{C}^3$ we can associate in a canonical way the degree-ℓ homogeneous polynomial

$$P_T \equiv P_{a_1 \cdots a_\ell} x_{a_1} \cdots x_{a_\ell}. \tag{43}$$

This contraction is sensitive only to the part of P which is totally symmetric in all indices, so restricting our attention to such symmetric tensors is exactly the restriction that makes this map injective without cutting down its image. Better yet, since every polynomial can be written in this way, this mapping is a bijection. This polynomial has Euclidean Laplacian

$$\begin{aligned} \nabla^2_{\mathbb{R}^3} P_T &= g^{ab}\partial_{x_a}\partial_{x_b} P_{a_1 \cdots a_\ell} x_{a_1} \cdots x_{a_\ell} \\ &= \ell(\ell+1) g^{ab} P_{aba_3 \cdots a_\ell} \end{aligned} \tag{44}$$

so P_T is harmonic iff P is trace-free[4]. So we have yet another equivalence

$$P_T \in \mathcal{H}_\ell(\mathbb{R}^3) \leftrightarrow P \in \mathrm{Sym}^\ell_0(\mathbb{C}^3) \equiv \{\text{symmetric traceless rank-}\ell \text{ polynomials over } \mathbb{C}^3\}. \tag{45}$$

[4]Note that since P is totally symmetric, there is no ambiguity in which pairing of indices we are using for the trace; any one pairing vanishes iff every pairing vanishes.

5 Multipole Moments

Legendre polynomials arise from from large-distance moments of inverse-distance potentials. In the approximation $r \equiv |\vec{x}| \gg |\vec{x}_0| \equiv r_0$ we have

$$\begin{aligned}
\frac{1}{|\vec{x}-\vec{x}_0|} &= \frac{1}{\sqrt{r^2+r_0^2-2rr_0\cos\gamma}} \\
&= \frac{1}{r}\left(1-\frac{2r_0\cos\gamma}{r}+\frac{r_0^2}{r^2}\right)^{-\frac{1}{2}} \\
&= \frac{1}{r}\left(1-\frac{1}{2}\left(\frac{r_0^2}{r^2}-\frac{2r_0\cos\gamma}{r}\right)+\frac{3}{8}\left(\frac{r_0^2}{r^2}-\frac{2r_0\cos\gamma}{r}\right)^2+\cdots\right) \\
&= \frac{1}{r}\left(1+\frac{r_0}{r}\cos\gamma+\frac{r_0^2}{r^2}\frac{3\cos^2\gamma-1}{2}+\cdots\right) \\
&= \sum_{n=0}^{\infty}\frac{r_0^n}{r^{n+1}}P_n(\cos\gamma).
\end{aligned} \tag{46}$$

Here we intuitively think of $\vec{x}_0$ as the location of a source and $\vec{x}$ as the location of a distant probe, while γ is the (necessarily unsigned) angle between the two position vectors. More generally, given a compactly supported source $\rho : \mathbb{R}^3 \to \mathbb{R}$ for the Poisson equation

$$\nabla^2_{\mathbb{R}^3}\phi = \rho \tag{47}$$

we can expand in components, organizing our result in according to scaling w.r.t. r, to find

$$\begin{aligned}
\phi(\vec{x}) &= \int_{\mathbb{R}^3} d^3\vec{x}'\,\frac{\rho(\vec{x}')}{|\vec{x}-\vec{x}'|} \\
&= \int_{\mathbb{R}^3} d^3\vec{x}'\rho(\vec{x}')\frac{1}{r}\left(1-\frac{2r'\cos\gamma}{r}+\frac{r'^2}{r^2}\right)^{-\frac{1}{2}} \\
&= \int_{\mathbb{R}^3} d^3\vec{x}'\rho(\vec{x}')\frac{1}{r}\left(1+\frac{r'^2-2\vec{x}\cdot\vec{x}'}{r^2}\right)^{-\frac{1}{2}} \\
&= \int_{\mathbb{R}^3} d^3\vec{x}'\rho(\vec{x}')\frac{1}{r}\left(1-\frac{1}{2r^2}\left(r'^2-2\vec{x}\cdot\vec{x}'\right)+\frac{3}{8r^4}\left(r'^2-2\vec{x}\cdot\vec{x}'\right)^2+\cdots\right) \\
&= \int_{\mathbb{R}^3} d^3\vec{x}'\rho(\vec{x}')\frac{1}{r}\left(1+\frac{\vec{x}\cdot\vec{x}'}{r^2}-\frac{r'^2}{2r^2}+\frac{3(\vec{x}\cdot\vec{x}')^2}{2r^4}+\mathcal{O}\left(\frac{1}{r^3}\right)\right) \\
&= \int_{\mathbb{R}^3} d^3\vec{x}'\rho(\vec{x}')\left(\frac{1}{r}+\frac{x_i'x_i}{r^3}+\frac{\left(3x_i'x_j'-r'^2\delta_{ij}\right)x_ix_j}{2r^5}+\mathcal{O}\left(\frac{1}{r^4}\right)\right) \\
&\equiv \frac{Q}{r}+\frac{Q_ix_i}{r^3}+\frac{Q_{ij}x_ix_j}{2r^5}+\frac{Q_{ijk}x_ix_jx_k}{3!r^7}+\cdots.
\end{aligned} \tag{48}$$

Notice that this implicit definition of that $Q's$ in terms of their results after contraction against x strings leaves exactly enough ambiguity to WLOG take the Q's to be totally symmetric; we will take this as a further part of their definition. As a sanity check, we can explicitly compute the Euclidean Laplacian of this expression. In fact

all terms exhibit the same behavior here, as exemplified by the fourth one:

$$\begin{aligned}
\nabla^2_{\mathbb{R}^3} \frac{Q_{ijk}x_i x_j x_k}{3! r^7} &= \delta_{ab} \frac{\partial^2}{\partial x_a x_b} \left(\frac{Q_{ijk}x_i x_j x_k}{3! r^7} \right) \\
&= \delta_{ab} \frac{\partial}{\partial x_a} \left(\frac{Q_{bjk}x_j x_k}{2! r^7} - \frac{7 Q_{ijk}x_i x_j x_k x_b}{3! r^9} \right) \\
&= \delta_{ab} \left(\frac{Q_{bak}x_k}{r^7} - \frac{7 Q_{bjk}x_j x_k x_a}{2! r^9} - \frac{7 Q_{ajk}x_j x_k x_b}{2! r^9} - \frac{7 Q_{ijk}x_i x_j x_k \delta_{ab}}{3! r^9} + \frac{63 Q_{ijk}x_i x_j x_k x_a x_b}{3! r^{11}} \right) \\
&= \frac{Q_{aak}x_k}{r^7} - \frac{7 Q_{ajk}x_j x_k x_a}{2! r^9} - \frac{7 Q_{ajk}x_j x_k x_a}{2! r^9} - \frac{7 Q_{ijk}x_i x_j x_k}{2! r^9} + \frac{63 Q_{ijk}x_i x_j x_k}{3! r^9} \\
&= \frac{Q_{aak}x_k}{r^7}.
\end{aligned} \tag{49}$$

In general, we have

$$\begin{aligned}
\nabla^2_{\mathbb{R}^3} \phi(\vec{x}) &= \delta^{ab} \frac{\partial^2}{\partial x_a x_b} \left(\frac{Q}{r} + \frac{Q_i x_i}{r^3} + \frac{Q_{ij} x_i x_j}{2 r^5} + \frac{Q_{ijk} x_i x_j x_k}{3! r^7} + \cdots \right) \\
&= \frac{Q_{aa}}{r^5} + \frac{Q_{aai} x_i}{r^7} + \frac{Q_{aaij} x_i x_j}{2! r^9} + \cdots + \frac{Q_{aai_3 \cdots i_\ell} x_{i_3} \cdots x_{i_\ell}}{(\ell - 2)! r^{2\ell + 1}} + \cdots .
\end{aligned} \tag{50}$$

If we are being careful this calculation has a restricted domain of validity. First, we have quietly assumed that $\vec{x} \neq 0$, since more generally

$$\nabla^2 \left(\frac{1}{r} \right) = -4\pi \delta^{(3)}(\vec{x}). \tag{51}$$

More interestingly, we know that $\phi(\vec{x})$ is harmonic away from the compact-by-assumption support of ρ, which implies that all of the traces must vanish.

$$Q_{aai_3 \cdots i_\ell} = 0. \tag{52}$$

But this implies our series expansion for $\nabla^2_{\mathbb{R}^3} \phi(\vec{x})$ vanishes uniformly. So it must be that the domain of validity of this series expansion (which we generically expect to be limited to sufficiently large r anyway for convergence reasons) is at most the exterior of the support of ρ, since inside this support we generally have $\nabla^2_{\mathbb{R}^3} \phi(\vec{x}) = \rho(\vec{x}) \neq 0$. So our series expansion holds exactly, but only for sufficiently large r.

In fact we never needed to talk about ρ; it was just a way of cooking up a function ϕ which, at large distances, is harmonic and decaying. Any scalar function which is harmonic and polynomial-decaying at large r can be written

$$\phi(\vec{x}) = \sum_{\ell=0}^{\infty} \frac{Q_{i_1 \cdots i_\ell} x_{i_1} \cdots x_{i_\ell}}{r^{2\ell+1}} \text{ at sufficiently large } r \tag{53}$$

where

$$\nabla_{\mathbb{R}^3} \phi(\vec{x}) = 0 \text{ at large } r \implies Q_{aai_3 \cdots i_\ell} = 0. \tag{54}$$

Pulling out factor of r as $\vec{x} = r\hat{x}$ lets us map each level-ℓ multipole is mapped invertibly to a combination of level-ℓ spherical harmonic, and we get the whole nice story above. For example, in the normalization

$$Y_{\ell m} = e^{im\phi} P_\ell^m(\cos\theta) \tag{55}$$

and by comparison with the $\ell = 1$ associated Legendre polynomials

$$\begin{aligned}
P_1^{-1}(x) &= \frac{1}{2}(1 - x^2)^{\frac{1}{2}}, \\
P_1^0(x) &= x, \\
P_1^1(x) &= -(1 - x^2)^{\frac{1}{2}}
\end{aligned} \tag{56}$$

we find

$$\begin{aligned}\frac{Q_i x_i}{r^3} &= \frac{Q_1 \sin\theta\cos\phi + Q_2 \sin\theta\sin\phi + Q_3\cos\theta}{r^2} \\ &= \frac{e^{i\phi}\frac{Q_1 - iQ_2}{2}\sin\theta + e^{-i\phi}\frac{Q_1+iQ_2}{2}\sin\theta + Q_3\cos\theta}{r^2} \\ &= \frac{Q_3 P_1^0(\cos\theta) - e^{i\phi}\frac{Q_1-iQ_2}{2}P_1^1(\cos\theta) + e^{-i\phi}(Q_1+iQ_2)P_1^{-1}(\cos\theta)}{r^2} \\ &= \frac{(Q_1+iQ_2)Y_{1-1}(\theta,\phi) + Q_3 Y_{10}(\theta,\phi) - \frac{Q_1-iQ_2}{2}Y_{11}(\theta,\phi)}{r^2}.\end{aligned} \tag{57}$$

6 Shifting the Origin

Take a harmonic function with expansion as above, but now interpret the coefficients as a function of the base point about which we expand, which in our first analysis was exactly the origin. Then we have slightly more explicitly

$$\phi(\vec{x}) = \frac{Q(0)}{r} + \frac{Q_i(0)x_i}{r^3} + \frac{Q_{ij}(0)x_i x_j}{2r^5} + \frac{Q_{ijk}(0)x_i x_j x_k}{3! r^7} + \cdots. \tag{58}$$

About a more general expansion point $\vec{y}$, we have

$$\phi(\vec{x}) = \frac{Q(\vec{y})}{|\vec{x}-\vec{y}|} + \frac{Q_i(\vec{y})(x_i-y_i)}{|\vec{x}-\vec{y}|^3} + \frac{Q_{ij}(\vec{y})(x_i-y_i)(x_j-y_j)}{2|\vec{x}-\vec{y}|^5} + \frac{Q_{ijk}(\vec{y})(x_i-y_i)(x_j-y_j)(x_k-y_k)}{3!|\vec{x}-\vec{y}|^7} + \cdots. \tag{59}$$

On the other hand, if we expand

$$\begin{aligned}\frac{1}{r^n} &= \frac{1}{|(\vec{x}-\vec{y})+\vec{y}|^n} \\ &= \frac{1}{|\vec{x}-\vec{y}|^n} + \frac{a_i^{(n)} y_i}{|\vec{x}-\vec{y}|^{n+1}} + \frac{a_{ij}^{(n)} y_i y_j}{|\vec{x}-\vec{y}|^{n+2}} + \cdots\end{aligned} \tag{60}$$

(the exact coefficients aren't important) then our expansion about the origin tells us that we can write[5] each $Q_{I_\ell}(\vec{y})$ as a function of the $Q_{J_\ell}(0)$'s for $\ell' \le \ell$. For example, $Q_{ij}(\vec{y})$ is a linear combination of $Q_0(0), Q_i(0), Q_{ij}(0)$. Conversely, $Q_{ij}(0)$ contributes to $Q_{i_1\cdots i_\ell}(\vec{y})$ for all $\ell \ge 2$.

Now we think of $Q_{I_\ell} \equiv Q_{i_1\cdots i_\ell}$ as a tensor field over $\mathbb{R}^3$, assigning a tensor value at each[6] $\vec{y} \in \mathbb{R}^3$. This has the peculiar property that, if we know $Q_{I_\ell}(\vec{y})$ for all $\vec{y} \in \mathbb{R}^3$, this fixes the same data for all lower $Q_{I_{\ell'}}$ with $\ell' < \ell$. Conversely, knowing all $Q_{I_{\ell'}}$ at just one fixed point fixes Q_{I_ℓ} as a tensor field. So knowing all moments up to level ℓ at one fixed point is equivalent to knowing Q_{I_ℓ} everywhere.

This action of translation linearly mixes up up the Q's with coefficients determined only by the relative positions of the two base points, and in such a way that each new level-ℓ charge only gets contributions from the old charges at level $\le \ell$. Algebraically, this means we have a representation of the group of 3D translations (composed in the usual way by vector addition) schematically of the form

$$\begin{bmatrix} Q(\vec{y}) \\ Q_i(\vec{y}) \\ Q_{ij}(\vec{y}) \\ \vdots \end{bmatrix} = \begin{bmatrix} \#Q(\vec{x}) \\ \#Q(\vec{x}) + \#Q_i(\vec{x}) \\ \#Q(\vec{x}) + \#Q_i(\vec{x}) + \#Q_{ij}(\vec{x}) \\ \vdots \end{bmatrix} = \begin{bmatrix} \# & 0 & 0 & \cdots & \\ \# & \# & 0 & \cdots & \\ \# & \# & \# & \cdots & \\ \vdots & \vdots & \vdots & & \ddots \end{bmatrix} \begin{bmatrix} Q(\vec{x}) \\ Q_i(\vec{x}) \\ Q_{ij}(\vec{x}) \\ \vdots \end{bmatrix} \tag{61}$$

where the black-boxed coefficients # are dressed with appropriate indices and depend only $\vec{x}-\vec{y}$.

[5] Here I is a multi-index.

[6] Really, each $\vec{y}$ large enough for convergence.

7 Multipole Moments as Conformal Killing Tensors

We take our space V to be the complement in $\mathbb{R}^3$ of some smooth, solid region which is topologically a solid ball, which we think of as the subtracted-off support of the source charge density. Topologically, V is then $S^2 \times \mathbb{R}$. Let φ be any harmonic function on V. Write the metric on V as h_{ab}, and let $\xi = \xi^a \partial_a$ be a conformal Killing vector

$$\nabla_{(a}\xi_{b)} = \frac{1}{3} h_{ab} \nabla^m \xi_m \tag{62}$$

where we take the good convention for symmetrizing

$$A_{(ab)} \equiv \frac{1}{2}(A_{ab} + A_{ba}). \tag{63}$$

Then for all $s \geq 0$, the dual vector field $U^{(s)} = U_m dx^m$ with components

$$U^{(s)}_m = \xi^{a_1} \cdots \xi^{a_s} \nabla_m \nabla_{a_1} \cdots \nabla_{a_s} \varphi \tag{64}$$

is harmonic. To see this, we repeatedly use commutativity of covariant derivatives in flat space to compute directly

$$\begin{aligned} \nabla^m U^{(s)}_m &= s(\nabla^m \xi^{a_1}) \cdots \xi^{a_s} \nabla_m \nabla_{a_1} \cdots \nabla_{a_s} \varphi + \xi^{a_1} \cdots \xi^{a_s} \nabla^m \nabla_m \nabla_{a_1} \cdots \nabla_{a_s} \varphi \\ &= s(\nabla^{(m} \xi^{a_1)}) \cdots \xi^{a_s} \nabla_{a_2} \cdots \nabla_{a_s} \nabla_m \nabla_{a_1} \varphi + \xi^{a_1} \cdots \xi^{a_s} \nabla_{a_1} \cdots \nabla_{a_s} \underbrace{\nabla^m \nabla_m \varphi}_{=0} \\ &= \frac{s}{3} h^{m a_1} \cdots \xi^{a_s} \nabla_{a_2} \cdots \nabla_{a_s} \nabla_m \nabla_{a_1} \varphi \\ &= \frac{s}{3} \cdots \xi^{a_s} \nabla_{a_2} \cdots \nabla_{a_s} \underbrace{\nabla_m \nabla^m \varphi}_{=0}. \end{aligned} \tag{65}$$

Then as long as K any outward-oriented smooth surface homeomorphic to S^2 and enclosing the "hole" in V, the integral

$$Q^{(s)} \equiv Q^{(s)}[\xi] \equiv \frac{1}{4\pi} \int_K U^{(s)} \tag{66}$$

is independent of the particular choice of K. This integral therefore associates a real number to any conformal Killing vector; we note that the latter form a real 10-dimensional Lie algebra (3 translations, 3 rotations, 3 special conformal transformations, plus dilatation). Geroch [**?**] claims that "This collection ($s = 0, 1, 2, \ldots$) of multilinear mappings contains precisely the information of the multipole moments of φ." To make this clearer, we denote by $\mathcal{C}$ the real vector space of conformal Killing vectors. As just mentioned, this has a natural basis

$$\{\varepsilon_\alpha\}_{\alpha=1}^{10} \{\varepsilon_1, \ldots, \varepsilon_{10}\} = \{\text{translations, rotations, SCTs, dilatation}\} \tag{67}$$

though there are infinitely many other choices, and all of the following will be agnostic w.r.t. the choice of basis. Crucially, here α (and moving forward, other Greek indices) enumerate basis elements and components in this finite-dimensional space; they are not tensor field indices. In the mildly unfortunate notation used by [**?**], we accordingly expand as, *e.g.*,

$$\xi = \xi^\alpha \varepsilon_\alpha \in \mathcal{C}. \tag{68}$$

If we extend the mapping $Q^{(s)}$ in the unique totally symmetric way to take in s independent conformal Killing vectors

$$Q^{(s)} : \mathcal{C} \otimes \cdots \otimes \mathcal{C} \to \mathbb{R}, \quad Q^{(s)}[\xi_1, \ldots, \xi_s] \equiv \frac{1}{4\pi} \int_K \xi_1^{a_1} \cdots \xi_s^{a_s} \nabla_m \nabla_{a_1} \cdots \nabla_{a_s} \varphi \tag{69}$$

the result can be thought of as a symmetric s-tensor over $\mathcal{C}$, with action in $\mathcal{C}$-components written as

$$Q^{(s)}[\xi_1, \ldots, \xi_s] = Q^{(s)}_{\alpha_1 \cdots \alpha_s} \xi_1^{\alpha_1} \cdots \xi_s^{\alpha_s} \equiv Q_{\alpha_1 \cdots \alpha_s} \xi_1^{\alpha_1} \cdots \xi_s^{\alpha_s} \in \mathbb{R} \tag{70}$$

where the presence of s indices lets us drop the superscript (s) without fear of ambiguity. The corresponding intrinsic object is the totally symmetric rank-s covariant tensor

$$Q_{\alpha_1 \cdots \alpha_s} (\varepsilon *)^{\alpha_1} \cdots (\varepsilon *)^{\alpha_s} \in \bigotimes_{\text{sym}}^{s} \mathcal{C}^* \tag{71}$$

A major goal of this exposition is to explain how these Q's relate to the usual symmetric tensors which characterize the multipole expansion of φ. [**?**] claims that the former totally characterizes the latter, which from all of our earlier work is immensely plausible.

The CKVs are richer than just a vector space, as they form a Lie algebra under vector field commutation. In any basis, we have associated structure constants

$$([\xi, \xi'])^\mu = C^\mu_{\alpha\beta} \xi^\alpha \xi'^\beta \tag{72}$$

from which we can built the usual Killing form

$$G_{\alpha\beta} \equiv -C^\mu_{\ \nu\alpha} C^\nu_{\ \mu\beta}. \tag{73}$$

A concrete calculation using the known structure of the CKVs of $\mathbb{R}^3$ show that this symmetric quadratic form is non-singular, with signature $(+,+,+,+,+,+,-,-,-,-)$. We write the corresponding inverse metric with components $G^{\alpha\beta}$. With this we can raise the $\mathcal{C}$-indices to product a totally symmetric rank-s contravariant tensor

$$Q^{\alpha_1 \cdots \alpha_s} \epsilon_{\alpha_1} \cdots \varepsilon_{\alpha_s} \in \bigotimes_{\text{sym}}^s \mathcal{C}. \tag{74}$$

The linear space of such tensors is spanned by uniform strings $\xi \otimes \cdots \otimes \xi$ for $\xi \in \mathcal{C}$. The number of linearly independent such strings will be written as D_s (the particular value is unimportant here; maybe we'll add it as an appendix). In any case we can therefore write (perhaps excessively verbosely)

$$\begin{aligned} Q^{\alpha_1 \cdots \alpha_s} &= \sum_{i=1}^{D_s} \xi_i^{\alpha_1} \cdots \xi_i^{\alpha_s} \\ \implies Q^{\alpha_1 \cdots \alpha_s} \epsilon_{\alpha_1} \cdots \varepsilon_{\alpha_s} &= \sum_{i=1}^{D_s} \xi_i^{\alpha_1} \cdots \xi_i^{\alpha_s} \epsilon_{\alpha_1} \cdots \varepsilon_{\alpha_s} \\ &= \sum_{i=1}^{D_s} \xi_i \otimes \cdots \otimes \xi_i \in \bigotimes_{\text{sym}}^s \mathcal{C}. \end{aligned} \tag{75}$$

To this we can associate the rank-s contravariant spatial object, namely the tensor field with components

$$Q^{a_1 \cdots a_s}(x) = \sum_{i=1}^{D} \xi_i^{a_1}(x) \cdots \xi_i^{a_s}(x). \tag{76}$$

8 Conformal Killing Tensors and Trace Subtraction

Per [**?**] appendix A, a conformal Kiling vector $P = P^a \partial_a$ may be defined as a vector field such that, for any null geodesic with affine tangent vector $\ell = \ell^a \partial_a$, $\ell_a P^a$ is constant along the geodesic. Why is that consistent with other definitions? An null geodesic affinely parametrized as $x(\lambda)$ has affine tangent vector

$$\ell^a = \frac{dx^a}{d\lambda} \tag{77}$$

and the geodesic condition is variously

$$0 = \frac{D}{d\lambda} \frac{dx^a}{d\lambda} = \ell^b \nabla_b \ell^a = (\nabla_\ell \ell)^a \,. \tag{78}$$

$\ell_a P^a = \langle \ell, P \rangle = g(\ell, P)$ is then constant along the curve iff

$$0 = \nabla_\ell g(\ell, P) = \ell^a \nabla_a (\ell_b P^b) = \ell^a \left(P^b \nabla_a \ell_b + \ell^b \nabla_a P_b \right). \tag{79}$$

Still not clear, but in our space of interest we have no null vectors anyway! In this case, [**?**] says we simply take a conformal Killing tensor to be any tensor $P^{a_1\cdots a_s}$ such that

$$\nabla^{(m}P^{a_1\cdots a_s)} = h^{(ma_1}T^{a_2\cdots a_s)} \tag{80}$$

for some T. Does this agree with our definition of a CKV? At $s = 1$ it reduces to

$$\nabla^{(m}P^{a)} = h^{ma}T \tag{81}$$

for T some scalar, which we can solve for by taking the trace

$$T = \frac{1}{3}h_{ab}\nabla^{(a}P^{b)} = \frac{1}{3}h_{ab}\nabla^{a}P^{b} = \frac{1}{3}\nabla_{a}P^{a} \tag{82}$$

So this is equivalent to the CKV definition

$$\nabla_{(a}P_{b)} = \frac{1}{3}h_{ab}\nabla_{c}P^{c}. \tag{83}$$

What about $s = 2$? There we have

$$\nabla^{(m}P^{ab)} = h^{(ma}T^{b)} \tag{84}$$

Can we think of this in terms of being killed by trace subtraction? Given a totally symmetric tensor A, the trace

$$t_a \equiv h^{bc}A_{abc} \tag{85}$$

is independent of the pair of indices we choose to contract. Naively trace-subtracting A gives

$$A'^{abc} = A^{abc} - \frac{1}{n}h^{ab}t^{c} \tag{86}$$

which is trace-free only necessarily in ab.

$$g_{ab}A'^{abc} = h_{ab}A^{abc} - \frac{1}{n}h^{ab}h_{ab}t_c = t_c - t_c = 0. \tag{87}$$

The fix is to symmetrize in the subtracted-off term

$$A^{abc}_{TF} = A^{abc} - \#h^{(ab}t^{c)} \tag{88}$$

where # is a coefficient TBD. We can fix it by demanding that the RHS has the desired vanishing trace.

$$\begin{aligned} h_{ab}A^{abc} &= \#h_{ab}h^{(ab}t^{c)} \\ \implies t^c &= \frac{\#}{3}h_{ab}\left(h^{ab}t^{c} + h^{ac}t^{b} + h^{bc}t^{a}\right) \\ &= \frac{\#}{3}\left(nt^{c} + t^{c} + t^{c}\right) \\ \implies \# &= \frac{3}{n+2} \end{aligned} \tag{89}$$

so bonafide trace-subtracted tensor is

$$A^{abc}_{TF} = A^{abc} - \frac{3}{n+2}h^{(ab}t^{c)} = A^{abc} - \frac{h^{ab}t^{c} + h^{ac}t^{b} + h^{bc}t^{a}}{n+2}. \tag{90}$$

So we see that, in $s = 2$ and by identical logic for higher ranks, [**?**]'s definition is indeed a pure-trace property. More generally, our trace term looks like

$$t_{a_3\cdots a_s} \equiv h^{a_1a_2}A_{a_1a_2\cdots a_s}, \tag{91}$$

our symmetric trace-subtracted tensor should be

$$A^{a_1\cdots a_s}_{\text{TF}} \stackrel{?}{=} A^{a_1\cdots a_s} - \#h^{(a_1a_2}t^{a_3\cdots a_s)}, \tag{92}$$

and the coefficient is found (after carefully enumerating the distinct types of contractions resulting from pairing up h against the symmetrized product ht) as

$$\begin{aligned}
t^{a_3\cdots a_s} &= h_{ab}A^{aba_3\cdots a_s} \\
&= \#h_{ab}h^{(ab}t^{a_3\cdots a_s)} \\
&= \#h_{ab}\frac{2!(s-2)!h^{ab}t^{a_3\cdots a_s} + 2!2(s-2)(s-2)!h^{b(a_3}t^{(a_4a_5\cdots a_s)a} + (s-2)(s-3)(s-2)!h^{(a_3a_4}t^{a_5\cdots a_s)ab}}{s!} \\
&= \#h_{ab}\frac{2h^{ab}t^{a_3\cdots a_s} + 4(s-2)h^{b(a_3}t^{a_4a_5\cdots a_s)a} + (s-2)(s-3)h^{(a_3a_4}t^{a_5\cdots a_s)ab}}{s(s-1)} \\
&= \#\frac{2nt^{a_3\cdots a_s} + 4(s-2)t^{a_3a_4a_5\cdots a_s} + (s-2)(s-3)h_{ab}h^{(a_3a_4}t^{a_5\cdots a_s)ab}}{s(s-1)}.
\end{aligned} \tag{93}$$

This is slightly funny! It's not clear what to do with this last term, which isn't obviously proportionate to $t^{a_3\cdots a_s}$, but rather to the metric times a second-traced object, and which only appears for $s \geq 4$. There is nothing wrong in our calculation, rather our trace-subtraction method was too naive for $s = 4$, roughly because the trace itself can have a nonzero trace. Said in a different order, for $s \geq 4$ we find that our hoped trace-free difference has trace

$$\begin{aligned}
&h_{ab}\left(A^{aba_3\cdots a_s} - \#h^{(ab}t^{a_3\cdots a_s)}\right) \\
&= t^{a_3\cdots a_s} - \#\frac{2nt^{a_3\cdots a_s} + 4(s-2)t^{a_3a_4a_5\cdots a_s} + (s-2)(s-3)h_{ab}h^{(a_3a_4}t^{a_5\cdots a_s)ab}}{s(s-1)} \\
&= t^{a_3\cdots a_s} - \#\frac{2n+4s-8}{s(s-1)}t^{a_3a_4a_5\cdots a_s} - \#\frac{h_{ab}h^{(a_3a_4}t^{a_5\cdots a_s)ab}}{s(s-1)}.
\end{aligned} \tag{94}$$

So no choice of # actually gives a trace-free result. For $s = 4$ exactly, it suffices to add just one secondary-trace term into our ansatz

$$A^{abcd}_{\text{TF}} \stackrel{?}{=} A^{abcd} - \alpha h^{(ab}t^{cd)} - \beta h^{(ab}h^{cd)}t' \tag{95}$$

where t' is the second-trace

$$t' \equiv h_{ab}t^{ab} = h_{ab}h_{cd}A^{abcd}. \tag{96}$$

Demanding that the trace of this quantity give zero gives the condition

$$\begin{aligned}
t^{cd} &= h_{ab}A^{abcd} \\
&= \alpha h_{ab}h^{(ab}t^{cd)} + \beta h_{ab}h^{(ab}h^{cd)}t' \\
&= \alpha h_{ab}\frac{4h^{ab}t^{cd} + 16h^{ac}t^{bd} + 4h^{cd}t^{ab}}{4!} + \beta h_{ab}\frac{8h^{ab}h^{cd} + 16h^{ac}h^{bd}}{4!}t' \\
&= \alpha\frac{nt^{cd} + 4t^{cd} + h^{cd}t'}{3!} + \beta\frac{nh^{cd} + 2h^{cd}}{3}t' \\
&= \alpha\frac{n+4}{6}t^{cd} + \left(\frac{\alpha}{3!} + \frac{\beta(n+2)}{3}\right)t'
\end{aligned} \tag{97}$$

from which we read off the solution

$$\alpha = \frac{6}{n+4}, \quad \beta = -\frac{\alpha}{2n+4} = -\frac{3}{(n+2)(n+4)}. \tag{98}$$

At $s = 4$ our trace-subtracted object is therefore

$$A^{abcd}_{\text{TF}} = A^{abcd} - \frac{6}{n+4}h^{(ab}t^{cd)} + \frac{3}{(n+2)(n+4)}h^{(ab}h^{cd)}t'. \tag{99}$$

If we repeat the above story in a slightly different order, we find that we can still think of [**?**]'s condition as saying that a tensor if a CKV iff its symmetric derivative is trace-free. For a general-spin symmetric tensor, the trace-subtracted version is

$$A^{a_1\cdots a_s}_{\text{TF}} = A^{a_1\cdots a_s} + \#h^{(a_1a_2}\left(\text{Tr}A\right)^{a_3\cdots a_s)} + \#h^{(a_1a_2}h^{a_3a_4}\left(\text{Tr}^2A\right)^{a_5\cdots a_s)} + \cdots \tag{100}$$

for # coefficients depending only on n and s, which we don't need for our argument and so will black-box. In terms of the (notable symmetric and associated operation) $\cap$ used by [**?**], this is

$$A_{\mathrm{TF}} = A + \#h \cap \mathrm{Tr}A + \#h \cap h \cap \mathrm{Tr}^2 A + \cdots \tag{101}$$

In fact all of the correction terms on the RHS are unique. More precisely, given any totally symmetric A of rank s, there exists a unique totally symmetric U of rank $s-2$ such that

$$A^{a_1 \cdots a_s} = A_{\mathrm{TF}}^{a_1 \cdots a_s} + h^{(a_1 a_2} U^{a_3 \cdots a_s)}. \tag{102}$$

or equivalently

$$A = A_{\mathrm{TF}} + h \cap U. \tag{103}$$

Uniqueness of U given A immediately implies A_{TF} is likewise unique. To say that A is pure-trace, therefore, is to say that its trace-free part vanishes, or equivalently that A has the form

$$A \text{ "pure trace"} \implies A = 0 + h \cap U. \tag{104}$$

But this is exactly [**?**]'s condition on the symmetric derivative

$$\nabla^{(m} P^{ab \cdots c)} = h^{(ma} T^{b \cdots c)} \tag{105}$$

for some totally symmetric T.