

Sparse somatosensory coding: towards explaining and predicting the response properties of rodent afferent pathway neurons

Mathew H Evans

Psychology Department, University of Sheffield, UK

E-mail: mat.evans@sheffield.ac.uk

Early sensory encoding appears to follow principles of efficiency and sparsity in both vision and audition [1]. Can a similar sparse coding approach describe the response properties of rodent afferent pathway neurons? The connectivity and general receptive field properties of this system have been characterized, barrel cortex in particular. However the response properties of the afferent pathway neurons have not been described beyond the trigeminal ganglion and VPM neurons of the first lemniscal pathway, partly due to the multi-whisker connectivity of these remaining pathways and the difficulty of stimulating such a system. Primary afferent neurons have been labeled as slowly adapting and rapidly adapting units [2], Figure 1A, and VPM neurons have been described as a population of diverse, precise kinetic feature detectors [3] Figure 1C. Can these responses be explained by sparse coding principles, tuned to object properties in the world? Early work applying a sparse ICA to spectrogram images from artificial whisker data predicted spectro-temporal receptive fields in cortex [4], of which there is tentative evidence [5]. However, rodents are susceptible to velocity-amplitude discrimination ‘illusions’, suggesting that whisker deflections are not encoded with spectro-temporal precision [5]. We tested a non-linear sparse coding approach [6] on multiple sets of artificial whisker time series data [7,8]. Initial results are shown in Figure 1. The first 3 principle eigenvectors from PCA analysis (Figure 1B) resemble primary afferent response properties (Figure 1A), and some sparsely coded basis functions (Figure 1D), resemble VPM thalamus responses (Figure 1C). We hope to evaluate this approach by verifying model and neural response similarity, and by extension to multi-whisker stimuli.




Figure 1. Neuron and model responses to whisker deflection. **A** Primary afferent responses. **B** First 3 principal eigenvectors from PCA on artificial data. **C** VPM thalamus responses. **D** Subset of sparse bases

Acknowledgements. This work is supported by an EPSRC Doctoral Prize Fellowship.

References

1. Olshausen, B. A., & Field, D. J. **Sparse coding of sensory inputs.** *Current opinion in neurobiology*, 2004, **14(4)**: 481-487.
2. Szwed, M., Bagdasarian, K., & Ahissar, E. **Encoding of vibrissal active touch.** *Neuron*, 2003, **40(3)**: 621-630.
3. Petersen, R. S., Brambilla, M., Bale, M. R., Alenda, A., Panzeri, S., Montemurro, M. A., & Maravall, M. **Diverse and temporally precise kinetic feature selectivity in the VPM thalamic nucleus.** *Neuron*, 2008, **60(5)**: 890-903.
4. Hafner, V. V., Fend, M., König, P., Körding, K. P., & Paul, K. **Predicting properties of the rat somatosensory system by sparse coding.** *Neural Information Processing Letters and Reviews*, 2004, **4(1)**: 11-18.
5. Diamond, M. E., & Arabzadeh, E. **Whisker sensory system—From receptor to decision.** *Progress in Neurobiology*. 2012.
6. Lee, H., Battle, A., Raina, R., & Ng, A. Y. **Efficient sparse coding algorithms.** *Advances in neural information processing systems*, 2007, **19**: 801.
7. Evans, M., Fox, C. W., Pearson, M. J., & Prescott, T. J. **Spectral template based classification of robotic whisker sensor signals in a floor texture discrimination task.** *Proceedings TAROS*, 2009, 19-24.
8. Sullivan, J., Mitchinson, B., Pearson, M. J., Evans, M., Lepora, N. F., Fox, C. W., Melhuish, C., & Prescott, T. J. **Tactile discrimination using active whisker sensors.** *Sensors Journal, IEEE*, 2012, **12(2)**: 350-362.