

Inequalities

Youming Zhao

Email: yoming0.zhao@gmail.com

First draft: January 1, 2022 Last update: October 14, 2023

Contents

1	Definitions	1
1.1	Norm	1
1.2	Vector p -norms	2
1.3	Dual norm	2
2	Quadratic mean, average mean, geometric mean, and harmonic mean	3
2.1	An inequality about the difference between AM and GM	6
2.2	Power mean inequality	6
2.3	Weighted power mean inequality	6
2.4	Applications	7
3	Young inequality	7
4	Hölder inequality	7
4.1	Generalized Hölder inequality	8
4.2	Cauchy-Schwarz inequality	10
4.3	Two variants of Cauchy-Schwarz inequality	10
5	Minkowski inequality	10
6	Rearrangement inequalities	11
6.1	Chebyshev's inequality	12

1 Definitions

1.1 Norm

A function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ with $\text{dom} f = \mathbf{R}^n$ is called a norm if

- f is nonnegative: $f(x) \geq 0$ for all $x \in \mathbf{R}^n$
- f is definite: $f(x) = 0$ if and only if $x = 0$
- f is homogeneous: $f(tx) = |t|f(x)$, for all $x \in \mathbf{R}^n$ and $t \in \mathbf{R}$
- f satisfies the triangle inequality: $f(x + y) \leq f(x) + f(y)$, for all $x, y \in \mathbf{R}^n$.

1.2 Vector p -norms

A vector p -norm denoted $\|\cdot\|_p$ is defined as,

$$\|\mathbf{x}\|_p := (|x_1|^p + |x_2|^p + \cdots + |x_n|^p)^{\frac{1}{p}} = \left(\sum_{i=1}^n |x_i|^p\right)^{\frac{1}{p}}, \quad p \geq 1, \mathbf{x} \in \mathbf{R}^n.$$

The most commonly used vector p -norms are ℓ_1 -norm, ℓ_2 -norm, and ℓ_∞ -norm.

$$\begin{aligned} \|\mathbf{x}\|_1 &:= |x_1| + |x_2| + \cdots + |x_n| = \sum_{i=1}^n |x_i| \\ \|\mathbf{x}\|_2 &:= \sqrt{|x_1|^2 + |x_2|^2 + \cdots + |x_n|^2} = \sqrt{\sum_{i=1}^n |x_i|^2} = \sqrt{x^T x} \\ \|\mathbf{x}\|_\infty &:= \max_{1 \leq i \leq n} |x_i|. \end{aligned}$$

where the $|\cdot|$ sign is not omitted in the definition of $\|\cdot\|_2$ to emphasize the importance of the $|\cdot|$ operation in the calculations of all p -norms. $\|\mathbf{x}\|_2$ is also known as Euclidean norm.

1.3 Dual norm

Let $\|\cdot\|$ be a norm on \mathbf{R}^n . The associated dual norm, denoted $\|\cdot\|_*$, is defined as

$$\|z\|_* = \sup\{z^T x \mid \|x\| \leq 1\}.$$

It is easy to show that the dual norm satisfies all properties of a norm, so the dual norm is a norm.

Proposition 1 (Inner product, norm, and dual norm). *Let $\|\cdot\|$ be a norm on \mathbf{R}^n . The associated dual norm $\|z\|_*$ satisfies*

$$z^T x \leq \|x\| \|z\|_*,$$

where $x, z \in \mathbf{R}^n$.

Proof. From the definition of dual norm, we have

$$z^T x \leq \|x\| \|z\|_*$$

with all x satisfying $\|x\| \leq 1$. So the inequality also holds for $\|x\| = 1$,

$$z^T x \leq \|x\| \|z\|_* = \|x\| \|z\|_*.$$

Let $x = ty$ with $t > 0$. Then we get

$$z^T(ty) \leq \|ty\| \|z\|_* = t\|y\| \|z\|_* \iff z^T y \leq \|y\| \|z\|_*,$$

as desired. Note that there is no requirement on the value of $\|y\|$. \square

Proposition 2 (Dual norm of Euclidean norm is Euclidean norm). *The dual norm of Euclidean norm is Euclidean norm, i.e.,*

$$\|z\|_2 = \sup\{z^T x \mid \|x\|_2 \leq 1\}.$$

46 *Proof.* According to Cauchy-Schwarz inequality, we get

$$z^T y \leq \|y\|_2 \|z\|_2.$$

47 Given z , the equality holds if and only if $y = z$. If $\|y\|_2$ is required to be not greater than 1, then
 48 $y = \frac{z}{\|z\|_2}$ which maximizes $z^T y$ and the maximum is $z^T \frac{z}{\|z\|_2} = \|z\|_2$. Thus, $\|z\|_2 = \sup\{z^T x \mid \|x\|_2 \leq 1\}$,
 49 which is exactly the definition of dual norm. \square

50 **Proposition 3 (Dual norm of ℓ_∞ -norm is ℓ_1 -norm).** *The dual norm of the ℓ_∞ -norm is the*
 51 *ℓ_1 -norm.*

52 *Proof.* Since $\|x\|_\infty \leq 1$, $x_i = \text{sgn}(z_i)$ for each $i \in \{1, \dots, n\}$ can maximize $z^T x$, where $\text{sgn}(t)$ is the
 53 sign function which outputs 1 for positive inputs, -1 for negative inputs, and 0 for zero inputs,
 54 respectively. Thus,

$$\sup\{z^T x \mid \|x\|_\infty \leq 1\} = \sum_{i=1}^n |z_i| = \|z\|_1,$$

55 as desired. \square

56 **Proposition 4 (Dual norm of ℓ_1 -norm is ℓ_∞ -norm).** *The dual norm of the ℓ_1 -norm is the*
 57 *ℓ_∞ -norm.*

58 *Proof.* We find the maximum of all $|z_i|, \forall i \in \{1, \dots, n\}$ and denote it by $|z_j|$. Then we let $x_j = 1$
 59 and $x_i = 0$ with $i \neq j$, which satisfies the requirement of $\|x\|_1 \leq 1$. Then,

$$\sup\{z^T x \mid \|x\|_1 \leq 1\} = |z_j| = \|z\|_\infty,$$

60 as desired. \square

61 **Proposition 5 (Dual norm of ℓ_p -norm is ℓ_q -norm).** *The dual norm of the ℓ_p -norm is the*
 62 *ℓ_q -norm, where $p, q \geq 1$ and $1/p + 1/q = 1$. That is,*

$$z^T x \leq \|x\|_p \|z\|_q.$$

63 This proposition is similar to Hölder inequality.

64 2 Quadratic mean, average mean, geometric mean, and har- 65 monic mean

66 The content in this section is largely taken from Xicheng Peng et al. Exploring Inequalities. 2016. In
 67 this section, we give the relationships between quadratic mean (QM), average mean (AM), geometric
 68 mean (GM), and harmonic mean (HM). The result is $QM \geq AM \geq GM \geq HM$.

69 **Theorem 6.** *For any positive real number a_1, a_2, \dots, a_n , we have the following inequalities*

$$\underbrace{\frac{1}{\frac{1}{a_1} + \frac{1}{a_1} + \dots + \frac{1}{a_n}}}_{\text{Harmonic Mean}} \leq \underbrace{\sqrt[n]{a_1 a_2 \dots a_n}}_{\text{Geometric Mean}} \leq \underbrace{\frac{a_1 + a_2 + \dots + a_n}{n}}_{\text{Arithmetic Mean}} \leq \underbrace{\sqrt{\frac{a_1^2 + a_2^2 + \dots + a_n^2}{n}}}_{\text{Quadratic Mean}} \quad (1)$$

70 where the equalities hold if and only if $a_1 = a_2 = \dots = a_n$. This inequality chain is also denoted as
 71 $H(n) \leq G(n) \leq A(n) \leq Q(n)$.

72 We will provide two proofs. The first proof employs Jensen's inequality and is simpler compared
 73 with the second proof.

⁷⁴ *Proof.* Let $f(x) = -\ln x$, then $f''(x) = \frac{1}{x^2} > 0$ which implies that f is convex. For all $a_i > 0$, by
⁷⁵ Jensen's inequality,

$$\begin{aligned} f\left(\frac{\frac{1}{a_1} + \frac{1}{a_2} + \cdots + \frac{1}{a_n}}{n}\right) &\leq \frac{1}{n} \left(f\left(\frac{1}{a_1}\right) + f\left(\frac{1}{a_2}\right) + \cdots + f\left(\frac{1}{a_n}\right) \right) \\ -\ln\left(\frac{\frac{1}{a_1} + \frac{1}{a_2} + \cdots + \frac{1}{a_n}}{n}\right) &\leq \frac{1}{n} \left(-\ln\left(\frac{1}{a_1}\right) - \ln\left(\frac{1}{a_2}\right) + \cdots - \ln\left(\frac{1}{a_n}\right) \right) \\ \ln\left(\frac{n}{\frac{1}{a_1} + \frac{1}{a_2} + \cdots + \frac{1}{a_n}}\right) &\leq \frac{1}{n} (\ln a_1 + \ln a_2 + \cdots + \ln a_n) \\ \ln\left(\frac{n}{\frac{1}{a_1} + \frac{1}{a_2} + \cdots + \frac{1}{a_n}}\right) &\leq \ln \sqrt[n]{a_1 a_2 \cdots a_n} \end{aligned}$$

⁷⁶ Since \ln is increasing,

$$\frac{n}{\frac{1}{a_1} + \frac{1}{a_2} + \cdots + \frac{1}{a_n}} \leq \sqrt[n]{a_1 a_2 \cdots a_n}$$

⁷⁷ where the equality holds if and only if $a_1 = a_2 = \cdots = a_n$. The “if” part is obvious. For the “only if”
⁷⁸ part, suppose the equality holds with $a_1 \neq a_2 = a_3 = \cdots = a_n$, then let $a_1 = k a_2$ ($k > 0$) we have

$$\frac{n}{(n-1 + \frac{1}{k}) \frac{1}{a_2}} = a_2 \sqrt[n]{k} \iff \frac{n}{(n-1 + \frac{1}{k})} = \sqrt[n]{k} \iff n = \sqrt[n]{k} (n-1 + \frac{1}{k})$$

⁷⁹ Let $x = \sqrt[n]{k}$ and it is clear that $x > 0$. Then

$$n x^{n-1} = (n-1) x^n + 1 \iff (n-1) x^n - n x^{n-1} + 1 = 0$$

⁸⁰

$$\begin{aligned} (n-1) x^n - n x^{n-1} + 1 &= (n-1) x^n - (n-1) x^{n-1} - x^{n-1} + 1 \\ &= (n-1) x^{n-1} (x-1) - (x^{n-1} - 1) \\ &= (n-1) x^{n-1} (x-1) - (x-1)(x^{n-2} + x^{n-3} + \cdots + 1) \\ &= (x-1)[(n-1) x^{n-1} - (x^{n-2} + x^{n-3} + \cdots + 1)] \\ &= (x-1)[(n-1) x^{n-1} - (n-1) x^{n-2} + (n-2) x^{n-2} - x^{n-3} - \cdots - 1] \\ &= (x-1)[(n-1) x^{n-2} (x-1) + (n-2) x^{n-2} - x^{n-3} - \cdots - 1] \\ &= (x-1)[(n-1) x^{n-2} (x-1) + (n-2) x^{n-2} - (n-2) x^{n-3} + (n-3) x^{n-3} - \cdots - 1] \\ &= (x-1)[(n-1) x^{n-2} (x-1) + (n-2) x^{n-3} (x-1) + (n-3) x^{n-3} - \cdots - 1] \\ &= (x-1)[(n-1) x^{n-2} (x-1) + (n-2) x^{n-3} (x-1) + (n-3) x^{n-4} (x-1) + \cdots + (x-1)] \\ &= (x-1)^2 [(n-1) x^{n-2} + (n-2) x^{n-3} + (n-3) x^{n-4} + \cdots + 1] \end{aligned}$$

⁸¹ Due to the fact that any polynomial that has positive coefficients cannot have roots on the nonnegative
⁸² real axis¹, 1 is the only roots which contradicts the supposition that $a_1 \neq a_2$. Hence, $a_1 = a_2 = \cdots =$
⁸³ a_n is the necessary and sufficient condition for the equality to hold.

⁸⁴ Now we show $\text{GM} \leq \text{AM}$,

$$\begin{aligned} f\left(\frac{a_1 + a_2 + \cdots + a_n}{n}\right) &\leq \frac{1}{n} (f(a_1) + f(a_2) + \cdots + f(a_n)) \\ -\ln\left(\frac{a_1 + a_2 + \cdots + a_n}{n}\right) &\leq \frac{1}{n} (-\ln(a_1) - \ln(a_1) + \cdots - \ln(a_1)) \end{aligned}$$

¹<https://mtns2018.hkust.edu.hk/media/files/0073.pdf>. Besides that paper, we can also prove this fact via contradiction. Suppose this kind of polynomial $P(x)$ has some positive real roots, say $P(x_0) = 0$, then each term of $P(x_0)$ is positive which leads to $P(x_0) > 0$, a contradiction. Thus, x_0 does not exist.

$$\begin{aligned}\ln\left(\frac{a_1 + a_2 + \cdots + a_n}{n}\right) &\geq \frac{1}{n}(\ln a_1 + \ln a_2 + \cdots + \ln a_n) \\ \ln\left(\frac{a_1 + a_2 + \cdots + a_n}{n}\right) &\geq \ln \sqrt[n]{a_1 a_2 \cdots a_n} \\ \frac{a_1 + a_2 + \cdots + a_n}{n} &\geq \sqrt[n]{a_1 a_2 \cdots a_n}\end{aligned}$$

85 Let $g(x) = x^2$, then $g''(x) = 2 > 0$ which indicates that g is convex.

86 Now we show AM \leq QM. By Jensen's inequality, we have

$$\begin{aligned}g\left(\frac{a_1 + a_2 + \cdots + a_n}{n}\right) &\leq \frac{1}{n}(g(a_1) + g(a_2) + \cdots + g(a_n)) \\ \left(\frac{a_1 + a_2 + \cdots + a_n}{n}\right)^2 &\leq \frac{1}{n}(a_1^2 + a_2^2 + \cdots + a_n^2) \\ \frac{a_1 + a_2 + \cdots + a_n}{n} &\leq \sqrt{\frac{a_1^2 + a_2^2 + \cdots + a_n^2}{n}}\end{aligned}$$

87 This completes our proof. □

88 We first introduce a lemma for the second proof.

89 **Lemma 7.** *If $a_i > 0, i = 1, 2, \dots, n$, and $a_1 a_2 \cdots a_n = 1$, then $a_1 + a_2 + \cdots + a_n \geq n$ where the*
90 *equality holds iff $a_1 = a_2 = \cdots = a_n = 1$.*

91 *Proof.* Let $x_i = \ln a_i, i = 1, 2, \dots, n$.

$$a_1 a_2 \cdots a_n = 1 \iff \ln a_1 + \ln a_2 + \cdots + \ln a_n = 0 \iff x_1 + x_2 + \cdots + x_n = 0$$

92 Since $e^x \geq x + 1$,

$$a_1 + a_2 + \cdots + a_n = e^{x_1} + e^{x_2} + \cdots + e^{x_n} \geq (x_1 + 1) + (x_2 + 1) + \cdots + (x_n + 1) = n$$

93 where the equality holds iff $x_i = 0, \forall i = 1, 2, \dots, n$, i.e., $a_i = 1, \forall i = 1, 2, \dots, n$ due to the fact that
94 $e^x = x + 1$ iff $x = 0$. Thus, $a_1 + a_2 + \cdots + a_n = n$ iff $a_1 = a_2 = \cdots = a_n = 1$. □

95 Next, we prove this inequality chain using Lemma 7.

96 *Proof.* 1. $H(n) \leq G(n)$

$$\frac{n}{\frac{1}{a_1} + \frac{1}{a_2} + \cdots + \frac{1}{a_n}} \leq \sqrt[n]{a_1 a_2 \cdots a_n} \iff \frac{\sqrt[n]{a_1 a_2 \cdots a_n}}{a_1} + \frac{\sqrt[n]{a_1 a_2 \cdots a_n}}{a_2} + \cdots + \frac{\sqrt[n]{a_1 a_2 \cdots a_n}}{a_n} \geq n$$

97 We observe that $\frac{\sqrt[n]{a_1 a_2 \cdots a_n}}{a_1} \cdot \frac{\sqrt[n]{a_1 a_2 \cdots a_n}}{a_2} \cdots \frac{\sqrt[n]{a_1 a_2 \cdots a_n}}{a_n} = 1$. By Lemma 7, the
98 above holds. The equality holds iff $\frac{\sqrt[n]{a_1 a_2 \cdots a_n}}{a_1} = \frac{\sqrt[n]{a_1 a_2 \cdots a_n}}{a_2} = \cdots = \frac{\sqrt[n]{a_1 a_2 \cdots a_n}}{a_n}$, i.e.,
99 $a_1 = a_2 = \cdots = a_n$.

100 2. $G(n) \leq A(n)$

$$\sqrt[n]{a_1 a_2 \cdots a_n} \leq \frac{a_1 + a_2 + \cdots + a_n}{n} \iff \frac{a_1}{\sqrt[n]{a_1 a_2 \cdots a_n}} + \frac{a_2}{\sqrt[n]{a_1 a_2 \cdots a_n}} + \cdots + \frac{a_n}{\sqrt[n]{a_1 a_2 \cdots a_n}} \geq n$$

101 We observe that $\frac{a_1}{\sqrt[n]{a_1 a_2 \cdots a_n}} \cdot \frac{a_2}{\sqrt[n]{a_1 a_2 \cdots a_n}} \cdots \frac{a_n}{\sqrt[n]{a_1 a_2 \cdots a_n}} = 1$. By Lemma 7, the
102 above holds. The equality holds iff $\frac{a_1}{\sqrt[n]{a_1 a_2 \cdots a_n}} = \frac{a_2}{\sqrt[n]{a_1 a_2 \cdots a_n}} = \cdots = \frac{a_n}{\sqrt[n]{a_1 a_2 \cdots a_n}}$, i.e.,
103 $a_1 = a_2 = \cdots = a_n$.

3. $A(n) \leq Q(n)$. Let $c = \frac{a_1+a_2+\dots+a_n}{n}$ and $a_i = c + \alpha_i, \forall i = 1$. Then

$$\begin{aligned} a_1 + a_2 + \dots + a_n &= nc + \alpha_1 + \alpha_2 + \dots + \alpha_n \\ &= a_1 + a_2 + \dots + a_n + (\alpha_1 + \alpha_2 + \dots + \alpha_n) \end{aligned}$$

So, $\alpha_1 + \alpha_2 + \dots + \alpha_n = 0$.

$$\begin{aligned} \sqrt{\frac{a_1^2 + a_2^2 + \dots + a_n^2}{n}} &= \sqrt{\frac{(c + \alpha_1)^2 + (c + \alpha_2)^2 + \dots + (c + \alpha_n)^2}{n}} \\ &= \sqrt{\frac{nc^2 + 2c(\alpha_1 + \alpha_2 + \dots + \alpha_n) + \alpha_1^2 + \alpha_2^2 + \dots + \alpha_n^2}{n}} \\ &= \sqrt{\frac{nc^2 + \alpha_1^2 + \alpha_2^2 + \dots + \alpha_n^2}{n}} \\ &= \sqrt{c^2 + \frac{\alpha_1^2 + \alpha_2^2 + \dots + \alpha_n^2}{n}} \\ &\geq \sqrt{c^2} = c \end{aligned}$$

104 where the third equality follows from $\alpha_1 + \alpha_2 + \dots + \alpha_n = 0$ and $c = \frac{a_1+a_2+\dots+a_n}{n}$.
 105 Lemma 7 is not involved with the proof of $A(n) \leq Q(n)$. □

106 *Note.* $AM \geq GM \geq HM$ can be proved using induction; see Theorem 1.2.2 of Jixiu Chen et al. *Mathe-*
 107 *matical Analysis*, third edition.

108 2.1 An inequality about the difference between AM and GM

109 **Proposition 8.** *Given $a, b, c \in \mathbf{R}_+$, show $3(\frac{a+b+c}{3} - \sqrt[3]{abc}) \geq 2(\frac{a+b}{2} - \sqrt{ab})$.*

110 *Proof.* After simple algebra, it suffices to show

$$c + 2\sqrt{ab} \geq 3\sqrt[3]{abc}$$

111 If we consider $2\sqrt{ab}$ as $\sqrt{ab} + \sqrt{ab}$ on the LHS, we can use AM-GM to get

$$c + \sqrt{ab} + \sqrt{ab} \geq 3\sqrt[3]{\sqrt{ab} \cdot \sqrt{ab} \cdot c} = 3\sqrt[3]{abc}.$$

112 This completes our proof. □

113 This result can be easily generalized to the general case, namely,

$$(n+1)\left(\frac{a_1^2 + a_2^2 + \dots + a_{n+1}^2}{n+1} - \sqrt[n+1]{a_1 a_2 \dots a_{n+1}}\right) \geq n\left(\frac{a_1^2 + a_2^2 + \dots + a_n^2}{n} - \sqrt[n]{a_1 a_2 \dots a_n}\right).$$

114 2.2 Power mean inequality

115 Refer to Page 106 of Xicheng Peng et al. *Exploring Inequalities*. 2016. This will be done later.

116 2.3 Weighted power mean inequality

117 Refer to Page 108 of Xicheng Peng et al. *Exploring Inequalities*. 2016. This will be done later.

2.4 Applications

Proposition 9. Given positive reals a, b, c and $a + b + c = 1$, show that $ab + bc + ca \leq \frac{1}{3}$.

Proof. Since the geometric mean is not less than the average mean, we have

$$\sqrt{\frac{a^2 + b^2 + c^2}{3}} \geq \frac{a + b + c}{3} = \frac{1}{3} \iff a^2 + b^2 + c^2 \geq \frac{1}{3}.$$

Furthermore, we have

$$ab + bc + ca = \frac{(a + b + c)^2 - (a^2 + b^2 + c^2)}{2} = \frac{1 - (a^2 + b^2 + c^2)}{2} \leq \frac{1 - 1/3}{2} = \frac{1}{3},$$

as desired. \square

3 Young inequality

Theorem 10. Given $x, y \geq 0$, $p, q \geq 1$, and $1/p + 1/q = 1$, the inequality

$$xy \leq \frac{1}{p}x^p + \frac{1}{q}y^q,$$

where the equality holds if and only if $x^p = y^q$.

Proof. The claim is obvious for the case when either $x = 0$ or $y = 0$. When x and y are positive reals, we let $f(t) = e^t$, then $f'' > 0$. So f is convex. By the Jensen's inequality,

$$f\left(\frac{1}{p} \ln x^p + \frac{1}{q} \ln y^q\right) \leq \frac{1}{p} f(\ln x^p) + \frac{1}{q} f(\ln y^q)$$

$$\Downarrow$$

$$e^{\frac{1}{p} \ln x^p + \frac{1}{q} \ln y^q} \leq \frac{1}{p} e^{\ln x^p} + \frac{1}{q} e^{\ln y^q} \iff e^{\frac{1}{p} \ln x^p} e^{\frac{1}{q} \ln y^q} \leq \frac{1}{p} x^p + \frac{1}{q} y^q \iff xy \leq \frac{1}{p} x^p + \frac{1}{q} y^q,$$

as desired. The equality follows from the condition for the equality of Jensen's inequality to hold with any convex function. \square

4 Hölder inequality

A classic result concerning p -norms is the **Hölder inequality in inner-product form**:

Theorem 11 (Hölder inequality). For any $\mathbf{x}, \mathbf{y} \in \mathbf{R}^n$ and $p, q > 1$ satisfying $1/p + 1/q = 1$, the following

$$|\mathbf{x}^T \mathbf{y}| \leq \|\mathbf{x}\|_p \|\mathbf{y}\|_q, \quad \frac{1}{p} + \frac{1}{q} = 1$$

holds. The equality holds if and only if $\frac{|x_1|^p}{|y_1|^q} = \frac{|x_2|^p}{|y_2|^q} = \dots = \frac{|x_n|^p}{|y_n|^q}$ holds.

We provide two proofs here. The first proof employs Young inequality and the second proof makes use of the fact that the dual norm of p -norm is q -norm with $p, q \geq 0$ and $1/p + 1/q = 1$.

Proof. The claim is trivial either $\mathbf{x} = \mathbf{0}$ or $\mathbf{y} = \mathbf{0}$. Suppose that $\mathbf{x} \neq \mathbf{0}$ or $\mathbf{y} \neq \mathbf{0}$. For any $i \in \{1, 2, \dots, n\}$, letting $s = \frac{|x_i|}{(\sum_{i=1}^n |x_i|^p)^{1/p}}$ and $t = \frac{|y_i|}{(\sum_{i=1}^n |y_i|^q)^{1/q}}$, by Young's inequality, we have

$$st = \frac{|x_i|}{(\sum_{i=1}^n |x_i|^p)^{1/p}} \frac{|y_i|}{(\sum_{i=1}^n |y_i|^q)^{1/q}} \leq \frac{1}{p} \left(\frac{|x_i|}{(\sum_{i=1}^n |x_i|^p)^{1/p}} \right)^p + \frac{1}{q} \left(\frac{|y_i|}{(\sum_{i=1}^n |y_i|^q)^{1/q}} \right)^q$$

$$\begin{aligned} & \Updownarrow \\ & \frac{|x_i y_i|}{(\sum_{i=1}^n |x_i|^p)^{1/p} (\sum_{i=1}^n |y_i|^q)^{1/q}} \leq \frac{1}{p} \frac{|x_i|^p}{\sum_{i=1}^n |x_i|^p} + \frac{1}{q} \frac{|y_i|^q}{\sum_{i=1}^n |y_i|^q} \end{aligned}$$

Summing up over i ,

$$\begin{aligned} & \frac{\sum_{i=1}^n |x_i y_i|}{(\sum_{i=1}^n |x_i|^p)^{1/p} (\sum_{i=1}^n |y_i|^q)^{1/q}} \leq \frac{1}{p} \frac{\sum_{i=1}^n |x_i|^p}{\sum_{i=1}^n |x_i|^p} + \frac{1}{q} \frac{\sum_{i=1}^n |y_i|^q}{\sum_{i=1}^n |y_i|^q} \\ & \Updownarrow \\ & \frac{\sum_{i=1}^n |x_i y_i|}{(\sum_{i=1}^n |x_i|^p)^{1/p} (\sum_{i=1}^n |y_i|^q)^{1/q}} \leq \frac{1}{p} + \frac{1}{q} = 1 \\ & \Updownarrow \\ & |\mathbf{x}^T \mathbf{y}| \leq \sum_{i=1}^n |x_i y_i| \leq \left(\sum_{i=1}^n |x_i|^p \right)^{1/p} \left(\sum_{i=1}^n |y_i|^q \right)^{1/q}, \end{aligned} \tag{2}$$

as desired. \square

Now we present the second proof.

Proof. Recall the definition of dual norm,

$$\|x\|_p = \max_{\|z\|_q \leq 1} x^T z.$$

Thus, $x^T z \leq \|x\|_p$ holds for any z satisfying $\|z\|_q \leq 1$, including $\|z\|_q = 1$. When $\|z\|_q = 1$, we have

$$x^T z \leq \|x\|_p \|z\|_q$$

Now let $z = ty$ with $t > 0$. Thus, we have

$$x^T (ty) \leq \|x\|_p \|ty\|_q \iff x^T y \leq \|x\|_p \|y\|_q$$

where $\|y\|_q = \|z/t\|_q = 1/t \|z\|_q = 1/t > 0$. This completes the proof. \square

The key part of the above proof is using the dual representation of the ℓ_p norm, namely, $\|x\|_p = \max_{\|z\|_q \leq 1} x^T z$.

4.1 Generalized Hölder inequality

Setting $\lambda_1 = 1/p$ and $\lambda_2 = 1/q$ in (2) yields

$$\left(\sum_{i=1}^n |x_i|^{1/\lambda_1} \right)^{\lambda_1} \left(\sum_{i=1}^n |y_i|^{1/\lambda_2} \right)^{\lambda_2} \geq \sum_{i=1}^n (|x_i|^{1/\lambda_1})^{\lambda_1} (|y_i|^{1/\lambda_2})^{\lambda_2},$$

where $0 < \lambda_1, \lambda_2 < 1$ and $\lambda_1 + \lambda_2 = 1$. Continuing on this notational trick by setting $a_{i1} = |x_i|^{1/\lambda_1}$ and $a_{i2} = |y_i|^{1/\lambda_2}$ for any $i \in \{1, 2, \dots, n\}$ gives

$$\left(\sum_{i=1}^n a_{i1} \right)^{\lambda_1} \left(\sum_{i=1}^n a_{i2} \right)^{\lambda_2} \geq \sum_{i=1}^n (a_{i1})^{\lambda_1} (a_{i2})^{\lambda_2}.$$

where $a_{i1}, a_{i2} \geq 0$ for any $i \in \{1, 2, \dots, n\}$. This form is beautiful. If it can be generalized to λ_m ($m > 2$), it will be wonderful. Yes, it is. We formalize it into the following theorem.

Theorem 12 (Generalized Hölder inequality). Given a matrix $A \in \mathbf{R}^{n \times m}$ with all nonnegative entries a_{ij} and $0 < \lambda_j < 1$ satisfying $\lambda_1 + \lambda_2 + \dots + \lambda_m = 1$, it holds that

$$\begin{aligned} & \left(\sum_{i=1}^n a_{i1} \right)^{\lambda_1} \left(\sum_{i=1}^n a_{i2} \right)^{\lambda_2} \dots \left(\sum_{i=1}^n a_{im} \right)^{\lambda_m} \\ & \geq a_{11}^{\lambda_1} a_{12}^{\lambda_2} \dots a_{1m}^{\lambda_m} + a_{21}^{\lambda_1} a_{22}^{\lambda_2} \dots a_{2m}^{\lambda_m} + \dots + a_{n1}^{\lambda_1} a_{n2}^{\lambda_2} \dots a_{nm}^{\lambda_m} \\ & = \sum_{i=1}^n a_{i1}^{\lambda_1} a_{i2}^{\lambda_2} \dots a_{im}^{\lambda_m} \end{aligned}$$

Compactly,

$$\prod_{j=1}^m \left(\sum_{i=1}^n a_{ij} \right)^{\lambda_j} \geq \sum_{i=1}^n \prod_{j=1}^m a_{ij}^{\lambda_j}.$$

Proof. We prove this generalized Hölder inequality by mathematical induction. When $m = 2$, it is exactly Theorem 11, i.e., the canonical Hölder inequality. Suppose the claim holds when $m = k$. When $m = k + 1$, let $\lambda_1 + \lambda_2 + \dots + \lambda_k = s$ and denote $t_i = \lambda_i/s$, $i = 1, 2, \dots, k$. Then

$$\begin{aligned} \sum_{i=1}^n a_{i1}^{\lambda_1} a_{i2}^{\lambda_2} \dots a_{im}^{\lambda_m} &= \sum_{i=1}^n (a_{i1}^{t_1} a_{i2}^{t_2} \dots a_{ik}^{t_k})^s a_{i,k+1}^{\lambda_{k+1}} \\ &\leq \left(\sum_{i=1}^n ((a_{i1}^{t_1} a_{i2}^{t_2} \dots a_{ik}^{t_k})^s)^{\frac{1}{s}} \right)^s \left(\sum_{i=1}^n (a_{i,k+1}^{\lambda_{k+1}})^{\frac{1}{\lambda_{k+1}}} \right)^{\lambda_{k+1}} \\ &= \left(\sum_{i=1}^n a_{i1}^{t_1} a_{i2}^{t_2} \dots a_{ik}^{t_k} \right)^s \left(\sum_{i=1}^n a_{i,k+1}^{\lambda_{k+1}} \right)^{\lambda_{k+1}} \\ &\leq \left(\left(\sum_{i=1}^n a_{i1} \right)^{t_1} \left(\sum_{i=1}^n a_{i2} \right)^{t_2} \dots \left(\sum_{i=1}^n a_{ik} \right)^{t_k} \right)^s \left(\sum_{i=1}^n a_{i,k+1}^{\lambda_{k+1}} \right)^{\lambda_{k+1}} \\ &= \left(\sum_{i=1}^n a_{i1} \right)^{\lambda_1} \left(\sum_{i=1}^n a_{i2} \right)^{\lambda_2} \dots \left(\sum_{i=1}^n a_{ik} \right)^{\lambda_k} \left(\sum_{i=1}^n a_{i,k+1}^{\lambda_{k+1}} \right)^{\lambda_{k+1}} \end{aligned}$$

where the first inequality and the second inequality follow from the classic Hölder inequality and the induction assumption, respectively. This completes the proof. \square

The generalized Hölder inequality can also be applied to the case where $\lambda_1 + \lambda_2 + \dots + \lambda_m < 1$. For this, we have the following corollary.

Corollary 13. Given a matrix $A \in \mathbf{R}^{n \times m}$ with all nonnegative entries a_{ij} and $0 < \lambda_j < 1$ satisfying $\lambda_1 + \lambda_2 + \dots + \lambda_m = r < 1$, it holds that

$$\begin{aligned} & \left(\sum_{i=1}^n a_{i1} \right)^{\lambda_1} \left(\sum_{i=1}^n a_{i2} \right)^{\lambda_2} \dots \left(\sum_{i=1}^n a_{im} \right)^{\lambda_m} \\ & \geq n^{r-1} a_{11}^{\lambda_1} a_{12}^{\lambda_2} \dots a_{1m}^{\lambda_m} + n^{r-1} a_{21}^{\lambda_1} a_{22}^{\lambda_2} \dots a_{2m}^{\lambda_m} + \dots + n^{r-1} a_{n1}^{\lambda_1} a_{n2}^{\lambda_2} \dots a_{nm}^{\lambda_m} \\ & = n^{r-1} \sum_{i=1}^n a_{i1}^{\lambda_1} a_{i2}^{\lambda_2} \dots a_{im}^{\lambda_m} \end{aligned}$$

Compactly,

$$\prod_{j=1}^m \left(\sum_{i=1}^n a_{ij} \right)^{\lambda_j} \geq n^{r-1} \sum_{i=1}^n \prod_{j=1}^m a_{ij}^{\lambda_j}.$$

163 *Proof.* By the generalized Hölder inequality, letting $\alpha = 1 - r$ yields

$$\begin{aligned} & (1 + 1 + \dots + 1)^\alpha \left(\sum_{i=1}^n a_{i1} \right)^{\lambda_1} \left(\sum_{i=1}^n a_{i2} \right)^{\lambda_2} \dots \left(\sum_{i=1}^n a_{im} \right)^{\lambda_m} \\ & \geq a_{11}^{\lambda_1} a_{12}^{\lambda_2} \dots a_{1m}^{\lambda_m} + a_{21}^{\lambda_1} a_{22}^{\lambda_2} \dots a_{2m}^{\lambda_m} + \dots + a_{n1}^{\lambda_1} a_{n2}^{\lambda_2} \dots a_{nm}^{\lambda_m} \\ & = \sum_{i=1}^n a_{i1}^{\lambda_1} a_{i2}^{\lambda_2} \dots a_{im}^{\lambda_m}. \end{aligned}$$

164 Equivalently,

$$\left(\sum_{i=1}^n a_{i1} \right)^{\lambda_1} \left(\sum_{i=1}^n a_{i2} \right)^{\lambda_2} \dots \left(\sum_{i=1}^n a_{im} \right)^{\lambda_m} \geq n^{r-1} \sum_{i=1}^n a_{i1}^{\lambda_1} a_{i2}^{\lambda_2} \dots a_{im}^{\lambda_m}$$

165

□

166 4.2 Cauchy-Schwarz inequality

A very important special case of Hölder inequality is **Cauchy-Schwarz inequality**:

$$|x^T y| \leq \|x\|_2 \|y\|_2.$$

167 which can also be expressed as $(x^T y)^2 \leq \|x\|_2^2 \|y\|_2^2$. Here, $x = (x_1, x_2, \dots, x_n)$ and $y = (y_1, y_2, \dots, y_n)$.

168 Thus,

$$\begin{aligned} (x_1 y_1 + x_2 y_2 + \dots + x_n y_n)^2 & \leq (x_1^2 + x_2^2 + \dots + x_n^2)(y_1^2 + y_2^2 + \dots + y_n^2) \\ \left(\sum_{i=1}^n x_i y_i \right)^2 & = \sum_{i=1}^n x_i^2 \sum_{i=1}^n y_i^2 \end{aligned}$$

169 4.3 Two variants of Cauchy-Schwarz inequality

170 Given $a_i \in \mathbf{R}$ and $b_i > 0 (i = 1, 2, \dots, n)$, let $y_i = \sqrt{b_i}$ and $x_i = \frac{a_i}{\sqrt{b_i}}$. Then we have an important
171 **variant** of Cauchy-Schwarz inequality.

$$\left(\sum_{i=1}^n \frac{a_i}{\sqrt{b_i}} \sqrt{b_i} \right)^2 = \left(\sum_{i=1}^n a_i \right)^2 \leq \left(\sum_{i=1}^n \frac{a_i^2}{b_i} \right) \sum_{i=1}^n b_i \iff \sum_{i=1}^n \frac{a_i^2}{b_i} \geq \frac{(\sum_{i=1}^n a_i)^2}{\sum_{i=1}^n b_i}.$$

172 For $a_i > 0$ and $b_i > 0 (i = 1, 2, \dots, n)$, letting $x_i = \sqrt{\frac{a_i}{b_i}}$ and $y_i = \sqrt{a_i b_i}$ gives

$$\sum_{i=1}^n x_i^2 \cdot \sum_{i=1}^n y_i^2 \geq \left(\sum_{i=1}^n x_i y_i \right)^2 \iff \sum_{i=1}^n \frac{a_i}{b_i} \cdot \sum_{i=1}^n a_i b_i \geq \left(\sum_{i=1}^n a_i \right)^2 \iff \sum_{i=1}^n \frac{a_i}{b_i} \geq \frac{(\sum_{i=1}^n a_i)^2}{\sum_{i=1}^n a_i b_i}.$$

173 5 Minkowski inequality

174 **Theorem 14 (Minkowski inequality).** Given $x_i, y_i \geq 0, i = 1, 2, \dots, n$ and $p \geq 1$, the following

$$\left(\sum_{i=1}^n |x_i + y_i|^p \right)^{\frac{1}{p}} \leq \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} + \left(\sum_{i=1}^n |y_i|^p \right)^{\frac{1}{p}}$$

175 holds. By the definition of p -norm (see section 1.2), the equivalent vector form

$$\|\mathbf{x} + \mathbf{y}\|_p \leq \|\mathbf{x}\|_p + \|\mathbf{y}\|_p$$

176 holds.

177 Since the following proof employs Hölder inequality in which $1/p + 1/q = 1$ is required and
 178 $q \rightarrow +\infty$ as $p \rightarrow 1$, we need to consider the case of $p = 1$ separately. The triangle inequality is
 179 enough to show the case of $p = 1$.

180 *Proof.* We observe that when either $x_i = 0$ or $y_i = 0$, or $x_i + y_i = 0, i = 1, 2, \dots, n$, the result is
 181 trivial. Now we show the case when $\sum_{i=1}^n x_i + y_i \neq 0$.

182 We first prove the case of $p = 1$. By the triangle inequality, we have

$$\sum_{i=1}^n |x_i + y_i| \leq \sum_{i=1}^n (|x_i| + |y_i|) = \sum_{i=1}^n |x_i| + \sum_{i=1}^n |y_i| \iff \|\mathbf{x} + \mathbf{y}\|_1 \leq \|\mathbf{x}\|_1 + \|\mathbf{y}\|_1.$$

183 Now we turn to the case of $p > 1$.

$$\sum_{i=1}^n |x_i + y_i|^p = \sum_{i=1}^n |x_i + y_i| |x_i + y_i|^{p-1} \leq \sum_{i=1}^n |x_i| |x_i + y_i|^{p-1} + \sum_{i=1}^n |y_i| |x_i + y_i|^{p-1},$$

184 where the last inequality follows from the triangle inequality.

185 Applying Hölder inequality on both terms of the RHS gives

$$\begin{aligned} \sum_{i=1}^n |x_i| |x_i + y_i|^{p-1} &\leq \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} \left(\sum_{i=1}^n |x_i + y_i|^{q(p-1)} \right)^{\frac{1}{q}} = \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} \left(\sum_{i=1}^n |x_i + y_i|^p \right)^{\frac{1}{q}} \\ \sum_{i=1}^n |y_i| |x_i + y_i|^{p-1} &\leq \left(\sum_{i=1}^n |y_i|^p \right)^{\frac{1}{p}} \left(\sum_{i=1}^n |x_i + y_i|^{q(p-1)} \right)^{\frac{1}{q}} = \left(\sum_{i=1}^n |y_i|^p \right)^{\frac{1}{p}} \left(\sum_{i=1}^n |x_i + y_i|^p \right)^{\frac{1}{q}} \end{aligned}$$

186 where the equalities follow from the fact that $q(p-1) = p$ due to Hölder inequality's condition, i.e.,
 187 $1/p + 1/q = 1$. Adding the above two inequalities up yields,

$$\begin{aligned} \sum_{i=1}^n |x_i + y_i|^p &\leq \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} \left(\sum_{i=1}^n |x_i + y_i|^p \right)^{\frac{1}{q}} + \left(\sum_{i=1}^n |y_i|^p \right)^{\frac{1}{p}} \left(\sum_{i=1}^n |x_i + y_i|^p \right)^{\frac{1}{q}} \\ &\iff \\ &\frac{\sum_{i=1}^n |x_i + y_i|^p}{\left(\sum_{i=1}^n |x_i + y_i|^p \right)^{\frac{1}{q}}} \leq \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} + \left(\sum_{i=1}^n |y_i|^p \right)^{\frac{1}{p}} \\ &\iff \\ \left(\sum_{i=1}^n |x_i + y_i|^p \right)^{1-\frac{1}{q}} &\leq \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} + \left(\sum_{i=1}^n |y_i|^p \right)^{\frac{1}{p}} \iff \left(\sum_{i=1}^n |x_i + y_i|^p \right)^{\frac{1}{p}} \leq \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} + \left(\sum_{i=1}^n |y_i|^p \right)^{\frac{1}{p}}, \end{aligned}$$

188 as desired. □

189 6 Rearrangement inequalities

190 For two sequences $a_1 \leq a_2 \leq \dots \leq a_n$ and $b_1 \leq b_2 \leq \dots \leq b_n$ with reals a_i and b_i , the following
 191 inequalities

$$\underbrace{a_1 b_n + a_2 b_{n-1} + \dots + a_n b_1}_{\text{reverse sum}} \leq \underbrace{a_1 b_{\pi(1)} + a_2 b_{\pi(2)} + \dots + a_n b_{\pi(n)}}_{\text{disordered sum}} \leq \underbrace{a_1 b_1 + a_2 b_2 + \dots + a_n b_n}_{\text{sequential sum}}$$

192 hold, where $\pi(1), \pi(2), \dots, \pi(n)$ is any permutation of $1, 2, \dots, n$. Simply speaking, **reverse sum** \leq
 193 **disordered sum** \leq **sequential sum**.

194 *Proof.* ² We want to prove that the identity permutation maximizes $a_1b_{\pi(1)} + a_2b_{\pi(2)} + \cdots + a_nb_{\pi(n)}$.
 195 Suppose for the sake of contradiction $\pi(i)$ is the smallest integer such that $\pi(i) \neq i$, then $\pi(i) = j (j > i)$
 196 (since $1, \dots, i-1$ have been assigned). Meanwhile, there exists a number $k > i$ such that $\pi(k) = i$ as
 197 some number must be assigned to i .

198 Now, since $i < j$, it follows that $b_i \leq b_j$. Likewise, since $i < k$, it follows that $a_i \leq a_k$. Thus,

$$(a_k - a_i)(b_j - b_i) \geq 0 \implies a_kb_j + a_ib_i \geq a_ib_j + a_kb_i,$$

199 which demonstrates that the sum $a_1b_{\pi(1)} + a_2b_{\pi(2)} + \cdots + a_nb_{\pi(n)}$ is not decreased by changing
 200 $\pi(j) = i$ and $\pi(k) = i$ to $\pi(i) = i$ and $\pi(k) = j$. This implies the identity permutation gives the
 201 maximum possible value of the sum $a_1b_{\pi(1)} + a_2b_{\pi(2)} + \cdots + a_nb_{\pi(n)}$, as desired. Moreover, the reverse
 202 identity permutation gives the minimum possible value of the sum $a_1b_{\pi(1)} + a_2b_{\pi(2)} + \cdots + a_nb_{\pi(n)}$. \square

203 6.1 Chebyshev's inequality

204 For two sequences $a_1 \leq a_2 \leq \cdots \leq a_n$ and $b_1 \leq b_2 \leq \cdots \leq b_n$ with reals a_i and b_i , by the
 205 rearrangement inequalities, we always have the following inequalities

$$\begin{aligned} x_1y_n + x_2y_{n-1} + \cdots + x_ny_1 &\leq x_1y_1 + x_2y_2 + \cdots + x_ny_n \leq x_1y_1 + x_2y_2 + \cdots + x_ny_n \\ x_1y_n + x_2y_{n-1} + \cdots + x_ny_1 &\leq x_1y_2 + x_2y_3 + \cdots + x_ny_1 \leq x_1y_1 + x_2y_2 + \cdots + x_ny_n \\ x_1y_n + x_2y_{n-1} + \cdots + x_ny_1 &\leq x_1y_3 + x_2y_4 + \cdots + x_ny_2 \leq x_1y_1 + x_2y_2 + \cdots + x_ny_n \\ &\vdots \\ x_1y_n + x_2y_{n-1} + \cdots + x_ny_1 &\leq x_1y_n + x_2y_1 + \cdots + x_ny_{n-1} \leq x_1y_1 + x_2y_2 + \cdots + x_ny_n \end{aligned}$$

206 Summing up the above inequalities gives

$$\begin{aligned} n(x_1y_n + x_2y_{n-1} + \cdots + x_ny_1) &\leq (x_1 + x_2 + \cdots + x_n)(y_1 + y_2 + \cdots + y_n) \leq n(x_1y_1 + x_2y_2 + \cdots + x_ny_n) \\ \frac{1}{n}(x_1y_n + x_2y_{n-1} + \cdots + x_ny_1) &\leq \frac{1}{n^2}(x_1 + x_2 + \cdots + x_n)(y_1 + y_2 + \cdots + y_n) \leq \frac{1}{n}(x_1y_1 + x_2y_2 + \cdots + x_ny_n) \\ \frac{1}{n} \sum_{i=1}^n x_i y_{n+1-i} &\leq \frac{1}{n} \sum_{i=1}^n x_i \cdot \frac{1}{n} \sum_{i=1}^n y_i \leq \frac{1}{n} \sum_{i=1}^n x_i y_i \end{aligned}$$

207 which is called Chebyshev's inequality.

²<https://brilliant.org/wiki/rearrangement-inequality/>