

A proof for a generalization of the inequality from the 42nd International Mathematical Olympiad

Youming Zhao
Email: youming0.zhao@gmail.com

October 29, 2023

Abstract

In this paper, we present a proof for a generalization of the inequality from the 42nd International Mathematical Olympiad. The proved inequality relates to a sum involving square roots of fractions. It has various applications in mathematical analysis, optimization, or statistics. In the field of mathematical analysis, it can be used in the study of convergence. In terms of optimization, it may help establish bounds or relationships between the variables involved.

In this paper, we will present a proof for a generalization of the inequality from the 42nd International Mathematical Olympiad. The original inequality corresponds to the case when $n = 3$. We aim to prove

$$\begin{aligned} & \sqrt{\frac{x_1^{n-1}}{x_1^{n-1} + (n^2 - 1)x_2x_3 \cdots x_n}} + \sqrt{\frac{x_2^{n-1}}{x_2^{n-1} + (n^2 - 1)x_1x_3 \cdots x_n}} \\ & + \cdots + \sqrt{\frac{x_n^{n-1}}{x_n^{n-1} + (n^2 - 1)x_1x_2 \cdots x_{n-1}}} \geq 1 \end{aligned}$$

for all positive real numbers $x_i, i = 1, 2, \dots, n$ with $n \geq 2$.

We can observe that the left-hand side (LHS) of the inequality is homogeneous. Therefore, we can assume, without loss of generality, that $\sum_{i=1}^n x_i^p = 1$, without changing the value of the LHS of the inequality. To see this, suppose $\sum_{i=1}^n x_i^p = m$ where $m > 0$, let $\tilde{x}_i = x_i/m^{\frac{1}{p}}$ which ensures $\sum_{i=1}^n \tilde{x}_i^p = 1$. The LHS of the inequality can be expressed as $h(x_1, x_2, \dots, x_n) = \sum_{i=1}^n h_i(x_i)$, in which $h_i(x_i)$ is defined as

$$h_i(x_i) = \sqrt{\frac{x_i^{n-1}}{x_i^{n-1} + (n^2 - 1)x_1x_2 \cdots x_{i-1}x_{i+1} \cdots x_n}}$$

Now we show $h_i(\tilde{x}_i) = h_i(x_i)$.

$$\begin{aligned}
h_i(\tilde{x}_i) &= \sqrt{\frac{\left(\frac{x_i}{m^{1/p}}\right)^{n-1}}{\left(\frac{x_i}{m^{1/p}}\right)^{n-1} + (n^2 - 1) \cdot \frac{x_1}{m^{1/p}} \frac{x_2}{m^{1/p}} \cdots \frac{x_{i-1}}{m^{1/p}} \frac{x_{i+1}}{m^{1/p}} \cdots \frac{x_n}{m^{1/p}}}} \\
&= \sqrt{\frac{\frac{x_i^{n-1}}{m^{(n-1)/p}}}{\frac{x_i^{n-1}}{m^{(n-1)/p}} + \frac{(n^2-1)x_1x_2\cdots x_{i-1}x_{i+1}\cdots x_n}{m^{(n-1)/p}}}} \\
&= \sqrt{\frac{x_i^{n-1}}{x_i^{n-1} + (n^2 - 1)x_1x_2\cdots x_{i-1}x_{i+1}\cdots x_n}} = h_i(x_i)
\end{aligned}$$

which demonstrates that arbitrary combination of x_i can be transformed into \tilde{x}_i such that $\sum_{i=1}^n \tilde{x}_i = 1$ without changing the value of the left-hand side.

Consider the function $f(x) = \frac{1}{\sqrt{x}}$ with $x > 0$. Taking the first and second derivative, we find that $f'(x) = -\frac{1}{2}x^{-\frac{3}{2}} < 0$ and $f''(x) = \frac{3}{4}x^{-\frac{5}{2}} > 0$. Hence, $f(x)$ is a monotonically decreasing and convex function. Using the homogeneity of the inequality, we can suppose $\sum_{i=1}^n x_i^{\frac{n-1}{2}} = 1$ and then apply Jensen's inequality as follows.

$$\begin{aligned}
&\sqrt{\frac{x_1^{n-1}}{x_1^{n-1} + (n^2 - 1)x_2x_3\cdots x_n}} + \sqrt{\frac{x_2^{n-1}}{x_2^{n-1} + (n^2 - 1)x_1x_3\cdots x_n}} \\
&+ \cdots + \sqrt{\frac{x_n^{n-1}}{x_n^{n-1} + (n^2 - 1)x_1x_2\cdots x_{n-1}}} \\
&= x_1^{\frac{n-1}{2}} f(x_1^{n-1} + (n^2 - 1)x_2x_3\cdots x_n) + x_2^{\frac{n-1}{2}} f(x_2^{n-1} + (n^2 - 1)x_1x_3\cdots x_n) \\
&+ \cdots + x_n^{\frac{n-1}{2}} f(x_n^{n-1} + (n^2 - 1)x_1x_2\cdots x_{n-1}) \\
&\geq f\left(x_1^{\frac{n-1}{2}} (x_1^{n-1} + (n^2 - 1)x_2x_3\cdots x_n) + x_2^{\frac{n-1}{2}} (x_2^{n-1} + (n^2 - 1)x_1x_3\cdots x_n) \right. \\
&\quad \left. + \cdots + x_n^{\frac{n-1}{2}} (x_n^{n-1} + (n^2 - 1)x_1x_2\cdots x_{n-1})\right) \\
&= f\left(x_1^{\frac{3(n-1)}{2}} + x_2^{\frac{3(n-1)}{2}} + \cdots + x_n^{\frac{3(n-1)}{2}} + (n^2 - 1)(x_1^{\frac{n-1}{2}} x_2x_3\cdots x_n \right. \\
&\quad \left. + x_1x_2^{\frac{n-1}{2}} x_2\cdots x_n + x_1x_2\cdots x_{n-1}x_n^{\frac{n-1}{2}})\right)
\end{aligned}$$

where since $f(x)$ is decreasing and $f((\sum_{i=1}^n x_i^{\frac{n-1}{2}})^3) = f(1) = 1$, proving that the last line is greater than or equal to 1 is equivalent to showing the following inequality

$$\begin{aligned}
\left(\sum_{i=1}^n x_i^{\frac{n-1}{2}}\right)^3 &\geq x_1^{\frac{3(n-1)}{2}} + x_2^{\frac{3(n-1)}{2}} + \cdots + x_n^{\frac{3(n-1)}{2}} + (n^2 - 1)(x_1^{\frac{n-1}{2}} x_2x_3\cdots x_n \\
&\quad + x_1x_2^{\frac{n-1}{2}} x_2\cdots x_n + x_1x_2\cdots x_{n-1}x_n^{\frac{n-1}{2}}).
\end{aligned}$$

After expanding the left hand side and subtracting the terms containing $x_i^{\frac{3(n-1)}{2}}$ from both sides, there are $n^3 - n = n(n^2 - 1)$ terms remaining on the LHS, which can be divided into

n groups with each group containing $n^2 - 1$ terms, as follows.

$$\begin{aligned} & \sum_{i=1}^n \left(x_i^{n-1} \sum_{j \neq i}^n x_j^{\frac{n-1}{2}} + x_i^{\frac{n-1}{2}} \sum_{j \neq i}^n x_j^{n-1} + \sum_{j=1}^{n-1} x_{G_{i,j}}^{n-1} x_{G_{i,j+1}}^{\frac{n-1}{2}} + x_i^{\frac{n-1}{2}} \sum_{j \neq i}^n x_j^{\frac{n-1}{2}} \sum_{k \neq i,j}^n x_k^{\frac{n-1}{2}} \right) \\ & \geq (n^2 - 1) \sum_{i=1}^n x_1 x_2 \cdots x_{i-1} x_i^{\frac{n-1}{2}} x_{i+1} \cdots x_n \end{aligned} \quad (1)$$

where $G_i = (1, 2, \dots, i-1, i+1, \dots, n)$ denotes a cyclic ordered sequence, in which $G_{i,j} = j$ for $0 < j < i$, $G_{i,j} = j+1$ for $i \leq j < n$, otherwise $G_{i,n} = G_{i,1}$. By applying GM inequality to each summand for the LHS of (1) and adding them together, we can get the right-hand side (RHS) of (1). For better presentation, we show the derivation for the first group and the other $n-1$ groups follow the same logic.

$$\begin{aligned} & x_1^{n-1} x_2^{\frac{n-1}{2}} + x_1^{\frac{n-1}{2}} x_2^{n-1} + x_2^{n-1} x_3^{\frac{n-1}{2}} + x_1^{n-1} x_3^{\frac{n-1}{2}} + x_1^{\frac{n-1}{2}} x_3^{n-1} \\ & + x_3^{n-1} x_4^{\frac{n-1}{2}} + \cdots + x_1^{n-1} x_n^{\frac{n-1}{2}} + x_1^{\frac{n-1}{2}} x_n^{n-1} + x_n^{n-1} x_2^{\frac{n-1}{2}} \\ & + x_1^{\frac{n-1}{2}} x_2^{\frac{n-1}{2}} x_3^{\frac{n-1}{2}} + x_1^{\frac{n-1}{2}} x_3^{\frac{n-1}{2}} x_2^{\frac{n-1}{2}} + x_1^{\frac{n-1}{2}} x_2^{\frac{n-1}{2}} x_4^{\frac{n-1}{2}} \\ & + x_1^{\frac{n-1}{2}} x_4^{\frac{n-1}{2}} x_2^{\frac{n-1}{2}} + \cdots + x_1^{\frac{n-1}{2}} x_{n-1}^{\frac{n-1}{2}} x_n^{\frac{n-1}{2}} + x_1^{\frac{n-1}{2}} x_n^{\frac{n-1}{2}} x_{n-1}^{\frac{n-1}{2}} \\ & \geq (n^2 - 1) \sqrt[n^2-1]{x_1^{\frac{3(n-1)^2}{2} + (n-1)(n-2) \cdot \frac{n-1}{2}} x_2^{3(n-1) + 2(n-2) \cdot \frac{n-1}{2}} \cdots x_n^{3(n-1) + 2(n-2) \cdot \frac{n-1}{2}}} \\ & = (n^2 - 1) \sqrt[n^2-1]{x_1^{\frac{(n-1)^2(n+1)}{2}} x_2^{n^2-1} \cdots x_n^{n^2-1}} \\ & = (n^2 - 1) x_1^{\frac{(n-1)^2(n+1)}{2(n^2-1)}} x_2 \cdots x_n \\ & \geq (n^2 - 1) x_1^{\frac{n-1}{2}} x_2 x_3 \cdots x_n \end{aligned}$$

which completes the proof.