

TOWARDS NIELSEN-THURSTON CLASSIFICATION FOR SURFACES OF INFINITE TYPE: WELL-TEMPERED HOMEOMORPHISMS

MLADEN BESTVINA, FEDERICA FANONI, AND JING TAO

ABSTRACT. We introduce and study tempered mapping classes of surfaces of infinite type. These are maps for which curves under iteration do not accumulate onto geodesic laminations with non-proper leaves, but only on unions of possibly intersecting curves or proper lines. Assuming an additional finiteness condition on the accumulation set, we prove a Nielsen-Thurston-type classification theorem. We prove that for such maps there is a canonical decomposition of the surface into invariant subsurfaces on which the first return is either periodic or a translation.

1. INTRODUCTION

In this paper, we study mapping classes – that is, isotopy classes of self-homeomorphisms – of a connected, orientable surface S of *infinite* type. For finite-type surfaces, the Nielsen–Thurston classification gives a complete picture. Every mapping class is periodic, reducible, or pseudo-Anosov. Periodic classes are represented by isometries of suitable hyperbolic metrics via Nielsen realization; Pseudo-Anosov classes preserve a pair of transverse measured laminations, expanding one and contracting the other; reducible classes decompose the surface along a multicurve, reducing the dynamics to subsurfaces on which the first-return maps are again periodic or pseudo-Anosov.

The long-term goal of this project is to extend this understanding to surfaces of infinite type. Here, we take a step in that direction: we introduce the notion of *tempered* mapping classes and prove a structure theorem for the subclass of *well-tempered* mapping classes.

To motivate our definitions, we briefly recall several approaches to the Nielsen-Thurston theorem and assess their prospects in the infinite setting. Thurston’s original proof [28, 1] is to find a fixed point in the compactified Teichmüller space. Bers’ proof [10] also uses Teichmüller space, but from the point of view of extremal quasiconformal maps. Casson’s proof [14] produces an invariant geodesic lamination by iterating a curve, while Nielsen’s approach [23, 24, 25], completed by Miller [22] and Handel-Thurston [16], analyzes the dynamics on the circle at infinity of lifts of the map to the universal cover. The Bestvina-Handel proof [11]

produces invariant train tracks from efficient spines of the surfaces and relies on Perron-Frobenius theory.

In the infinite-type setting, however, Teichmüller space is much more complicated: it is infinite-dimensional, has uncountably many connected components, and some mapping classes do not preserve any component. So it is hard to imagine how Thurston's or Bers' approach could be adapted. Likewise, train-track methods would require infinite-sized matrices without a suitable analog of Perron-Frobenius theory. By contrast, the viewpoints of Casson and Nielsen-Handel-Thurston seem more amenable. In this paper, we take Casson's approach.

Let us say more carefully what Casson's method requires, because the difficulties in infinite type are already visible here. Given $f : S \rightarrow S$ and a curve α , one takes a subsequence $f^{n_i}(\alpha)$, pulls each iterate tight in some hyperbolic metric, and attempts to extract a geodesic lamination λ as a limit. For λ to be useful—specifically for $\bigcup_n f^n(\lambda)$ to be an f -invariant lamination without transverse intersections, the convergence must be “robust”, meaning that more and more strands of $f^{n_i}(\alpha)$ should converge to the leaves of λ . On finite-type surfaces, this robustness is essentially automatic; it fails only when α intersects a subsurface on which f is periodic, up to isotopy. On an infinite-type surface, it can fail in ways that are genuinely new and geometrically interesting.

Three examples are worth keeping in mind. The first example comes from infinite-order isometries, such as the translation $(x, y) \mapsto (x + 1, y)$ on $\mathbb{R}^2 \setminus \mathbb{Z}^2$, which sends every curve to infinity. This is an instance of a general *translation*—a homeomorphism that generates an infinite cyclic group acting properly on the surface. For a more complicated example, consider the same surface $\mathbb{R}^2 \setminus \mathbb{Z}^2$ but consider a map that agrees with the translation on $y \leq 1/3$, the inverse of the translation on $y \geq 2/3$, and a continuous interpolation of the two by horizontal translations in the strip $1/3 < y < 2/3$. In this case, a curve that intersects the strip essentially will converge to a line homotopic into the strip, but the convergence is not robust. A different phenomenon is an *irrational rotation*. Start with a homeomorphism of the equator in S^2 whose minimal invariant set is a Cantor set C , and extend it to a homeomorphism f of $S^2 \setminus C$. Here, a convergent subsequence $f^{n_i}(\alpha)$ limits onto a line in a non-robust way, and the union of all such limits is a non-locally finite collection of lines, some of which intersect; in particular, it is not a geodesic lamination. These are not exotic counterexamples; they represent genuinely different dynamical behaviors, and any classification theorem must account for all of them.

With these examples in mind, we arrive at the central definition of *tempered* maps, which are precisely those mapping classes where robust convergence never occurs for any curve; in particular, a tempered mapping class does not exhibit pseudo-Anosov behavior anywhere on the surface. Concretely, $f : S \rightarrow S$ is tempered if for any two curves α, β on S , there is a uniform bound on the intersection numbers $i(f^n(\alpha), \beta)$ for all $n \in \mathbb{Z}$ (see Section 4). As shown in Section 4, if f is tempered, then a curve α either leaves every compact set, or has limit set $\mathcal{L}(\alpha)$

consisting of curves or properly embedded lines (rather than more complicated laminations). A tempered map is *well tempered* if, in addition, the limit set of every curve is a *finite* collection of lines and curves. These notions are well-defined on the mapping class of f .

On a finite-type surface, tempered is the same as periodic. On an infinite-type surface, the class is considerably richer. The basic examples are periodic maps, translations, and irrational rotations, with the first two being well tempered, and irrational rotations tempered but not well-tempered. More complicated examples can be built by gluing such maps together in a controlled way—for instance, by gluing two translations along a fixed line, as in the second example above—and this construction can be extended to infinitely many maps.

Our main theorem shows that well-tempered mapping classes are exactly those arising from this kind of construction. In particular, they admit a satisfying Nielsen–Thurston-type theory.

Theorem A. *Let ϕ be a well-tempered mapping class of a surface S of infinite type. Then there exists a representative $f \in \phi$ and an f -invariant 1-manifold $\lambda \subset S$ without compact components such that, for the closure X of each component of $S \setminus \lambda$, f returns to X , and its first return is isotopic either to a periodic map or a translation.*

It is natural to ask whether Theorem A admits an analogue for arbitrary tempered mapping classes. The full strength of the theorem can fail in several ways. First, one must allow for irrational-rotation behavior. Second, the fact that well-tempered maps are reduced along a 1-manifold is special to that setting; it ensures that the natural invariant pieces are always subsurfaces with boundary. For general tempered maps, by contrast, these pieces may have boundaries that accumulate in a complicated way, and hence need not be subsurfaces (see the *Strategy of proof* section for more details). Even assuming local finiteness of limit sets of curves is not enough to circumvent this issue (see Section 7.1). Third, the existence of first-return maps is itself a nontrivial feature of the well-tempered setting and need not to exist in general.

Despite these complications, one might still hope that every tempered mapping class can be decomposed into pieces modeled on periodic maps, translations, and irrational rotation. Indeed, in the first version of this article, we conjectured such a picture, but it turns out to be incomplete. In addition to the three types above, there is a fourth type of dynamics, which we call an *almost translation*. This is a map f such that, for every two curves α and β , the set $\{n \mid i(f^n(\alpha), \beta) \neq 0\}$ has zero density in $\mathbb{Z}$.

Taking these complications into account, we show in forthcoming work that for every tempered mapping class ϕ , there exists a representative $f \in \phi$ and an f -invariant lamination λ such that, for every complementary component of $S \setminus \lambda$, whenever a first return map exists, it is isotopic either to a periodic map, an almost translation, or an irrational rotation. We also show there that almost

translations do occur. Whether the leaves of λ can always be taken to be proper remains open.

Beyond the tempered case, if ϕ is not tempered, one could still try to construct an invariant geodesic lamination using the Casson technique. This approach was carried out successfully for the class of *irreducible end-periodic* maps by Handel and Miller [13]. In our vision, a comprehensive structure theorem for an arbitrary mapping class ϕ should begin by decomposing S into invariant pieces on which the first return is either tempered or preserves two transverse geodesic laminations. Therefore, a structure theorem for tempered maps is an essential part of the broader program, and Theorem A should be regarded as a first step.

Strategy of proof. Given a map $f : S \rightarrow S$, our strategy for decomposing S into f -invariant pieces is to group curves according to their dynamical behavior under f , and then analyze the subsurface *spanned* by curves of a given type, when that subsurface exists. For instance, let $\mathcal{C}_{\text{per}}$ be the collection of f -periodic curves. If the subsurface S_{per} spanned by $\mathcal{C}_{\text{per}}$ exists, then it is f -invariant (up to isotopy) and the action of f on each component of S_{per} is isotopic to a periodic map (see Section 9).

We apply the same idea to the collection $\mathcal{C}_{\infty}$ of *wandering* curves, namely curves whose iterates leave every compact set of S . With some work, one can show that if the subsurface S_{∞} spanned by $\mathcal{C}_{\infty}$ exists, then the action of f on each component of S_{∞} is isotopic to a translation. This may be viewed as an analogue of Brouwer's plane translation theorem ([12]) and it is proved in Section 8.

Theorem B. *Suppose ϕ is a mapping class of a surface S such that every curve of S is wandering. Then there is a representative $f \in \phi$ and a hyperbolic metric on S such that f is an isometric translation.*

Thus, once the existence of S_{∞} has been established, then Theorem B reduces the proof of the main theorem to showing that the first return map on each component of $S \setminus S_{\infty}$ exists and is periodic.

This strategy makes the existence of spanning subsurfaces a central technical issue. For a finite collection of curves, the span is always well defined: put the curves in general position, take the union of their regular neighborhoods, and cap off any components homeomorphic to a disk with at most one puncture. For an infinite collection, one may take the span S_n of the first n curves and study the limiting object as $n \rightarrow \infty$. On a finite-type surface, Euler characteristic considerations force the topology of S_n to stabilize, and this procedure essentially recovers the Nielsen-Thurston decomposition of f . On an infinite-type surface, however, the limiting object need not be a subsurface of S , but may instead be a complementary component of a geodesic lamination. Indeed, there are examples of tempered maps for which the collection of periodic or wandering curves fail to span a subsurface (see Section 7.1). Under the additional assumption that f is

well tempered, however, this bad behavior disappears, allowing the argument to go through.

Proposition C. *If f is well tempered, then the collection of f -periodic curves spans a subsurface S_{per} , and the collection of f -wandering curves spans a subsurface S_{∞} .*

Proposition C is proved in Section 7 and follows from Lemmas 7.2 and 7.4.

Note that periodic curves and wandering curves together do not, in general, span the whole surface – said differently, $S_{\infty} \cup S_{\text{per}}$ is not always S . We therefore need to study the complement of this union, denoted S_0 , and show that it contains no essential nonperipheral curves, and that the first-return map on each of its components exists and has finite order. In fact, Section 9 shows a more technical statement than Theorem A – namely, Theorem 9.1 – from which our main theorem is deduced.

Connection to actions on graphs. A related, and perhaps more tractable, variant of the Nielsen-Thurston classification theorem is to study the action of mapping classes on Gromov-hyperbolic graphs naturally associated to a surface, and to classify mapping classes by their dynamic types as graph isometries. For example, in the finite-type setting, pseudo-Anosov mapping classes act loxodromically on the curve graph of S , while all other mapping classes act elliptically.

In the infinite-type setting, the curve graph always has bounded diameter, reducing the classification of isometries to a trivial problem. However, many infinite-type surfaces admit other interesting Gromov-hyperbolic graphs of infinite diameter. One example is the ray graph, introduced by Bavard [8] for the plane minus a Cantor set $\mathbb{R}^2 \setminus C$, generalized to surfaces with one isolated end by Aramayona-Fossas-Parlier [5], and extended further by Bar-Natan-Verberne [7], who define the *grand-arc graph*. In [8], Bavard also constructed a map of $\mathbb{R}^2 \setminus C$ that acts loxodromically on the ray graph, and Abbott-Miller-Patel [2] generalized this construction to a large class of surfaces. However, a general classification of isometries remains open, and even the following question is unknown.

Question 1.1. Are there mapping classes of $\mathbb{R}^2 \setminus C$ that act as parabolic isometries of the ray graph? More generally, are there mapping classes that act as parabolic isometries of the grand arc graphs?

By our work above, well-tempered mapping classes are elliptic isometries of the ray graph, and our future work on tempered mapping classes suggests that they are also elliptic isometries.

Acknowledgments. We thank the American Institute of Mathematics for the hospitality during the workshop “Surfaces of infinite type”, where this project began. We are also grateful to Peter Feller, Diana Hubbard, and Hannah Turner for sharing their preprint [15], which was useful for refining our work on tempered maps. We thank Sanghoon Kwak for many useful comments on the first version

of this paper, and the referee for their careful reading and comments. M.B. gratefully acknowledges support from the National Science Foundation under grant DMS-1905720. J.T. gratefully acknowledges support from the National Science Foundation under grants DMS-1651963 and DMS-2304920. F.F. also thanks Peter Feller for useful conversations.

2. BACKGROUND

Standing assumptions. By a surface S we will mean a connected orientable 2-manifold, without boundary unless otherwise stated. When we need to consider surfaces with boundary, we will usually say *bordered* surface for emphasis. By a *hyperbolic metric* on S we always mean a complete hyperbolic metric without funnels or half-planes; equivalently, the hyperbolic surface coincides with its convex core, which is also equivalent to the metric being of the *first kind* - see [4]. For bordered surfaces, hyperbolic metrics are assumed to contain no funnels or half-planes and to have totally geodesic boundary.

A surface S has *finite type* if $\pi_1(S)$ is finitely generated and *infinite type* otherwise. By the classification of surfaces, S is of finite type if and only if S is homeomorphic to a closed genus g surface with finitely many points removed. A *pair of pants* is a (possibly bordered) surface homeomorphic to either a three-holed sphere, or a two-holed once-punctured sphere, or a twice-punctured disk.

Associated to S are its *ends*, which can be *planar* or *nonplanar*. A *puncture* is an isolated planar end. We refer to [6] for definitions and properties of ends of a surface.

The *mapping class group* of S is the group $\text{Map}(S)$ of isotopy classes of orientation-preserving homeomorphisms of S .

Curves on S are assumed to be simple, closed, and not homotopic to a point or a puncture. We will sometimes call such a curve *essential* for emphasis. If S has boundary, then a curve is called *peripheral* if it is homotopic to a boundary component. When S is endowed with a hyperbolic metric with geodesic boundary, every essential curve is realized by a unique geodesic α^* representing its homotopy class.

A *line* on S is the image of a proper embedding of $\mathbb{R} \rightarrow S$. We will usually assume a line is *essential*, meaning it does not bound a (topological) half-plane. We will usually consider lines up to proper isotopy relative to their ends. Similarly to curves, essential lines admit geodesic representatives:

Lemma 2.1. *Let ℓ be an essential line on a surface S endowed with a hyperbolic metric (of the first kind). Then ℓ is isotopic to a unique geodesic ℓ^* .*

Sketch of proof. Fix a pants decomposition of S , which yields an exhaustion of S by finite-type subsurfaces with compact totally geodesic boundary (see [4]). Pick a lift $\tilde{\ell}$ of ℓ in the universal cover. Since ℓ is proper and the metric is of the first kind, by looking at the lifts of the pants curves we can conclude that $\tilde{\ell}$ has two distinct limit points on the circle at infinity. Let $\tilde{\ell}^*$ be the unique geodesic joining

the two limit points and let ℓ^* be the projection of $\tilde{\ell}^*$ to S . Arguing as in the proof of Proposition 3.3, one shows that ℓ is isotopic to ℓ^* . Uniqueness follows because the limit points of $\tilde{\ell}$ determine $\tilde{\ell}^*$ and hence ℓ^* . $\square$

An *arc* on S is the image of an embedding $[0, 1] \rightarrow S$; in particular in our definition arcs are compact. If S has boundary, an arc with endpoints on ∂S is called *essential* if it doubles to an essential curve in the double of S .

A *ray* is the image of $[0, \infty)$ under a proper embedding.

A *geodesic lamination* is a closed subset of S that is a union of complete, simple, and pairwise disjoint geodesics (with respect to some hyperbolic metric on S), called the *leaves* of the lamination.

The *geometric intersection number* $i(\alpha, \beta)$ is the minimum number of intersections between representatives of α and β in their homotopy classes, where each of α and β could be either a line, curve, or a leaf of a lamination. Note that $i(\alpha, \beta)$ could be infinite, but we have the following characterization of lines.

Lemma 2.2. *Let S be a surface equipped with a hyperbolic metric. Then a complete simple geodesic $\ell \subset S$ is a line or a curve if and only if $i(\ell, \alpha) < \infty$ for all curves α .*

Proof. Clearly, if $i(\ell, \alpha) = \infty$ for some curve α , then ℓ cannot be proper. Conversely, suppose $i(\ell, \alpha) < \infty$ for all curves α . Let $K \subset S$ be a compact subsurface. Since $\sum_{\alpha \subset \partial K} i(\ell, \alpha) < \infty$, $\ell \cap K$ has a finite number of components. If a component $\tau \subset \ell \cap K$ had infinite length, then τ would accumulate onto some geodesic lamination $\lambda \subset K$, or a boundary component γ of K . In either case, any curve α intersecting λ (or γ) would satisfy $i(\ell, \alpha) \geq i(\tau, \alpha) = \infty$. Thus $\ell \cap K$ is always a finite union of arcs, and hence ℓ is proper. $\square$

2.1. Subsurfaces. A *subsurface* of S is a closed subset $X \subset S$ that is a two-dimensional (possibly disconnected) submanifold such that each boundary component of X is either an essential curve or an essential line of S . We also require that no connected component of X can be homotoped into another. In particular, no component of a subsurface can be a closed disk with at most one puncture. Moreover, X is a subsurface if and only if $Y = \overline{S \setminus X}$ is also a subsurface. We will usually consider a subsurface up to isotopy.

A connected subsurface X is *essential* if its double either has negative Euler characteristic or is of infinite type. Topologically, this rules out annuli and *strips* (i.e. closed disks with two points removed from the boundary).

Two subsurfaces in S are *disjoint* if they have disjoint representatives. Similarly, a curve or line or geodesic lamination is disjoint from a subsurface if they have disjoint representatives. Note that in this convention, the boundary of a subsurface is considered disjoint from the subsurface.

Fix a hyperbolic metric on S . Because we will usually consider a subsurface up to isotopy, it is convenient to pick out a canonical representative within each isotopy class. To this end, we will introduce the notion of an (*almost*) *geodesic*

representative of a subsurface $X \subset S$, subject to the hyperbolic metric on S , which serve as an *almost* canonical representative of the isotopy class of X .

We begin with representatives of the interiors of subsurfaces. First suppose that X is connected. A *geodesic representative of the interior of X* is an open set X° defined as follows:

- If X is essential, then X° is the open set properly homotopic to $\text{int}(X)$ whose metric completion is a hyperbolic surface with totally geodesic boundary, homeomorphic to X .
- If X is an annulus or a strip with core curve/line α , then X° is the interior of any closed regular neighborhood of α^* , whose boundary components are proper curves or lines.

For a (not necessarily connected) subsurface X , a *geodesic representative of its interior* (also denoted X°) is the union, over the connected components of X , of geodesic representatives of their interiors. By construction, X° is an open subset properly homotopic to $\text{int}(X)$.

To get a representative for X , we proceed as follows. For each component Y of X , let $\overline{Y^\circ}$ be the closure of Y° , and set $\partial Y^\circ := \overline{Y^\circ} \setminus Y^\circ$. When Y is essential, then ∂Y° is a disjoint union of simple closed geodesics and geodesic lines; otherwise, ∂Y° consists of a pair of almost geodesic curves or lines, each homotopic to the core curve/line of Y . Set $\partial X^\circ := \bigcup_{Y \subset X} \partial Y^\circ$, and let $\partial_{sa} X^\circ$ consist of those components α of ∂X° such that both boundary components of a regular neighborhood of α are contained in X° (informally, boundary components which will be “self-adjacent”). We define an *almost geodesic representative* of X by

$$X^* = \overline{X^\circ} \setminus \bigcup_{\alpha \in \partial_{sa} X^\circ} N(\alpha)$$

where $N(\alpha)$ is an open regular neighborhood α .

An almost geodesic representative is well-defined up to our choices of regular neighborhoods of the components in $\partial_{sa}(X^\circ)$ and the representatives of the inessential components of X (and the hyperbolic metric on S). In particular, if $\partial_{sa}(X^\circ) = \emptyset$ and X has no inessential components, then X^* is a totally geodesic subsurface representing X . Moreover, $\overline{X^\circ}$ is homeomorphic to X^* if and only if $\partial_{sa} X^\circ = \emptyset$.




FIGURE 1. An example of a subsurface X so that $\overline{X^\circ}$ is not homeomorphic to X^* . The curves α and β are the geodesic representatives of the boundary components of X .

Lemma 2.3. *For any hyperbolic metric on S , any subsurface is properly homotopic to its almost geodesic representative.*

Sketch of proof. If $\partial_{sa}X^\circ = \emptyset$ and X has no inessential components, then the almost geodesic representative agrees with the geodesic representative of X , and the claim follows by properly homotoping all boundary components of X at once to their geodesic representatives (see Lemma 2.1 and [14, Lemma 2.6]). If $\partial_{sa}X^\circ \neq \emptyset$ or X has annular or strip components, then the same argument applies with a minor modification to handle pairs of parallel boundary components. $\square$

A consequence is that “inclusion up to proper homotopy” defines a partial order on the set of equivalence classes of subsurfaces defined up to proper homotopy:

Corollary 2.4. *Let S be a surface equipped with a hyperbolic metric. If X_1 and X_2 are subsurfaces such that each is properly homotopic into the other, then X_1 and X_2 are properly homotopic.*

Proof. We first prove that inessential components of X_1 and X_2 are the same, up to proper homotopy. Indeed, suppose Y is a inessential component of X_1 . Since X_1 is properly homotopic into X_2 , Y properly homotopes into some component Z of X_2 . Since X_2 properly homotopes into X_1 , Z is properly homotopic into some component Y' of X_1 . Thus Y properly homotopes into Y' , and by the definition of subsurface (no component homotopes into another), we must have $Y = Y'$. In particular, Z properly homotopes into a strip or an annulus and is hence a strip or an annulus itself. By repeating the same argument for the inessential components of X_2 we get the claim.

We may therefore assume that X_1 and X_2 have no inessential components. With respect to the hyperbolic metric on S , X_1° and X_2° are uniquely defined. We claim they are the same. Otherwise, after swapping indices if necessary, there exists a component Y of $X_1^\circ \setminus X_2^\circ$. Since X_1 properly homotopes into X_2 , such Y can only be a curve or a line, so $Y \subset \partial_{sa}X_1^\circ$. Choose an essential curve or line $\alpha \subset X_1^\circ$ that intersects Y essentially. Again, because X_1 properly homotopes into X_2 , α can be homotoped into X_2 and hence must be disjoint from Y , a contradiction. This shows $X_1^\circ = X_2^\circ$, so X_1 and X_2 have the same almost geodesic representative. Lemma 2.3 then implies that X_1 and X_2 are properly homotopic. $\square$

2.2. Some results on isometries of surfaces. We say that a collection $\mathcal{C}$ of curves *fills* a (sub)surface X if every nonperipheral curve in X has positive intersection number with some curve in $\mathcal{C}$. The following well-known statement follows from the proof of [14, Theorem 2.7].

Lemma 2.5. *Let F be a possibly bordered finite-type surface of negative Euler characteristic, whose boundary is either empty or compact. Let f be a mapping class of F . If F is filled by a finite collection of f -periodic curves, then f has finite order.*

Remark 2.6. The order of f in Lemma 2.5 can be arbitrarily larger than the periods of the filling curves. For instance, consider the torus $\mathbb{R}^2/\mathbb{Z}^2$ and the curves α and β given by the projection of the lines $y = nx$ and $y = -nx$, for some positive integer n . Puncture the torus at the points that are the projections of $(\frac{1+2k}{2n}, 0)$ and $(\frac{k}{n}, \frac{1}{2})$, for $k = 0, \dots, n-1$, which do not meet α and β . Then the map induced by $(x, y) \mapsto (x + \frac{1}{2n}, y + \frac{1}{2})$ has order $2n$, but it fixes both α and β .

We will also need a realization result for periodic mapping classes of infinite-type surfaces; the proof is essentially borrowed from Afton-Calegari-Chen-Lyman ([3]).

Proposition 2.7. *Let S be an infinite-type surface with boundary, and f a homeomorphism of S such that f^n homotopic to the identity for some $n > 0$. For each compact boundary component α of S , choose a positive number $\ell(\alpha)$ that is f -invariant, i.e. $\ell(f(\alpha)) = \ell(\alpha)$. Then there exists a hyperbolic metric on S such that each compact boundary component α has length $\ell(\alpha)$ and f is isotopic to a periodic isometry of S .*

Proof. Suppose first that S has only compact boundary components. Using Nielsen's realization theorem for finite-type surfaces with compact boundary [26] and the argument of Afton-Calegari-Chen-Lyman (see the proof of [3, Theorem 2]), one obtains the desired metric and periodic representative in this case.

Now suppose S has noncompact boundary components. Let DS be the double of S along its noncompact boundary and extend f to a homeomorphism $\hat{f}$ of DS . Let i be the involution of DS with quotient S , which commutes with $\hat{f}$ by construction. By [3] and the discussion above, there is an i -invariant metric on DS realizing the prescribed lengths for the compact components, and a periodic isometry g isotopic to $\hat{f}$ that commutes with i . Since i is an isometry, the noncompact boundaries of S are realized as geodesics in DS , so that the metric descends to a metric on S with totally geodesic boundary. Since g commutes with i , it descends to a periodic isometry of S isotopic to f . $\square$

3. LIMIT SETS

Throughout this section, S is a surface without boundary equipped with a hyperbolic metric, though everything we say will be independent of the particular choice of metric. Indeed, the following holds (see [29, Theorem 3.6] for the infinite-type case). Recall that all hyperbolic metrics are of the first kind.

Theorem 3.1. *Let m and m' be two hyperbolic metrics on S . The identity map on S induces a natural identification between the boundaries at infinity of the universal covers of (S, m) and (S, m') . This identification yields a homeomorphism between the spaces of complete geodesics in the universal covers of (S, m) and (S, m') , which descends to a homeomorphism between complete m -geodesics and complete m' -geodesics on S .*

Given a complete m -geodesic ℓ , we call the corresponding m' -geodesic the m' -straightening of ℓ . For a collection of m -geodesics, its m' -straightening is the union of the m' -straightenings of its geodesics. A consequence of Theorem 3.1 is the following:

Lemma 3.2. *Let m and m' be two hyperbolic metrics on S . If two complete m -geodesics ℓ_1 and ℓ_2 are simple and disjoint, then the m' -straightenings ℓ'_1 and ℓ'_2 are also simple and disjoint. If λ is an m -geodesic lamination, its m' -straightening is an m' -geodesic lamination.*

The reason why the lemma holds is that (self-)intersections correspond to linked pairs of endpoints at infinity of lifts and convergence of geodesics corresponds to convergence of pairs of endpoints at infinity of lifts. Moreover, we have:

Proposition 3.3. *Let m and m' be hyperbolic metrics on S and λ an m -geodesic lamination. Then there is a homeomorphism of S , isotopic to the identity, that maps λ leaf-by-leaf to its m' -straightening λ' .*


Proof. Pick an m -geodesic pants decomposition $\mathcal{P}$ that contains no leaf of λ (this can be done by choosing a pants decomposition and performing elementary moves until none of its curves lie in λ). In particular, $\mathcal{P}$ is in minimal position with respect to λ . Let $\mathcal{P}'$ be the m' -straightening of $\mathcal{P}$. There is an ambient isotopy H_1 of S sending $\mathcal{P}$ to $\mathcal{P}'$. Let $\lambda_1 := H_1(\lambda)$. Then $\mathcal{P}'$ is in minimal position with respect to both λ_1 and λ' , and contains no leaf of either.

Now pick a curve α in $\mathcal{P}'$, and lift α , λ_1 , and λ' to the universal cover of (S, m') . Choose and orient a lift β of α . Given a lift ℓ_1 of a leaf in λ_1 , its m' -straightening is the unique lift ℓ'_1 of the m' -straightened leaf in λ' with the same endpoints; in particular, ℓ_1 intersects β if and only if ℓ'_1 intersects β . Moreover, if ℓ_1 and ℓ_2 are two such lifts intersecting β , and ℓ'_1 and ℓ'_2 are their m' -straightenings, then the intersection points $\ell_1 \cap \beta$ and $\ell_2 \cap \beta$ occur in the same order along β as $\ell'_1 \cap \beta$ and $\ell'_2 \cap \beta$. Hence there is an order-preserving homeomorphism

$$\tilde{\lambda}_1 \cap \beta \rightarrow \tilde{\lambda}' \cap \beta$$

sending $\ell \cap \beta$ to $\ell' \cap \beta$, for every leaf $\ell \subset \tilde{\lambda}_1$, where ℓ' is its m' -straightening. Extending this to an equivariant isotopy on a small neighborhood of β , and doing so equivariantly for every lift of every curve α in $\mathcal{P}'$ on pairwise disjoint neighborhoods, produces an equivariant isotopy of the universal cover. This descends to an ambient isotopy H_2 of S , sending λ_1 to a lamination λ_2 , such that $\lambda_2 \cap \alpha = \lambda' \cap \alpha$ for all α in $\mathcal{P}'$.

Now let P be a pair of pants of $\mathcal{P}'$. Since λ_2 and λ' are transverse to ∂P , $\lambda_2 \cap P$ and $\lambda' \cap P$ are unions of arcs with endpoints on ∂P . We claim that there is an isotopy of P , fixing ∂P pointwise, sending $\lambda_2 \cap P$ to $\lambda' \cap P$. Partition the arcs of $\lambda_2 \cap P$ into (at most three) homotopy classes relative to the boundary. Do the same for the arcs of $\lambda' \cap P$; note that the same classes occur for both. For a class of $\lambda_2 \cap P$, choose a rectangle $R \subset P$, whose two opposite sides lie in ∂P and whose two other sides are distinct arcs of $\lambda_2 \cap P$, such that R contains all arcs of $\lambda_2 \cap P$

FIGURE 2. Lifting a curve in $\mathcal{P}'$

in that class. Construct a corresponding rectangle $R' \subset P$ for the same class of $\lambda' \cap P$, arranged so that R and R' have the same four corners on ∂P . Then an isotopy of P fixing ∂P pointwise can be chosen to send R to R' and the arcs of $\lambda_2 \cap R$ to the corresponding arcs of $\lambda' \cap R'$. Doing this for each homotopy class of arcs yields the desired isotopy of P .

Repeating the procedure in each pair of pants produces an ambient isotopy H_3 of S , fixing each curve of $\mathcal{P}'$ pointwise, and sending λ_2 to λ' . The composition $H_3 \circ H_2 \circ H_1$ is the required isotopy. $\square$




FIGURE 3. Isotoping rectangles

Let $\lambda \subset S$ be a closed subset. We say that a sequence of curves $\{\alpha_n \mid n \in \mathbb{N}\}$ *converges* to λ , and that λ is the *limit* of $\{\alpha_n \mid n \in \mathbb{N}\}$, if for every finite-type subsurface $K \subset S$ with compact boundary, $\alpha_n^* \cap K$ converges to $\lambda \cap K$ with respect to the Hausdorff distance. We will also denote this by $\alpha_n \rightarrow \lambda$. The same definition applies to sequences of geodesic arcs. Since convergence is tested on finite-type subsurfaces, the standard argument (see e.g. [14]) implies:

Lemma 3.4. *Let $\lambda \subset S$ be a closed subset. If λ is the limit of a sequence of curves $\{\alpha_n \mid n \in \mathbb{N}\}$, then λ is a geodesic lamination. If λ is the limit of a sequence of geodesic arcs, it is a union of pairwise disjoint geodesic arcs, geodesic rays and/or complete simple geodesics.*

The *limit set* of a sequence of curves $\{\alpha_n \mid n \in \mathbb{N}\}$ is the set $\mathcal{L}(\{\alpha_n \mid n \in \mathbb{N}\})$ consisting of all complete geodesics that occur as leaves of the limit of some subsequence of $\{\alpha_n \mid n \in \mathbb{N}\}$. Similarly we can define the limit set of a sequence of geodesic arcs. Changing the metric on S does not change the limit set, in the following sense.

Lemma 3.5. *Let m and m' be two hyperbolic metrics on S . For a sequence $\{\alpha_n\}$ of curves, let $\mathcal{L}(\{\alpha_n \mid n \in \mathbb{N}\}, m)$ and $\mathcal{L}(\{\alpha_n \mid n \in \mathbb{N}\}, m')$ be their respective limit sets in the two metrics. Then a subsequence $\{\alpha_{n_j}\}_{j \in \mathbb{N}}$ converges to $\lambda \subset \mathcal{L}(\{\alpha_n \mid n \in \mathbb{N}\}, m)$ if and only if $\{\alpha_{n_j}\}_{j \in \mathbb{N}}$ converges to $\lambda' \subset \mathcal{L}(\{\alpha_n \mid n \in \mathbb{N}\}, m')$, where λ' is the m' -straightening of λ .*

Proof. As convergence of geodesics corresponds to convergence of pairs of end-points at the boundary at infinity, this follows from Theorem 3.1. $\square$

This justifies our notation $\mathcal{L}(\{\alpha_n \mid n \in \mathbb{N}\})$, which makes no reference to the metric on S .

3.1. Limit sets under a homeomorphism. For a homeomorphism f of S and a curve α , we say α is *f-periodic* if there exists $n \geq 1$ such that $f^n(\alpha)$ is isotopic to α . The smallest such n is called the *f-period* of α . We say α is *f-forward wandering* if $\{f^n(\alpha)^*\}_{n \geq 0}$ leaves every compact set of S , and *f-backward wandering* if it is f^{-1} -forward wandering. The curve α is *f-wandering* if it is both *f-forward* and *f-backward* wandering. These properties are independent of the hyperbolic metric on S as well as perturbing f by an isotopy. Thus, we will also adopt the same definition for a mapping class of S . We will usually suppress the reference to f when the context is clear.

Note that a curve is *f-wandering* if and only if for every compact set $C \subset S$, $f^n(\alpha)$ can be homotoped to be disjoint from C for all sufficiently large $|n|$. With this in mind, we can extend the definition of *f-wandering* to lines or subsurfaces of S , by which we mean that for every compact set $C \subset S$, the images under f^n can be homotoped away from C for any n with sufficiently large $|n|$.

For a curve α , let $\mathcal{L}^+(\alpha) := \mathcal{L}(\{f^n(\alpha) \mid n \geq 0\})$ be the *forward limit* of α under f , and $\mathcal{L}^-(\alpha) := \mathcal{L}(\{f^n(\alpha) \mid n \leq 0\})$ the *backward limit*. Set $\mathcal{L}(\alpha) := \mathcal{L}^+(\alpha) \cup \mathcal{L}^-(\alpha)$. When α is *f-periodic*, then

$$\mathcal{L}^\pm(\alpha) = \{\alpha^*, f(\alpha)^*, \dots, f^{n-1}(\alpha)^*\},$$

where $n \geq 1$ is the *f-period* of α . Moreover, α is forward wandering if and only if $\mathcal{L}^+(\alpha) = \emptyset$, and it is backward wandering if and only if $\mathcal{L}^-(\alpha) = \emptyset$. See Corollary 4.3.

We establish some basic facts.

Lemma 3.6. *Let f be a homeomorphism and α a curve. Suppose λ is the limit of a sequence of iterates $\{f^{n_j}(\alpha) \mid j \in \mathbb{N}\}$. Then for any curve β , $i(\beta, \lambda) \neq 0$ if and only if $i(\beta, f^{n_j}(\alpha)) \neq 0$ for all sufficiently large j . Moreover, if $i(\beta, \lambda) \geq k$, then $i(\beta, f^{n_j}(\alpha)) \geq k$ for all sufficiently large j .*

Proof. If we look at an annular neighborhood K of β , $f^{n_j}(\alpha)^* \cap K$ converges to $\lambda \cap K$ in the Hausdorff metric, so the first statement follows. If β intersects λ at least k times, then we can find k arcs in $K \cap \lambda$. Choose sufficiently small neighborhood about each arc so that they are pairwise disjoint. Then for all sufficiently large j , $f^{n_j}(\alpha) \cap K$ will pass through each of these neighborhoods, so $i(\beta, f^{n_j}(\alpha)) \geq k$. $\square$

Lemma 3.7. *Let f be a homeomorphism. Then for any two (not necessarily distinct) curves α and β we have:*

$$i(\alpha, \mathcal{L}^+(\beta)) \neq 0 \iff i(\mathcal{L}^-(\alpha), \beta) \neq 0.$$

Proof. Suppose $i(\alpha, \mathcal{L}^+(\beta)) \neq 0$. Let $n_j \rightarrow \infty$ be a sequence so that $f^{n_j}(\beta) \rightarrow \lambda \subset \mathcal{L}^+(\beta)$ with $i(\alpha, \lambda) \neq 0$. By Lemma 3.6, for every j large enough,

$$i(\alpha, f^{n_j}(\beta)) \neq 0$$

and thus

$$i(f^{-n_j}(\alpha), \beta) \neq 0.$$

In particular, α is not backward wandering. By taking a further subsequence so that $f^{-n_{j_k}}(\alpha) \rightarrow \lambda' \subset \mathcal{L}^-(\alpha)$, we have, again by Lemma 3.6, $i(\lambda', \beta) \neq 0$, i.e. $i(\mathcal{L}^-(\alpha), \beta) \neq 0$. By replacing f by f^{-1} we get the other implication. $\square$

Lemma 3.8. *Let f be a homeomorphism. Then a curve α is forward wandering if and only if $i(\alpha, \mathcal{L}^-(\beta)) = 0$ for all curves β . Similarly, α is backward wandering if and only if $i(\alpha, \mathcal{L}^+(\beta)) = 0$ for all β .*

Proof. If α is not forward wandering, then we can choose a curve β with $i(\mathcal{L}^+(\alpha), \beta) \neq 0$. By Lemma 3.7, this implies $i(\alpha, \mathcal{L}^-(\beta)) \neq 0$. Conversely, if $i(\alpha, \mathcal{L}^-(\beta)) \neq 0$, then again by Lemma 3.7, $i(\mathcal{L}^+(\alpha), \beta) \neq 0$. In particular $\mathcal{L}^+(\alpha) \neq \emptyset$, so α is not forward wandering. $\square$

Corollary 3.9. *Let f be a homeomorphism. If α is a periodic curve and β is a wandering curve, $i(\alpha, \beta) = 0$.*

Proof. If $i(\alpha, \beta) \neq 0$, $i(\mathcal{L}^+(\alpha), \beta) \neq 0$ (as $\alpha \in \mathcal{L}^+(\alpha)$), so β is not wandering by Lemma 3.8. $\square$

4. TEMPERED MAPS

As usual, assume S is a surface without boundary endowed with a hyperbolic metric. We say a homeomorphism f of S is *tempered* if for every finite-type subsurface $K \subset S$ with compact boundary and every curve α there are $N = N(K, \alpha) \in \mathbb{N}$ and finitely many isotopy (relative to ∂K) classes of arcs from ∂K to ∂K and curves in K such that, for every $n \in \mathbb{N}$, the intersection $f^n(\alpha)^* \cap K$ has at most N components and each is in one of the given isotopy classes. Being tempered can be characterized by looking at intersections of curves, as the following lemma shows.

Lemma 4.1. *Let f be a homeomorphism of S . The following statements are equivalent.*

- (1) f is tempered.
- (2) f^{-1} is tempered.
- (3) For curves α and β , $i(f^n(\alpha), \beta)$ is uniformly bounded for all $n \geq 0$.
- (4) For curves α and β , $i(f^n(\alpha), \beta)$ is uniformly bounded for all $n \leq 0$.
- (5) For curves α and β , $i(f^n(\alpha), \beta)$ is uniformly bounded for all n .

In particular, being tempered is independent of the hyperbolic metric on S .

Proof. The equivalence of (3), (4), and (5) are immediate as the action of f preserves geometric intersection number. That (1) implies (3) follows from taking the compact surface to be an annular neighborhood of the curve β . On the other hand, if f is not tempered, then there exists a finite-type subsurface K with compact boundary, a curve α , and a subsequence $\{n_j\} \subset \mathbb{N}$ such that $f^{n_j}(\alpha)^* \cap K$ has N_j components with $N_j \rightarrow \infty$ and $j \rightarrow \infty$, or includes components with increasing multiplicity. Then there exists a curve $\beta \subset K$ such that $i(f^{n_j}(\alpha), \beta) \rightarrow \infty$. This shows the equivalence of (1) and (3), as well as (2) and (4) by symmetry, whence the equivalence of all statements. Property (5) is independent of the hyperbolic metric on S , so the same is true of being tempered. $\square$

We say a mapping class $\phi \in \text{Map}(S)$ is *tempered* if it has a tempered representative. From the characterization of Lemma 4.1, if ϕ has a tempered representative, then all of its representatives are tempered. We call a tempered homeomorphism f *well tempered* if the limit set $\mathcal{L}(\alpha)$ is finite for every curve α . This property is also inherited by the mapping class of f .

As a point of reference, on a finite-type surface the following are equivalent: being tempered, well tempered, periodic, and isotopic to an isometry for some hyperbolic structure. In the infinite-type case, the situation is more complicated. Being periodic implies being isotopic to an isometry (with respect to some hyperbolic structure), which implies being well tempered, and well-tempered maps are tempered by definition. None of these implications is an equivalence.

For instance, consider a translation of a surface S , which – as mentioned in the introduction – is a map f that generates an infinite cyclic group acting properly on S . If S has infinite type, then the quotient surface $X = S/\langle f \rangle$ has negative Euler characteristic or has infinite type, so we can lift a hyperbolic metric from X to S so that f acts by an (infinite-order) isometry. In this case, every curve on S is wandering, so f is (well) tempered.

Another example is what we call an *irrational rotation*. Start with a homeomorphism of the circle with minimal invariant subset a Cantor set C (see the Denjoy construction [19, Section 12.2]). View the circle as the equator of the two-sphere S^2 , and extend the map to a homeomorphism of $S = S^2 \setminus C$. This map is tempered but *not* well tempered: every curve has an infinite limit set, consisting of lines which may intersect. More complicated examples of (well) tempered maps can be found in Sections 7.1 and 9.

On the other hand, a Dehn twist is not tempered: if τ is the twist about a curve α , let β be a curve intersecting α essentially. If K is a finite-type subsurface with compact boundary containing α and β , the curves $\tau^n(\beta) \cap K = \tau^n(\beta)$ are all distinct. Another example of a map which is not tempered is a homeomorphism which restricts to a pseudo-Anosov on some finite-type subsurface (with compact boundary).

The goal of the remainder of this section is to describe properties of limit sets of curves and lines under tempered mapping classes. Our first result is that if a curve or line is not periodic, then its limit set can only contain lines.

Proposition 4.2. *Let f be a tempered homeomorphism and α a nonperiodic curve. If α is not forward wandering, then $\mathcal{L}^+(\alpha)$ is a non-empty collection of lines. Similarly, if α is not backward wandering, then $\mathcal{L}^-(\alpha)$ is a non-empty collection of lines.*

Proof. Assume α is not forward wandering, and for convenience, write $\alpha_n = f^n(\alpha)^*$. Suppose $\mathcal{L}^+(\alpha) = \emptyset$, then for every compact subsurface $K \subset S$, the set $\bigcup_{n \geq 0} \alpha_n \cap K$ has no accumulation points. Equivalently, $\bigcup_{n \geq 0} \alpha_n \cap K$ consists of finitely many arcs or curves. But two geodesics that coincide along a non-trivial arc must be equal, and α is not f -periodic, we must have $\alpha_n \cap K = \emptyset$ for all sufficiently large n . But this contradicts the assumption that α is not forward wandering. By Lemma 3.4, $\mathcal{L}^+(\alpha)$ is a non-empty union of complete simple geodesics. If $\mathcal{L}^+(\alpha)$ contains a non-proper leaf ℓ , then by Lemma 2.2 there exists a curve β such that $i(\beta, \ell) = \infty$. Let $\lambda \subset \mathcal{L}^+(\alpha)$ be the limit of some subsequence $\{\alpha_{n_j}\}$ containing ℓ . By Lemma 3.6 $i(f^{n_j}(\alpha), \beta) = i(\alpha_{n_j}, \beta) \rightarrow \infty$, contradicting the characterization of being tempered given by Lemma 4.1. If instead $\mathcal{L}^+(\alpha)$ contains a compact leaf β , then since $\mathcal{L}^+(\alpha)$ has no non-proper leaves, β arises as the limit of some subsequence $\{\alpha_{n_i}\}$. By the tempered condition, such a sequence has to be eventually constant, but this contradicts the nonperiodicity of α . Therefore $\mathcal{L}^+(\alpha)$ has no compact or non-proper leaves, so it is a non-empty union of lines. By replacing f by f^{-1} we get the second statement. $\square$

As $\mathcal{L}^\pm(\alpha)$ is well defined for mapping classes, we obtain the following immediate corollary.

Corollary 4.3. *For a tempered mapping class f and a curve or line α , the following statements hold.*

- (1) $\mathcal{L}^+(\alpha) = \emptyset$ if and only if α is forward wandering.
- (2) $\mathcal{L}^-(\alpha) = \emptyset$ if and only if α is backward wandering.
- (3) $\mathcal{L}(\alpha) = \emptyset$ if and only if α is wandering.
- (4) $\mathcal{L}(\alpha)$ contains a curve if and only if α is a periodic curve, in which case $\mathcal{L}(\alpha) = \mathcal{L}^+(\alpha) = \mathcal{L}^-(\alpha)$ is the orbit of α under f .

Another consequence of being tempered is that limits of iterates of curves can intersect only finitely many times.

Lemma 4.4. *Let f be a tempered homeomorphism. For any two (not necessarily distinct) curves α and β , if $f^{n_j}(\alpha) \rightarrow \lambda \subset \mathcal{L}(\beta)$ and $f^{m_k}(\beta) \rightarrow \lambda' \subset \mathcal{L}(\beta)$, for some sequences $\{n_j\}$ and $\{m_k\}$, then $i(\lambda, \lambda') < \infty$.*

Proof. If $i(\lambda, \lambda') = \infty$, then for every N there are j, k such that $i(f^{n_j}(\alpha), f^{m_k}(\beta)) \geq N$, by Lemma 3.6. Thus, $i(\alpha, f^{m_k - n_j}(\beta)) \geq N$, so f is not tempered by Lemma 4.1. $\square$

5. SUBSURFACES AND SPANNING SETS OF CURVES

Throughout this section, we fix a hyperbolic structure on S . Given a collection of curves $\mathcal{C}$, we say a curve α is a *concatenation* of curves in $\mathcal{C}$ if it is homotopic to a finite concatenation of geodesic arcs coming from the geodesic representatives of curves in $\mathcal{C}$. The collection $\mathcal{C}$ is *closed under concatenation* if every curve obtained as a concatenation of curves in $\mathcal{C}$ is again an element of $\mathcal{C}$.

We say that a subsurface Σ is *spanned* by $\mathcal{C}$ if:

- (1) every curve in $\mathcal{C}$ is contained in Σ , up to isotopy; and
- (2) every curve $\gamma \subset \Sigma$ is a concatenation of the curves in $\mathcal{C}$

and Σ is minimal (with respect to inclusion, up to proper homotopy) among all subsurfaces satisfying (1) and (2).

Note that if $\mathcal{C}$ spans a subsurface Σ , then necessarily $\mathcal{C}$ fills Σ . On the other hand, a (sub)surface Σ need not coincide with the subsurface spanned by its curves. For instance, on a once-punctured surface of genus two we can consider the subsurface Σ , homeomorphic to a one-holed torus with a point removed from the boundary, as in Figure 4. Here, the subsurface spanned by the curves of Σ is a one-holed torus with compact boundary.




FIGURE 4. A subsurface Σ not spanned by its curves.

We say that X is *spanned by its curves* if it coincides with the subsurface spanned by its curves.

When $\mathcal{C}$ is a finite collection, there is always a subsurface Σ (of finite type) spanned by $\mathcal{C}$. Namely, put the curves in $\mathcal{C}$ in general position, take a regular neighborhood of their union, and fill in any complementary disks with at most one puncture.

In general, one can show that if a subsurface spanned by a collection $\mathcal{C}$ of curves exists, then it is unique up to proper homotopy. To prove this fact, we introduce three subcollections of curves, which we will also use later:

$$\mathcal{C}_p := \{\gamma \in \mathcal{C} \mid \forall \gamma' \in \mathcal{C}, i(\gamma, \gamma') = 0\};$$

$$\mathcal{C}_{\text{int}} := \mathcal{C} \setminus \mathcal{C}_p;$$

$$\mathcal{C}_a := \{\gamma \in \mathcal{C}_p \mid \gamma \text{ is not a concatenation of curves in } \mathcal{C}_{\text{int}}\};$$

If a subsurface spanned by $\mathcal{C}$ exists, the curves in $\mathcal{C}_p$ are peripheral. Among them, the curves in $\mathcal{C}_a$ are precisely those that contribute annular components to the spanned subsurface, while the curves in $\mathcal{C}_p \setminus \mathcal{C}_a$ occur as boundary components

of non-annular components. The curves in $\mathcal{C}_{\text{int}}$ instead are in the interior of the subsurface.

Lemma 5.1. *Let $\mathcal{C}$ be a collection of curves in S . If Σ_1 and Σ_2 are subsurfaces spanned by $\mathcal{C}$, then they are properly homotopic.*

Proof. Fix a hyperbolic structure on S , and assume that all curves in $\mathcal{C}$ are realized as geodesics. Let Σ_i° be the geodesic representative of the interiors of Σ_i , chosen so that $\partial\Sigma_1^\circ$ and $\partial\Sigma_2^\circ$ are in minimal position. Let Σ be an almost geodesic representative of $\Sigma_1^\circ \cap \Sigma_2^\circ$. Then Σ is properly homotopic into both Σ_1 and Σ_2 . We claim that Σ contains all curves in $\mathcal{C}$ and that all curves in Σ are concatenations of curves in $\mathcal{C}$. Then, by minimality of Σ_1 and Σ_2 , they both are properly homotopic to Σ and hence to each other.

It remains to prove the claim.

(1) Let $\gamma \in \mathcal{C}$.

- If $\gamma \notin \mathcal{C}_p$, then γ is a nonperipheral curve of a subsurface contained both in Σ_i . Hence it is contained in both Σ_i° , and therefore $\gamma \subset \Sigma$, up to homotopy.
- If $\gamma \in \mathcal{C}_a$, then γ corresponds to an annular component contained in both Σ_i . Hence it is contained in both Σ_i° , and therefore $\gamma \subset \Sigma$, up to homotopy.
- If $\gamma \in \mathcal{C}_p \setminus \mathcal{C}_a$, then there are finitely many curves in $\mathcal{C} \setminus \mathcal{C}_p$, say $\beta_1, \dots, \beta_k$, such that γ is a concatenation of arcs of the β_i . In particular, γ is contained in the subsurface spanned by $\{\beta_1, \dots, \beta_k\}$, which is contained, up to homotopy, in both Σ_i and hence in Σ_i° . Therefore, $\gamma \subset \Sigma$, up to homotopy.

(2) Let $\gamma \subset \Sigma$. Then γ is up to homotopy contained in both Σ_i . Since each Σ_i is spanned by $\mathcal{C}$, γ is a concatenation of curves in $\mathcal{C}$. $\square$

Note that uniqueness requires the minimality assumption, even for finite collections of curves on finite-type surfaces. For example, in Figure 5, let $\mathcal{C}$ be a pair of curves α and β that intersect once. Then a one-holed torus with a point removed from the boundary satisfies conditions (1) and (2), but it is not minimal. The subsurface spanned by $\mathcal{C}$ is a one-holed torus (with compact boundary).




FIGURE 5. A (non-minimal) subsurface Σ whose curves are concatenations of α and β .

If $\mathcal{C}$ is an infinite collection of curves, then a subsurface spanned by $\mathcal{C}$ does not always exist – see for instance Figure 6. A simple condition that ensures the

existence of a spanned subsurface is that $\mathcal{C}$ be *locally finite*, meaning that every compact set of S intersects only finitely many curves in $\mathcal{C}$. The main goal of this section is to give a condition on $\mathcal{C}$ that guarantees it spans a subsurface. In fact, under the additional assumption that $\mathcal{C}$ is closed under concatenation, we give a necessary and sufficient condition for a collection $\mathcal{C}$ to span a subsurface (Proposition 5.5). This assumption holds for some natural collections of curves, such as curves which are periodic or wandering with respect to a mapping class:

Lemma 5.2. *Let f be a mapping class on S . Every curve obtained as a concatenation of f -periodic curves is periodic, and every curve obtained as a concatenation of wandering curves is wandering.*

Proof. We first consider periodic curves. A curve is a concatenation of curves $\gamma_1, \dots, \gamma_n$ if and only if it is contained in the subsurface F spanned by $\{\gamma_1, \dots, \gamma_n\}$. Note that F is connected and has negative Euler characteristic. Since each γ_i is f -periodic, we can take a suitable power so that f^k fixes γ_i for all i . Since F is spanned by $\{\gamma_i\}$, $f^k(F)$ is spanned by $\{f^k(\gamma_i)\} = \{\gamma_i\}$, so $f^k(F)$ is isotopic to F . Since F has negative Euler characteristic, f^k restricted to F is isotopic to a periodic map of F (see Lemma 2.5), so every curve in F is f -periodic.

The proof for wandering curves is similar. Let F be the subsurface spanned by a finite collection $\{\gamma_1, \dots, \gamma_n\}$ of wandering curves. It suffices to show that for all compact subsurface $K \subset S$, $f^{\pm k}(F)$ can be homotoped to be disjoint from K for all sufficiently large k . Since each γ_i is wandering, we can find k_0 such that $f^{\pm k}(\gamma_i)^* \cap K = \emptyset$ for all $k \geq k_0$ and all i . Then $f^{\pm k}(F)$ is spanned by $\{f^{\pm k}(\gamma_i)\}$, so $f^{\pm k}(F)$ can be homotoped away from K . $\square$




FIGURE 6. The beginning of a collection of curves not spanning a subsurface.

As mentioned before, if $\mathcal{C}$ is a finite collection of curves then there is an explicit way to construct the subsurface spanned by $\mathcal{C}$. Given any collection of curves, our goal is to describe a procedure that yields the subsurface spanned by them, if it exists, and to understand conditions under which it does exist.

Recall the sets $\mathcal{C}_p$, $\mathcal{C}_{\text{int}}$ and $\mathcal{C}_a$ defined before:

$$\mathcal{C}_p := \{\gamma \in \mathcal{C} \mid \forall \gamma' \in \mathcal{C}, i(\gamma, \gamma') = 0\}$$

$$\mathcal{C}_{\text{int}} := \mathcal{C} \setminus \mathcal{C}_p$$

$$\mathcal{C}_a := \{\gamma \in \mathcal{C}_p \mid \gamma \text{ is not a concatenation of curves in } \mathcal{C}_{\text{int}}\}$$

Enumerate the curves in $\mathcal{C}_{\text{int}} = \{\gamma_0, \gamma_1, \dots\}$. Up to reordering, we can assume that $i(\gamma_0, \gamma_1) \neq 0$. Then for every $i \geq 1$, let

$$\mathcal{C}_{\text{int}}^i := \{\gamma_j \mid j \leq i \text{ and } \exists j' \leq i \text{ such that } i(\gamma_j, \gamma_{j'}) \neq 0\}.$$

Note that $\mathcal{C}_{\text{int}}^1 = \{\gamma_0, \gamma_1\}$, $\mathcal{C}_{\text{int}}^i \subset \mathcal{C}_{\text{int}}^{i+1}$, and $\mathcal{C}_{\text{int}} = \bigcup_{i \geq 1} \mathcal{C}_{\text{int}}^i$. Moreover, if $\mathcal{C}_{\text{int}}^i \neq \mathcal{C}_{\text{int}}^{i+1}$, to obtain $\mathcal{C}_{\text{int}}^{i+1}$ we add curves which either intersect curves in $\mathcal{C}_{\text{int}}^i$ or intersect each other. In particular, each component of the subsurface spanned by $\mathcal{C}_{\text{int}}^i$ has negative Euler characteristic.

We now let $F^i = F(\mathcal{C}_{\text{int}}^i)$ be the geodesic representative of the *interior* of the subsurface spanned by $\mathcal{C}_{\text{int}}^i$ and define

$$F = F(\mathcal{C}_{\text{int}}) := \bigcup_{i \geq 1} F^i.$$

By construction every connected component of F^i is an open subset (homotopic to a subsurface), and thus F is an open subset of S . We will call F the *set spanned* by $\mathcal{C}_{\text{int}}$.

We also define $F(\mathcal{C})$ to be the union

$$F(\mathcal{C}) := F(\mathcal{C}_{\text{int}}) \cup \bigcup_{\alpha \in \mathcal{C}_a} A(\alpha),$$

where, for every α , $A(\alpha)$ is a regular neighborhood of the geodesic representative of α , chosen so that they are pairwise disjoint and disjoint from $F(\mathcal{C}_{\text{int}})$.

We will call a collection $\mathcal{C}$ of curves *good* if:

- (a) the curves in $\mathcal{C}_a$ when realized as geodesics, form a locally finite collection, i.e. they do not accumulate anywhere; and
- (b) for every $p \in \partial F$ there is a closed disk B centered at p such that $B \cap F$ is either B minus a diameter, or a connected component of B minus a diameter.




FIGURE 7. A collection of curves which is not good.

For instance, the collection of curves depicted in Figure 6 is not good, because the isolated curves accumulate somewhere. Similarly, the collection in Figure 7 is not good either, because condition (b) is not satisfied at the point p in the drawing (onto which the curves accumulate). In both examples, the collection of curves does not span a subsurface. Indeed, we will show that the condition of being good is equivalent to $\mathcal{C}$ admitting a spanning subsurface (provided $\mathcal{C}$ is closed under concatenation).

Let $K \subset S$ be a finite-type subsurface with compact totally geodesic boundary. Each F^i intersects K in a disjoint union of open subsurfaces and disks with at most one puncture. By a *rectangle* we mean a connected component R of $K \cap F^i$ that is homeomorphic to a closed disk and whose boundary is split into *horizontal sides* lying in ∂F^i and *vertical sides* lying in ∂K . All components of $K \cap F^i$ that are not rectangles are called *essential*. Let K_e^i be the union of all the essential components of $K \cap F^i$.




FIGURE 8. A compact subsurface K , whose boundary is in black, and the boundary of a subsurface F^i in gray. The shaded part is the intersection, formed by an essential component and a rectangle.

Lemma 5.3. *For all $i \geq 2$, $K_e^i \subset K_e^{i+1}$, and for large i it stabilizes. More precisely, there exists an open subsurface $K_e \subset K$ such that for all sufficiently large i , K_e^i is isotopic to K_e .*

Proof. This follows from Euler characteristic considerations. $\square$

So for any finite-type subsurface K with compact totally geodesic boundary, there exists I such that for all $i \geq I$, the topology of the essential components K_e^i stabilizes, and further increasing i only adds new rectangles or adjoins previous ones. We define:

$$\mathcal{R}(K) = \{R \mid R \subset K \cap F^i \text{ is a rectangle for some } i \geq I \text{ and is not contained in any } K_e^i\}.$$

Two parallel rectangles in $\mathcal{R}(K)$ are declared *equivalent* if they are contained in the same connected component of $F^i \cap K$ for some i . The following Lemma is not hard to prove, and we leave it as an exercise.

Lemma 5.4. *The relation defined above is an equivalence relation on $\mathcal{R}(K)$. Moreover, suppose R_1, R_2, R_3 are three parallel rectangles with R_2 contained in the rectangle between R_1 and R_3 . If R_1 and R_3 are equivalent, then all three are equivalent.*

This allows us to naturally organize parallel rectangles into equivalence classes. As i increases, the number of equivalence classes may increase. We have the following proposition.

Proposition 5.5. *Suppose $\mathcal{C}$ is a collection of curves closed under concatenation. The following statements are equivalent:*

- (1) *There is a subsurface spanned by $\mathcal{C}$.*

- (2) *The curves in $\mathcal{C}_a$ do not accumulate anywhere and for every compact subsurface K with totally geodesic boundary, the number of equivalence classes in $\mathcal{R}(K)$ is finite.*
- (3) *$\mathcal{C}$ is good.*

Proof. [(1) $\Rightarrow$ (2)] Let Σ be a surface spanned by $\mathcal{C}$. By contradiction, suppose the curves in $\mathcal{C}_a$ accumulate somewhere. Then we can find a sequence of curves $\gamma_j \in \mathcal{C}_a$ and a curve α such that $i(\gamma_j, \alpha) \neq 0$ for all j . Since Σ is a subsurface, $\Sigma \cap \alpha$ consists of finitely many arcs, with at least one denoted a -intersecting at least three curves, say $\gamma_{j_1}, \gamma_{j_2}$ and γ_{j_3} . Up to replacing a with a subarc and changing the roles of the curves, we can assume that $\gamma_{j_1} \cap a$ and $\gamma_{j_3} \cap a$ are reduced to a point and there is at least one intersection point of γ_{j_2} with a between $\gamma_{j_1} \cap a$ and $\gamma_{j_3} \cap a$.

Then the pair of pants containing $\gamma_{j_1} \cup a \cup \gamma_{j_3}$ is in Σ and contains a curve β which intersects γ_{j_2} essentially. As $\beta \subset \Sigma$, it is a concatenation of curves in $\mathcal{C}$, and since $\mathcal{C}$ is closed under concatenation we have $\beta \in \mathcal{C}$, contradicting the definition of $\mathcal{C}_a$. This shows that the curves in $\mathcal{C}_a$ cannot accumulate anywhere.

Next, suppose there is a finite-type subsurface K with compact totally geodesic boundary such that $\mathcal{R}(K)$ has infinitely many equivalence classes. Then we can find infinitely many parallel rectangles R_i that are not equivalent. Let γ_i be a curve in F passing through R_i . By construction, each γ_i is a concatenation of subarcs of curves in $\mathcal{C}$, hence $\gamma_i \subset \Sigma$. Since Σ is a subsurface, $\Sigma \cap K$ is a finite union of connected components. In particular, there is some component X of $\Sigma \cap K$ containing $\gamma_i \cap K$ for infinitely many i . This implies that we can find distinct curves γ_{i_1} and γ_{i_2} such that the rectangle R between $\gamma_{i_1} \cap K$ and $\gamma_{i_2} \cap K$ is included in Σ . This implies that R is contained in $F^l \cap K$ for some l , so R_{i_1} and R_{i_2} are equivalent, a contradiction.

[(2) $\Rightarrow$ (3)] We just need to verify property (b) in the definition of “good”. By construction, ∂F is a union of limits of simple closed geodesics. We claim that for every finite-type subsurface K with compact totally geodesic boundary, $\partial F \cap K$ has finitely many connected components. Otherwise, as K is of finite type, there are infinitely many components l_j of $\partial F \cap K$ parallel to a given arc from ∂K to ∂K . Then there are infinitely many rectangles between the l_j which are not in F and therefore infinitely many rectangles that are not equivalent, a contradiction.

Therefore, for every $p \in \ell \subset \partial F$ we can find a sufficiently small ball B such that $\ell \cap B$ coincided with $\partial F \cap B$ and it is a single arc from ∂B to ∂B . In particular, $B \setminus \partial F = B \setminus \ell$ has two components, each of which is either entirely contained in F or in $S \setminus F$. Moreover, since $\ell \subset \partial F$, at least one of the components intersects F , and hence is contained in F . So B is the required disk.

[(3) $\Rightarrow$ (1)] The condition on boundary points implies that ∂F contains no non-proper component and that its components do not accumulate anywhere. If $\gamma \subset \partial F$ is a compact boundary component, we can find some finite-type subsurface K with compact boundary components such that $\gamma \subset K$. As K_e^i stabilizes, γ is eventually a boundary component of some F^i , and therefore is an essential curve.

If $\gamma \subset \partial F$ is a non-compact boundary component, since it is the limit of simple closed geodesics, it is geodesic, and therefore it is a line. Moreover, since each component of ∂F is a simple closed geodesic or the limit of simple closed geodesics, for any $\ell \subset \partial F$, either

- for every $p \in \ell$ there is a closed disk B centered at p such that $B \cap F$ is a connected component of a disk minus a diameter, or
- for every $p \in \ell$ there is a closed disk B centered at p such that $B \cap F$ is a disk minus a diameter.

Let T_1 be the collection of lines and curves in ∂F for which the first condition holds, and T_2 be the collection of lines and curve for which the second holds.

For every $\ell \in T_2 \cup \mathcal{C}_a$ we can find a regular neighborhood $N(\ell)$ such that:

- $N(\ell)$ is open if $\ell \in T_2$ and closed if $\ell \in \mathcal{C}_a$;
- $N(\ell)$ is disjoint from $\partial F \cup \mathcal{C}_a \setminus \{\ell\}$;
- if $\ell_1 \neq \ell_2$, then $N(\ell_1) \cap N(\ell_2) = \emptyset$, and
- the neighborhoods don't accumulate anywhere.

We can find such neighborhoods by the properness of components of ∂F , together with the non-accumulation assumption on $\mathcal{C}_a$ (we can fix a compact exhaustion of the surface and construct the neighborhoods piece by piece).

Define

$$\Sigma := \left(F \cup \bigcup_{\ell \in T_1} \ell \cup \bigcup_{\gamma \in \mathcal{C}_a} N(\gamma) \right) \setminus \bigcup_{\ell \in T_2} N(\ell).$$

Then Σ is a two-dimensional submanifold. Its boundary components are either homotopic to boundary components of F (hence are essential curves or lines), or homotopic to curves in $\mathcal{C}_a$ (hence are essential curves). Therefore, Σ is a subsurface.

It is clear that Σ contains all curves in $\mathcal{C}$. Moreover, if γ is a curve in Σ , either it is contained in an annular component, and therefore is in $\mathcal{C}$, or it is homotopic into some F^i (by the same argument for compact components of ∂F), and therefore γ is a concatenation of curves in $\mathcal{C}$. Finally, Σ is properly homotopic to $F \cup \bigcup_{\gamma \in \mathcal{C}_a} N(\gamma)$. Since any surface containing all curves in $\mathcal{C}$ must contain $F \cup \bigcup_{\gamma \in \mathcal{C}_a} \gamma$ up to proper isotopy, Σ satisfies the minimality hypothesis, and hence is a subsurface spanned by $\mathcal{C}$. $\square$

Corollary 5.6. *If X and Y are subsurfaces spanned by two collections of curves $\mathcal{C}_X$ and $\mathcal{C}_Y$ and $i(\alpha, \beta) = 0$ for every $\alpha \in \mathcal{C}_X, \beta \in \mathcal{C}_Y$, then X and Y are disjoint.*

Proof. This follows from the explicit construction of a subsurface spanned by a set of curves given in the proof of Proposition 5.5, together with the uniqueness of spanned subsurfaces (Lemma 5.1). $\square$

We record a consequence of having a non-good collection of curves. This will be our main tool for showing certain collections of curves are good.

Lemma 5.7. *Let $\mathcal{C}$ be a collection of curves closed under finite concatenations. Suppose $\mathcal{C}$ is not good. Then there is a simple closed geodesic α and two infinite sequences of simple closed geodesics $\beta_j, \gamma_j, j \in \mathbb{N}$, with the following properties:*

- (1) $\alpha, \gamma_j \notin \mathcal{C}$ but $\beta_j \in \mathcal{C}$, for all j .
- (2) every β_j intersects α and the β_j 's are pairwise disjoint.
- (3) For every j , there is a subarc $\tau_j \subset \alpha$ joining β_j to β_{j+1} so that the regular neighborhood of the union $\tau_j \cup \beta_j \cup \beta_{j+1}$ is a pair of pants whose third boundary component is γ_j . Furthermore, τ_j 's are pairwise disjoint and τ_j is disjoint from β_k for $k < j$.

Proof. By Proposition 5.5, either there is a compact subsurface K with totally geodesic boundary such that $\mathcal{R}(K)$ contains infinitely many equivalence classes, or the curves in $\mathcal{C}_a$ accumulate somewhere.

In the first case, we can find a sequence $R_i \in \mathcal{R}(K)$ which are all parallel and pairwise not equivalent. Let α be a boundary curve of K containing a vertical side of each R_i . Up to passing to a subsequence, we can assume that there is an oriented arc $[p, q] \subset \alpha$ and a sequence of points $v_i \in R_i \cap [p, q]$ converging monotonically to q . For every i we can moreover find a curve $\delta_i \in \mathcal{C}$ which is a boundary component of some surface F^{l_i} and intersecting K in a collection of arcs that includes a horizontal arc of R_i . This boundary curve is in $\mathcal{C}$ since the collection is closed under concatenation. In this way all δ_i 's are pairwise disjoint since the F^{l_i} 's are nested.

After a subsequence, we can assume that the intersections $R_i \cap \alpha$ converge to a point $q \in \alpha$ and that the δ_i 's don't contain q . After a further subsequence there will be an oriented arc $[p, q] \subset \alpha$ that contains a vertical side of each R_i and the convergence to q is monotonic along $[p, q]$. Further, any arc of any $\delta_i \cap K$ that intersects $[p, q]$ is contained in some rectangle. Inductively, we can construct a subsequence δ_{i_j} such that

- (i) there is a strand of $\delta_{i_{j+1}}$ below (i.e. closer to q) all strands of δ_{i_j} , and
- (ii) between the lowest strand of δ_{i_j} and the first strand of $\delta_{i_{j+1}}$ below δ_{i_j} there are at least two other rectangles.

Let τ_j be the subarc of $[p, q]$ connecting the two strands of δ_{i_j} and $\delta_{i_{j+1}}$ in (ii). Let γ_j be the third boundary component of a small regular neighborhood of $\delta_{i_j} \cup \tau_j \cup \delta_{i_{j+1}}$. Since τ_j intersects another δ_k , γ_j is essential. Moreover, if $\gamma_j \in \mathcal{C}$, then the rectangles associated with the two strands of δ_k 's would be equivalent, which is a contradiction. Now rename $\beta_j := \delta_{i_j}$.

If the curves in $\mathcal{C}_a$ accumulate somewhere, we can find a curve α intersecting infinitely many $\delta_i \in \mathcal{C}_a$. As before, up to passing to a subsequence, we can find an arc $[p, q] \subset \alpha$, with some $v_i \in \delta_i \cap [p, q]$ converging to q monotonically. We then define δ_{i_j} , γ_j and τ_j as above. Since each γ_j intersects a curve in $\mathcal{C}_a$, $\gamma_j \notin \mathcal{C}$.

Note that in both cases there may be strands of β_k that intersect τ_j when $k > j + 1$. $\square$




FIGURE 9. On the left-hand side, the arc $a = [p, q]$ for the non-good collection of curves of Figure 6; on the right-hand side, the general situation in the proof of Lemma 5.7.

Remark 5.8. In the situation of the previous lemma, and given a homeomorphism f , fix an index j . Look at $f^n(\alpha)$, $f^n(\beta_j)$ and $f^n(\beta_{j+1})$ and consider an ambient isotopy which sends them to their geodesic representatives. The image through this isotopy of $f^n(\tau_j)$ is a geodesic arc of $f^n(\alpha)^*$ that we denote by τ_j^n . In particular, $f^n(\gamma_j)^*$ is homotopic to $f^n(\beta_j)^* \cup \tau_j^n \cup f^n(\beta_{j+1})^*$.

6. DIAGONAL CLOSURE OF CURVES AND LINES

In this section we discuss two subsurfaces associated to a finite collection of curves and lines: the subsurface spanned by the collection and the diagonal closure of the collection.

Given a finite collection of lines and curves L , the subsurface *spanned* by $\mathcal{L}$, denoted $\langle L \rangle$, is the subsurface obtained by putting the curves and lines in minimal position, taking a regular neighborhood of their union, and then filling in all complementary disks with at most one puncture.

Given two rays r_1 and r_2 , we say that they *cobound a half-strip* if there is a proper embedding $\varphi : \mathbb{R}_{\geq 0} \times [0, 1] \rightarrow S$ such that $\varphi(\mathbb{R}_{\geq 0} \times \{0\}) = r_1$ and $\varphi(\mathbb{R}_{\geq 0} \times \{1\}) = r_2$. We say that two lines ℓ_1 and ℓ_2 are *asymptotic* in some direction if they contain rays $r_1 \subset \ell_1$ and $r_2 \subset \ell_2$ which cobound a half-strip.




FIGURE 10. Two asymptotic lines with the half-strip cobounded by two rays.

Let L be a finite collection of curves or proper lines such two elements of L intersect only finitely many times, and no line in L ends in an isolated planar end.

We denote by $\langle\langle L \rangle\rangle$ the subsurface of S obtained as follows. First put the curves and lines in L in minimal position. Next form the subsurface $\langle L \rangle$ spanned by L . Then, for every two (not necessarily distinct) lines and every direction in which they are asymptotic, adjoin to $\langle L \rangle$ the corresponding half-strip cobounded by the asymptotic rays. Finally fill in any additional complementary disks with at most one puncture. We call $\langle\langle L \rangle\rangle$ the *diagonal closure* of L . The following lemma gives a characterization of $\langle\langle L \rangle\rangle$.

Lemma 6.1. *Let $L = \{\ell_1, \dots, \ell_N\}$ be a finite collection of curves and lines such that no element ends in an isolated planar end of S , and any two elements of L intersect only finitely many times. Then for any line ℓ that does not end in an isolated planar end, the following statements are equivalent.*

- (1) ℓ can be homotoped into $\langle\langle L \rangle\rangle$.
- (2) For any curve α , if $i(\alpha, \ell) \neq 0$, then $i(\alpha, \ell_j) \neq 0$ for some j .
- (3) For every compact subsurface K , ℓ can be homotoped to ℓ' such that $\ell' \cap K \subset \langle L \rangle$.

Moreover there is a finite union of curves μ such that if ℓ is a line in $\langle\langle L \rangle\rangle$, then $i(\mu, \ell) \neq 0$.

Proof. We first prove the equivalence of the three conditions.

[(1) $\Rightarrow$ (3)] We may assume $\ell \subset \langle L \rangle$. Since $\langle\langle L \rangle\rangle$ is formed from $\langle L \rangle$ by adjoining finitely many half-strips and disks, we can first homotope ℓ away from all the adjoined disks. For any compact subsurface K , there is a homotopy that pushes all the half-strips off of K . Let ℓ' be the image of ℓ under this homotopy. Then $\ell' \cap K \subset \langle L \rangle$.

[(3) $\Rightarrow$ (2)] Let α be a curve with $i(\alpha, \ell) \neq 0$. Let K be an annular neighborhood about α and ℓ' a line homotopic to ℓ so that $\ell' \cap K \subset \langle L \rangle$. Since ℓ' intersects α essentially, α crosses $\langle L \rangle$ essentially. Therefore, there is some j such that $i(\alpha, \ell_j) \neq 0$.

[(2) $\Rightarrow$ (1)] We prove the contrapositive: if ℓ cannot be homotoped into $\langle\langle L \rangle\rangle$, then there is some curve α that intersects ℓ but disjoint from all ℓ_j 's.

Assume all lines are geodesic and ℓ is in minimal position with respect to $\langle\langle L \rangle\rangle$. Note that $\langle\langle L \rangle\rangle$ is the union of a compact surface with finitely many half-strips. In particular the boundary of $\langle\langle L \rangle\rangle$ is a finite union of circles and lines. The same therefore holds for $\Sigma := \overline{S \setminus \langle\langle L \rangle\rangle}$. Moreover, Σ contains no pairs of boundary rays cobounding a half-strip in S , since any such half-strip would be included in the construction of $\langle\langle L \rangle\rangle$.

Let a be a component of $\ell \cap \Sigma$ and F the component of Σ containing a . If a is nonseparating in F , we can find a curve contained in F intersecting a once and we are done. If a is separating and both components of $F \setminus a$ are not contractible, choose two non-contractible curves α_1 and α_2 (which may be homotopic to a puncture), one in each component of $F \setminus a$, and a simple arc b connecting them. The regular neighborhood of $\alpha_1 \cup b \cup \alpha_2$ is a pair of pants whose third boundary component gives the required curve α . Thus we may assume that a is separating

that intersect $\langle L \rangle$ and other half-strips only in the regular nbhds of boundary rays –mb

Not sure about the comment –ff

and at least one component C of $F \setminus a$ is contractible. By the classification of contractible surfaces with boundary, $C \cup a$ is a closed disk with points removed from its boundary. Since $\partial\Sigma$ has only finitely many components, $C \cup a$ is in fact a closed disk with finitely many points removed from its boundary. As ℓ cannot be homotoped into $\langle\langle L \rangle\rangle$, one such point is not an end of a . It follows that Σ contains a half-strip cobounded by two boundary rays, a contradiction (see Figure 11). This finishes the proof of this case.

Finally, we prove the last statement. Write

$$\langle\langle L \rangle\rangle = K \sqcup S_1 \sqcup \cdots \sqcup S_k,$$

where K is a compact surface and the S_i are pairwise disjoint half-strips. For every $j = 1, \dots, k$, choose a ray r_j going through S_j . As no line ends in a puncture, there is a sequence of essential separating curves, pairwise not homotopic, converging to the end of r_j contained in S_j . Choose one such curve μ_j sufficiently far out so that μ_j intersects the ray r_j and is disjoint from K . Let

$$\mu = \bigcup_{j=1}^k \mu_j.$$

If ℓ is a line in $\langle\langle L \rangle\rangle$, then by properness ℓ has to exit K along some ray r_j , hence it intersects the corresponding μ_j . This shows $i(\mu, \ell) \neq 0$ as desired. $\square$




FIGURE 11. The situation in the proof of Lemma 6.1.

Remark 6.2. At first one might think that condition (2) in Lemma 6.1 is equivalent to ℓ being in $\langle L \rangle$ (up to proper homotopy). The example depicted in Figure 12 shows that this is not the case: condition (2) holds, but ℓ cannot be properly homotoped into $\langle L \rangle$.

The reason why we are interested in Lemma 6.1 is the following relationship between $\langle\langle \cdot \rangle\rangle$ and limits of concatenations of curves.

Lemma 6.3. *Suppose f is well tempered. Let $\alpha_1, \dots, \alpha_k$ be curves. If β is a concatenation of $\alpha_1, \dots, \alpha_k$, then*

$$\mathcal{L}^\pm(\beta)k \subset \left\langle \left\langle \bigcup_{i=1}^n \mathcal{L}^\pm(\alpha_i) \right\rangle \right\rangle.$$




FIGURE 12. L is the union of the gray-green lines and ℓ is the black one. Then ℓ is not properly homotopic into $\langle L \rangle$, but it is in $\langle\langle L \rangle\rangle$.

Proof. Let $\ell \in \mathcal{L}^+(\beta)$ and γ a curve intersecting ℓ . By Lemma 3.6 there are infinitely many $n \geq 0$, such that $i(\gamma, f^n(\beta)) \neq 0$. Since β is a finite concatenation of the α_i 's, there is some $j \in \{1, \dots, n\}$ and infinitely many $n \geq 0$ such that $i(f^n(\alpha_j), \gamma) \neq 0$. Applying Lemma 3.6 again, we obtain some $\ell' \in \mathcal{L}^+(\alpha_j)$ with $i(\ell', \gamma) \neq 0$. Since γ was arbitrary, Lemma 6.1 implies that $\ell \subset \langle\langle \bigcup_{i=1}^k \mathcal{L}^\pm(\alpha_i) \rangle\rangle$ up to proper homotopy. The same argument applies to $\mathcal{L}^-(\beta)$. $\square$

We end this section with a variant of Lemma 6.1.

Lemma 6.4. *Let $A \subset B$ be two finite collections of curves and lines such that any two elements intersect only finitely many times. Then there exist finitely many curves $\delta_1, \dots, \delta_s$ disjoint from lines in A , with the following property: if a line $\ell \subset \langle\langle B \rangle\rangle$ cannot be homotoped into $\langle\langle A \rangle\rangle$, then $i(\ell, \delta_i) \neq 0$ for some i .*

Proof. As in the proof of Lemma 6.1, we can write $\langle\langle B \rangle\rangle$ as the union of a finite-type subsurface K with compact boundary and pairwise disjoint half-strips $S_1, \dots, S_k$. Since $A \subset B$, we can homotope $\langle\langle A \rangle\rangle$ so that it is the union of a finite type subsurface $K' \subset K$ with compact boundary, together with some of the half-strips S_i ; up to renumbering, say $S_1, \dots, S_l$.

For each $l < j \leq k$, choose a curve δ_{j-l} associated to the half-strip S_j as in the proof of Lemma 6.1. We append to this collection the set of boundary curves $\delta_{k-l+1}, \dots, \delta_s$ of K' .

Now suppose a line ℓ in $\langle\langle B \rangle\rangle$ cannot be homotoped into $\langle\langle A \rangle\rangle$. If ℓ goes through a half-strip not contained in $\langle\langle A \rangle\rangle$, then ℓ intersects one of the curves δ_i with $i \leq k - l$. Otherwise, $\ell \setminus K$ is contained in $\langle\langle A \rangle\rangle$, so $\ell \cap K$ cannot be contained in K' and hence it intersects some δ_i with $i > k - l$. $\square$

7. WELL-TEMPERED MAPS: CANONICAL DECOMPOSITION

By a *decomposition* of S we mean a collection $\{X_i\}$ of pairwise disjoint subsurfaces of S (each of which may be disconnected), such that we can find representatives $\tilde{X}_i$ with $S = \bigcup \tilde{X}_i$ and such that the only overlap between $\tilde{X}_i$ and $\tilde{X}_j$, $i \neq j$, are common boundary components. When f is a homeomorphism of S , a decomposition $\{X_i\}$ of S is called *f-invariant* if $f(X_i)$ is isotopic to X_i for all i .

For a subsurface $X \subset S$, we say that f *returns to* X if there is a power $k > 0$ such that $f^k(X)$ is isotopic to X . Any such $k > 0$ is called a *return time* of f to X . Note that all return times of f to X are multiples of the smallest return time.

Our main proposition in this section is the following.

Proposition 7.1. *Suppose f is well tempered. Then there is an f -invariant decomposition of S into three subsurfaces, S_{per} , S_{∞} , and S_0 , with the following properties.*

- *A curve in S is periodic if and only if it can be homotoped into S_{per} .*
- *A curve in S is wandering if and only if it can be homotoped into S_{∞} .*
- *S_0 contains no essential, nonperipheral curves.*

We will call S_{per} , S_{∞} , and S_0 the canonical decomposition of f on S .

An explicit example of a well-tempered map is the map induced by the matrix $\begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}$ on $S = \mathbb{R}^2 \setminus \mathbb{Z}^2$. In this case, the decomposition is given by four invariant lines, which cobound a subsurface $X = S_0$ homeomorphic to a once-punctured disk with four points removed from the boundary. The rest of the surface is S_{∞} , consisting of four invariant components (see Figure 13).




FIGURE 13. The four invariant lines giving the decomposition of Proposition 7.1 for the map $(x, y) \mapsto (2x + y, x + y)$ on $S = \mathbb{R}^2 \setminus \mathbb{Z}^2$.

For any surface S and any well-tempered map f , the decomposition is defined as follows. Let $\mathcal{C}_{\text{per}}$ be the collection of f -periodic curves, and $\mathcal{C}_{\infty}$ the collection of f -wandering curves. Let S_{per} be the subsurface spanned by $\mathcal{C}_{\text{per}}$ (whose existence is proved in Lemma 7.4), S_{∞} the subsurface spanned by $\mathcal{C}_{\infty}$ (whose existence is proved in Lemma 7.2).

Fix a hyperbolic metric on S . We choose an almost geodesic representative for the components of S_{per} and S_{∞} as follows. For each non-annular component X , let X^* be the almost geodesic representative of X . Recall this means that we first take the geodesic representative X° of the interior of X and then remove a small regular neighborhood of a boundary component of X° that is homotopic to another boundary component of X° . We also assume that if two components share any boundary component, their almost geodesic representatives do as well.

By abuse of notation, we continue to use S_{per} and S_{∞} for the union of the representatives of their components. Define

$$S_0 := \overline{S \setminus (S_{\text{per}} \cup S_{\infty})},$$

whose isotopy class is independent of the choice of the hyperbolic metric on S .

As just seen, a fundamental step in proving Proposition 7.1 is to show that, for a well-tempered map, $\mathcal{C}_{\text{per}}$ and $\mathcal{C}_{\infty}$ span subsurfaces S_{per} and S_{∞} . We then need to show that S_0 has no interesting topology.

All three proofs proceed by contradiction. Namely, we will show that if the required statement is false, then Lemma 5.7 can be used to produce an essential curve α with infinite limit set $\mathcal{L}(\alpha)$. The arguments for S_{per} and S_{∞} are independent. The proof for S_0 is more technical: it relies on the fact that S_0 is already a subsurface, being the complement of two subsurfaces, and on the observation that no curve in S_0 can be periodic or wandering, hence any curve in S_0 must limit onto a non-empty collection of lines.

Let us now prove that S_{∞} and S_{per} are subsurfaces.

Lemma 7.2. *If f is well tempered, then $\mathcal{C}_{\infty}$ spans a subsurface S_{∞} . Every curve in S_{∞} is wandering, and if γ is any curve with $\mathcal{L}(\gamma) \neq \emptyset$, then $\mathcal{L}(\gamma)$ does not intersect S_{∞} essentially.*

Proof. Suppose there is no subsurface spanned by $\mathcal{C}_{\infty}$. By Lemma 5.2, $\mathcal{C}_{\infty}$ is closed under concatenation, so we can apply Lemma 5.7 and find curves α , β_j and γ_j and pairwise disjoint arcs $\tau_j \subset \alpha$ as in the lemma. Recall that $\beta_j \in \mathcal{C}_{\infty}$, but $\alpha, \gamma_j \notin \mathcal{C}_{\infty}$. In particular, $\mathcal{L}(\gamma_j)$ and $\mathcal{L}(\alpha)$ are nonempty. Since γ_j is homotopic to $\beta_j \cup \tau_j \cup \beta_{j+1}$ (and none of these is periodic, by assumption and Corollary 3.9), Lemma 6.3 gives

$$\mathcal{L}(\gamma_j) \subset \langle\langle \mathcal{L}(\alpha) \cup \mathcal{L}(\beta_j) \cup \mathcal{L}(\beta_{j+1}) \rangle\rangle = \langle\langle \mathcal{L}(\alpha) \rangle\rangle,$$

where the last equality uses the fact that the β_i are wandering, hence $\mathcal{L}(\beta_i) = \emptyset$. Since $\mathcal{L}(\alpha)$ is a finite collection of lines with finite pairwise intersections, Lemma 6.1 provides a finite union of curves μ such that every line in $\langle\langle \mathcal{L}(\alpha) \rangle\rangle$ intersects μ . Hence there is a curve $\delta \subset \mu$ that intersects infinitely many $\mathcal{L}(\gamma_j)$. By Lemma 3.7, $\mathcal{L}(\delta)$ intersects infinitely many γ_j . Since $\mathcal{L}(\delta)$ is finite, some line $\ell \in \mathcal{L}(\delta)$ intersects infinitely many γ_j .

If $i(\ell, \beta_j) \neq 0$, then $i(\delta, \mathcal{L}(\beta_j)) \neq 0$, which is a contradiction since β_j is wandering; the same applies to β_{j+1} . Therefore ℓ must intersect infinitely many of the arcs τ_j , and hence intersects α infinitely many times. This is impossible since ℓ is a proper line. This contradiction shows that $\mathcal{C}_{\infty}$ spans a subsurface S_{∞} .

For the second statement, if γ is a curve such that $\mathcal{L}(\gamma)$ intersects S_{∞} essentially, then there exists $\beta \subset S_{\infty}$ such that $i(\beta, \mathcal{L}(\gamma)) \neq 0$. But every curve in S_{∞} is a concatenation of wandering curves, hence β is wandering by Lemma 5.2. This contradicts Lemma 3.8. $\square$

We also need the following generalization.

Lemma 7.3. *Suppose f is well tempered. Let L^+ and L^- be two finite collections of lines with finite pairwise intersections, and such that each of L^+ and L^- has finite intersection with every $\mathcal{L}^\pm(\alpha)$. Define*

$$\mathcal{C}_L := \{\beta \mid \mathcal{L}^\pm(\beta) \subseteq \langle\langle L^\pm \rangle\rangle\}.$$

Then $\mathcal{C}_L$ spans a subsurface S_L of S .

Proof. Note that $\mathcal{C}_L$ contains all wandering curves, and is closed under concatenation by Lemma 6.3.

Assume that $\mathcal{C}_L$ does not span a subsurface. Then we have curves $\alpha, \beta_j, \gamma_j$ and arcs $\tau_j \subset \alpha$ as in Lemma 5.7, where $\beta_j \in \mathcal{C}_L$; $\alpha, \gamma_j \notin \mathcal{C}_L$; and γ_j is homotopic to $\beta_j \cup \tau_j \cup \beta_{j+1}$. In this situation,

$$\mathcal{L}^\pm(\gamma_j) \subset \langle\langle \mathcal{L}^\pm(\alpha) \cup \mathcal{L}^\pm(\beta_j) \cup \mathcal{L}^\pm(\beta_{j+1}) \rangle\rangle \subseteq \langle\langle \mathcal{L}^\pm(\alpha) \cup L^\pm \rangle\rangle.$$

Let $\{\delta_l^+\}_l$ and $\{\delta_m^-\}_m$ be the curves from Lemma 6.4 for the collections $L^\pm \subset L^\pm \cup \mathcal{L}^\pm(\alpha)$. After passing to a subsequence and possibly replacing f by f^{-1} and switching $+$ and $-$, we may assume that for every j there is at least one line in $\mathcal{L}^+(\gamma_j)$ that is not homotopic into $\langle\langle L^+ \rangle\rangle$, hence intersects one of the $\{\delta_l^+\}_l$. Passing to a further subsequence, we may assume that the same curve $\delta \in \{\delta_l^+\}_l$ intersects infinitely many $\mathcal{L}^+(\gamma_j)$. By construction, δ does not intersect any of $\mathcal{L}^+(\beta_j)$. It follows by Lemma 3.7 that $\mathcal{L}^-(\delta)$ must intersect α infinitely many times, which is impossible since the map is well tempered. $\square$

Lemma 7.4. *If f is well tempered, then $\mathcal{C}_{\text{per}}$ spans a subsurface S_{per} , and every curve in S_{per} is periodic.*

Proof. By Lemma 5.2, $\mathcal{C}_{\text{per}}$ is closed under concatenation. Suppose, for contradiction, there is no subsurface spanned by $\mathcal{C}_{\text{per}}$. Then we can find curves α, β_j and γ_j and arcs $\tau_j \subset \alpha$ as in Lemma 5.7. Recall that $\beta_i \in \mathcal{C}_{\text{per}}$, but $\alpha, \gamma_j \notin \mathcal{C}_{\text{per}}$. Moreover, since α intersects periodic curves, it is not wandering either. Thus $\mathcal{L}(\alpha)$ is non-empty and consists of lines.

Each line in $\mathcal{L}(\alpha)$ is the limit of a uniformly bounded number of strands of $f^{n_i}(\alpha)^*$. Let N be the cardinality of $\mathcal{L}(\alpha)$, counted with multiplicity, and set $J = N + 1$. Consider the finite collection of curves $\gamma_1, \dots, \gamma_J$. By replacing f by a power, we may assume $f(\beta_j) = \beta_j$ for all $j \leq J + 1$.

Recall that γ_j is homotopic to $\beta_j \cup \tau_j \cup \beta_{j+1}$. For each n , let τ_j^n be the geodesic subarc of $f^n(\alpha)^*$ which together with $f^n(\beta_j)^* = \beta_j$ and $f^n(\beta_{j+1})^* = \beta_{j+1}$ is homotopic to $f^n(\gamma_j)^*$ (see Remark 5.8). Since γ_j is not periodic, the arcs τ_j^n cannot be periodic. As there are only finitely many isotopy classes of arcs of bounded length between β_j and β_{j+1} , the length of τ_j^n cannot be bounded as $n \rightarrow \infty$.

Moreover, the angles of intersections of τ_j^n with β_j and β_{j+1} are bounded away from zero; otherwise some sequence of iterates of α would wrap around β_j or β_{j+1} more and more times, contradicting the tempered condition. Thus any convergent subsequence of τ_j^n contains a ray starting at β_j and a ray starting at β_{j+1} . By construction, each such ray is contained in a line of $\mathcal{L}(\alpha)$.

Choose a subsequence f^{n_i} so that each (oriented) $\tau_j^{n_i}$ converges. Then the union of all these limit sets contains at least $2J = 2N + 2$ rays. If a collection of k such limiting rays is nested within the same line in $\mathcal{L}(\alpha)$, then the multiplicity of that line is at least k . Taking into account both orientations of each line, we see that there can be at most $2N$ rays, a contradiction. This shows the existence of S_{per} . $\square$

Lemma 7.5. *If f is well tempered, then S_{per} and S_∞ are disjoint subsurfaces.*

Proof. Since they are subsurfaces spanned by curves, if they are not disjoint, by Corollary 5.6 there are curves α in S_{per} and β in S_∞ with $i(\alpha, \beta) > 0$, contradicting Corollary 3.9. $\square$

Lemma 7.6. *If f is well tempered, $S_0 := \overline{S \setminus (S_{\text{per}} \cup S_\infty)}$ has no essential, non-peripheral curves.*

Proof. Suppose, by contradiction, that there is an essential, nonperipheral curve δ in S_0 . Then δ is neither wandering nor periodic, so $L := \mathcal{L}(\delta)$ is a nonempty collection of lines. Let $L^\pm := \mathcal{L}^\pm(\delta)$ and let S_L be the subsurface from Lemma 7.3 spanned by the collection of curves $\{\gamma\}$ with $\mathcal{L}^\pm(\gamma) \subset \langle\langle L^\pm \rangle\rangle$. Put S_L and S_0 in minimal position and let $\tilde{S}_L = S_0 \cap S_L$. Note that $S_\infty \subseteq S_L$ and $S_{\text{per}} \cap S_L$ is a compact surface, so the boundaries of S_L and S_0 intersect in only finitely many points.

Up to replacing f by its inverse, we may assume L^+ is nonempty. By Lemma 6.1, there exists a finite collection of curves $\{\delta_1, \dots, \delta_k\}$ intersecting all lines in $\langle\langle L^+ \rangle\rangle$. Then Lemma 3.7 implies that for every curve γ in S_L , either

$$(a) \mathcal{L}^+(\gamma) = \emptyset, \quad \text{or} \quad (b) i(\gamma, \mathcal{L}^-(\delta_i)) \neq 0 \text{ for some } i.$$

Claim: There exists an essential, nonperipheral curve in $\tilde{S}_L$ satisfying (a).

Indeed, otherwise the finite collection of curves and lines in $\bigcup_i \mathcal{L}^-(\delta_i)$ fills every curve in $\tilde{S}_L$. Since there are only finitely many intersection points between these curves and lines, it follows that each component of $\tilde{S}_L$ has finite type. Furthermore, since $\tilde{S}_L$ is f -invariant, if one of its components is not periodic, then its orbit is a subsurface of $\tilde{S}_L$. In particular the orbit of the component cannot accumulate anywhere; that is, it is wandering. Now consider the component that contains δ . If it is wandering, δ is wandering, which is impossible. Hence this component must be periodic. But then the first return map is a tempered mapping class of a finite-type surface; in particular, it is periodic and so δ is periodic as well, which is again impossible. This proves the claim.

By the claim, there exists an essential, nonperipheral curve δ' in S_0 that is forward wandering. Replace f by f^{-1} and run the whole construction on δ' , obtaining the subsurfaces $S_{L'}$ and $\tilde{S}_{L'}$, where $L' = L^\pm(\delta')$. Now, every curve γ in $S_{L'}$ is f -forward wandering (since $\mathcal{L}^+(\gamma) \subset \langle\langle \mathcal{L}^+(\delta') \rangle\rangle = \emptyset$). The above claim produces another essential, nonperipheral curve in $\tilde{S}_{L'}$ which is f -backward wandering, and thus wandering, a contradiction. $\square$

A consequence of the lemma is the following.

Corollary 7.7. *Any negative Euler characteristic component of S_0 is a pair of pants.*

Proof. Let Y be a component of S_0 with $\chi(Y) < 0$. If $\chi(Y) < -1$, then Y contains an essential nonperipheral curve, a contradiction. Hence $\chi(Y) = -1$. If Y were not a pair of pants, then it would be a pair of pants with some boundary points removed, and such a surface still contains an essential nonperipheral curve. This contradiction proves that Y is a pair of pants. $\square$

7.1. Examples. We end this section with some examples of tempered but not well tempered maps for which the conclusions of Proposition 7.1 are false.

Example 7.8. This example shows the existence of a tempered map f such that $\mathcal{L}(\alpha)$ is locally finite for every curve α , but the collection of periodic curves does not span a subsurface.

Consider the cylinder $\Sigma := S^1 \times \mathbb{R}$ and fix a Cantor set $K \subset S^1$. Let S be the surface obtained as

$$S := \Sigma \setminus \bigcup_{n=-\infty}^{\infty} K \times \{t_n\},$$

where $t_{\pm\infty} = \pm 1$, $t_0 = 0$, $t_n = -1 + \frac{1}{2^{-n}}$ for $n < 0$ and $t_n = 1 - \frac{1}{2^n}$ for $n > 0$ (see Figure 14).

Fix a monotone non-decreasing homeomorphism $s : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\begin{aligned} s(t_{\pm\infty}) &= t_{\pm\infty} \\ s(t_k) &= t_{k+1} \quad \forall k \in \mathbb{Z}. \end{aligned}$$

Then the map $\text{id} \times s : \Sigma \rightarrow \Sigma$ induces a tempered homeomorphism f of S , which has no wandering curves. The periodic curves are also fixed curves, which are those that can be homotoped to be vertical between height -1 and height 1 . In particular $\mathcal{C}_{\text{per}}$ does not span a subsurface.

Example 7.9. This example shows that there exists a tempered map f such that $\mathcal{L}(\alpha)$ is locally finite for every curve α , but $\mathcal{C}_{\infty}$ does not span a subsurface. It also shows that such an example can be constructed so that a single component of S_{∞} is not a subsurface.

In the plane, for any $k \in \mathbb{N}$, let ℓ_k be the boundary of

$$[k+1, \infty) \times \left[\frac{1}{k+1} + \varepsilon_k, \frac{1}{k} - \varepsilon_k \right],$$

where $\varepsilon_k = \frac{1}{10k^2}$ (see Figure 15 for a schematic picture). Let S be the plane punctured at the points $(m, \frac{1}{n})$, for $n \in \mathbb{N}$ and $m \in \mathbb{Z}$, at the points $(0, m)$, for $m \in \mathbb{Z}$, and for every line ℓ_k , $k \in \mathbb{N}$, at a sequence of points on the line not accumulating anywhere.

The homeomorphism f is defined as follows:




FIGURE 14. A sketch of the surface in Example 7.8, with a periodic curve (in gray) and the Cantor sets in light gray.




FIGURE 15. The lines ℓ_k in the construction of Example 7.9.

- on a small regular neighborhood (of width less than ε_k) of each horizontal line $y = \frac{1}{n}$ ($n \in \mathbb{N}$) and $y = 0$, the map is induced by $(x, y) \mapsto (x + 1, y)$;
- for each k , it performs a puncture shift supported on a small regular neighborhood of ℓ_k , tapered to the identity on the rest of the surface.

One can show that f is tempered. Moreover, $F(\mathcal{C}_\infty)$ is the disjoint union of one strip per line $y = \frac{1}{n}$, $n \in \mathbb{N}$, and of the strips $(k, \infty) \times \left(\frac{1}{k+1} + \frac{\varepsilon_k}{2}, \frac{1}{k} - \frac{\varepsilon_k}{2}\right)$. The closure of each component is a subsurface, but the horizontal strips accumulate onto $y = 0$, so $\mathcal{C}_\infty$ does not span a subsurface. Note moreover that there are no periodic curves.

Finally, we can modify the surface by adding handles connecting consecutive horizontal strips in a translation-invariant way (see Figure 16 for a schematic picture). The induced map on this new surface is still tempered and has no periodic curves. In this case, $F(\mathcal{C}_\infty)$ consists of one component per line ℓ_k and

a single nonplanar component. This nonplanar component is not homotopic to a subsurface.




FIGURE 16. The modified surface in Example 7.9.

8. CHARACTERIZATION OF TRANSLATIONS

Recall a homeomorphism $f : X \rightarrow X$ is called a *translation* if it generates an infinite cyclic group acting properly on X . In this section, we give a characterization of the mapping class of a translation in terms of the dynamics of its action on curves. The main application will be to a well tempered map f and its action on the invariant subsurface S_∞ in its canonical decomposition. Therefore, in this section, we need to work with surfaces with boundary, and we will prove a more general version of Theorem B. Everything in this section is independent of the previous sections.

Extending the notation for curves, when δ is an essential arc in a hyperbolic surface with totally geodesic boundary, we denote by δ^* the geodesic arc homotopic to δ which is perpendicular to the boundary at both endpoints.

Lemma 8.1. *Let X be a connected surface, possibly with boundary. Let f be a homeomorphism of X such that every curve in X is f -wandering. Then there are ends $e_\pm$ of X (possibly $e_+ = e_-$) such that for every curve α in X , $f^n(\alpha)^*$ converges to $e_\pm$ as $n \rightarrow \pm\infty$.*

We call e_+ and e_- the f -attracting and f -repelling ends of X , respectively.

Proof. Let α be a nonperipheral curve in X . We first show that $f^n(\alpha)^*$ converges to an end. Let

$$\alpha^* = \alpha_0, \alpha_1, \dots, \alpha_k = f(\alpha)^*$$

be a chain of geodesic curves in X connecting α^* and $f(\alpha)^*$, i.e. $i(\alpha_i, \alpha_{i+1}) \neq 0$ for every i . For any compact set $K \subset X$, there exists n_0 such that for all $n \geq n_0$ and all $i \in \{1, \dots, k\}$, the curves $f^n(\alpha_i)^*$ are disjoint from K . Since consecutive ones intersect, they must lie in the same connected component of $X \setminus K$. In particular, $f^n(\alpha)^*$ and $f^{n+1}(\alpha)^*$ lie in the same complementary component of K , for all $n \geq n_0$. As this holds for all compact sets of X , it follows that $f^n(\alpha)^*$ converges to a single end e_+ as $n \rightarrow \infty$. The same argument applied to f^{-1} shows that $f^n(\alpha)^*$ converges to an end e_- as $n \rightarrow -\infty$.

Now let β be another nonperipheral curve in X . Choose a chain of curves connecting α^* to β^* and repeat the same argument as above to deduce that $f^n(\beta)^* \rightarrow e_{\pm}$ as $n \rightarrow \pm\infty$.

Finally, if γ is a peripheral curve, then it can be expressed as a concatenation of nonperipheral curves (since X is spanned by its curves), all of which converge under forward (respectively backward) iteration to e_+ (respectively e_-), and hence so does γ . $\square$

Lemma 8.2. *Let X be a connected surface, possibly with boundary, and let e_+, e_- be two distinct ends of X . Then there is either a curve or a finite collection of disjoint arcs in X separating e_+ from e_- .*

Proof. Let DX be the double of X along its boundary (and set $DX = X$ if $\partial X = \emptyset$). Let $i : X \hookrightarrow DX$ be the inclusion map and $i_* : \text{Ends}(X) \rightarrow \text{Ends}(DX)$ the induced map. Note that i_* is injective: if $e_1 \neq e_2$ are ends of X , there is a compact subset K separating them, and therefore $i_*(e_1)$ and $i_*(e_2)$ are separated by the double of K in DX .

Since DX has no boundary, there is a curve $\hat{\delta} \subset DX$ separating $i_*(e_+)$ and $i_*(e_-)$. Assume $\hat{\delta}$ is in minimal position with respect to $i(\partial X)$ and let $\delta := i^{-1}(\hat{\delta})$. Then δ is either a curve in X or a finite collection of pairwise disjoint arcs. Moreover, δ separates e_+ and e_- , because if not, there would exist a line ℓ in X from e_+ to e_- disjoint from δ , and hence a line $i(\ell)$ from $i_*(e_+)$ to $i_*(e_-)$ disjoint from $\hat{\delta}$, contradicting that $\hat{\delta}$ separates the ends. $\square$

Proposition 8.3. *Let X be a connected bordered surface spanned by its curves. Let f be a homeomorphism of X such that every curve is f -wandering and every boundary line is f -periodic. Then every essential arc γ in X is f -wandering. Moreover, γ converges to the f -attracting end under forward iteration and to the f -repelling end under backward iteration.*

Proof. Fix a complete hyperbolic structure on X with totally geodesic boundary and positive injectivity radius. For any arc γ , denote by γ^* its geodesic representative which is orthogonal to the boundary of X . We must show that $f^n(\gamma)^*$ leaves every compact set of X .

It is enough to show the statement for arcs whose endpoints lie on noncompact boundary components of X . Indeed, suppose first that γ has endpoints on peripheral curves α and β , possibly $\alpha = \beta$. Then the regular neighborhood of $\alpha \cup \beta \cup \gamma$ contains an essential curve. Since every essential curve under iteration converges to the same attracting and repelling ends of X , it follows that γ is also wandering. Next suppose that γ has one endpoint on a peripheral curve α and the other endpoint on a boundary line ℓ . Then a regular neighborhood of $\alpha \cup \gamma$ contains an arc with both endpoints on ℓ . If γ' also converges to the same attracting/repelling ends of X as curves, then the same is true for γ .

Henceforth, we may assume that γ is an arc with both endpoints lying on noncompact boundary components of X . As for curves, denote by $\mathcal{L}^{\pm}(\gamma)$ the

Removed the assumption that there are no compact boundary and added a paragraph in the proof to deal with the new cases. – jt

accumulation sets of γ_n^* as $n \rightarrow \pm\infty$. Applying the same argument as in Lemma 3.7 to arcs, we have that

$$i(\mathcal{L}(\gamma), \beta) \neq 0 \iff i(\mathcal{L}(\beta), \gamma) \neq 0,$$

where β is any curve or arc in S . Since curves in X are wandering and X is spanned by its curves, $\mathcal{L}(\gamma)$ cannot contain any geodesic segment that meets the interior of X essentially. Hence, if $\mathcal{L}(\gamma) \neq \emptyset$, then it is a union of boundary components. It follows that the endpoints of γ_n^* must escape to infinity along ∂X , for otherwise $\mathcal{L}(\gamma)$ would contain an arc or a ray perpendicular to ∂X .

Case 1. Both endpoints of γ are on the same oriented boundary line ℓ and f preserves the oriented line ℓ . In this case, we close up γ along ℓ to obtain a curve α . Assume for contradiction that $\mathcal{L}^+(\gamma) \neq \emptyset$, i.e. γ is not forward wandering. Then there is a compact set $K \subset X$ and a sequence $n_i \rightarrow \infty$ such that $f^{n_i}(\gamma)^* \cap K \neq \emptyset$ for all i .

Note that the distance between the endpoints of $f^{n_i}(\gamma)^*$ goes to 0 as $i \rightarrow \infty$. Indeed, otherwise after a further subsequence the curve $f^{n_i}(\alpha)^*$ would lie in a fixed Hausdorff neighborhood of the arc $f^{n_i}(\gamma)^*$. This would imply that $f^{n_i}(\alpha)^*$ would intersect a fixed neighborhood of K , contrary to the assumption that α is wandering.

Up to homotopy, we can express $f^{n_i}(\gamma)^*$ as $\delta_{n_i} = \tau_{n_i} * f^{n_i}(\alpha)^* * \bar{\tau}_{n_i}$ for a geodesic arc τ_{n_i} perpendicular at the endpoints to ℓ and $f^{n_i}(\alpha)^*$ respectively. If the lengths of the τ_{n_i} are uniformly bounded from above, then δ_{n_i} also lies close to $f^{n_i}(\alpha)^*$, so δ_{n_i} must go to infinity since α is wandering. We may now assume that the lengths of the τ_{n_i} are going to infinity. The length of $f^{n_i}(\alpha)^*$ is bounded from below, by the injectivity radius assumption. Since the endpoints of $f^{n_i}(\gamma)^*$ lie close, by basic hyperbolic geometry, $f^{n_i}(\gamma)^*$ and δ_{n_i} must lie uniformly close to each other. As $f^{n_i}(\gamma)^*$ intersects K but $f^{n_i}(\alpha)^*$ goes to infinity, for all sufficiently large i , the arc τ_{n_i} must intersect a fixed neighborhood L of K .

Now define a possibly immersed curve β by joining δ_0 and δ_1 along an arc in ℓ and then taking a boundary of a regular neighborhood. The argument from Lemma 5.2 also shows that immersed curves are wandering. Therefore, for all sufficiently large i , the geodesic representative $f^{n_i}(\beta)^*$ lies outside of L . Let x_i be the intersection point of τ_{n_i} with $f^{n_i}(\alpha)^*$. The discussion above implies that for all sufficiently large i , there exists an arc η_i joining x_i to x_{i+1} lying outside of L , such that $f^{n_i}(\beta)^*$ is homotopic to $f^{n_i}(\alpha)^* * \eta_i * f^{n_{i+1}}(\alpha)^*$. On the other hand, $f^{n_i}(\beta)^*$ is homotopic to $\delta_{n_i} * l_i * \delta_{n_{i+1}}$ for some $l_i \subset \ell$. It follows that the arc $\bar{\tau}_{n_i} * l_i * \tau_{n_{i+1}}$ is homotopic to η_i relative to endpoints. In particular, the initial segment of τ_{n_i} from ℓ to its last intersection point with L is homotopic to the corresponding initial segment of $\tau_{n_{i+1}}$, relative to ℓ and ∂L . By repeating the same argument replacing n_i by $n_i + 1$, we see that the initial segment of τ_{n_i} is homotopic to the initial segment of τ_{n_j} for all $j \geq i$. This is impossible, since these segments meet ℓ orthogonally while their endpoints on ℓ escape to ∞ . This contradiction shows that γ is forward wandering.

rewrote this section. Can we change the arc η to join the endpoints of the τ arcs, and have the arc lie outside the set L ? Hmm, the index error seems to be a problem. -jt




FIGURE 17. Schematic pictures in Case 1

Case 2. Both endpoints of γ are on a periodic line ℓ . Then apply Case 1 to the power of f that preserves the oriented line ℓ .

General case. We have seen that the limit set of an arc γ , if non-empty, can only consist of boundary lines. If ℓ is a boundary line in $\mathcal{L}(\gamma)$, let β be an essential arc with both endpoints on ℓ . By the previous cases, β is wandering, so by Lemma 3.7

$$i(\mathcal{L}(\beta), \gamma) = 0 \iff i(\beta, \mathcal{L}(\gamma)) = 0.$$

However, β intersects $\ell \in \mathcal{L}(\gamma)$, a contradiction.

Finally, since every pair of arcs can be connected by a sequence of curves, it follows that every arc converges to the f -attracting end under forward iteration and to the f -repelling end under backward iteration. This finishes the proof. $\square$

Corollary 8.4. *Let X be a connected bordered surface spanned by its curves. Let f be a homeomorphism of X such that every curve is f -wandering and every boundary line is f -periodic. Let DX be the double of X , and let Df be the double of f to DX . Then:*

- (1) *every curve in DX is Df -wandering;*
- (2) *if e_+ and e_- are the f -attracting and f -repelling ends of X , then each determines a unique end $\hat{e}_+$ and $\hat{e}_-$ of DX , and these are precisely the Df -attracting and Df -repelling ends of DX , respectively.*

Proof. For (1), every curve of DX that is not contained in X meets each copy of X in a collection of arcs. By the previous proposition, every arc in this collection is wandering towards the same end of X . Therefore every curve in DX is wandering.

For (2), by Lemma 8.1, there are unique ends $\hat{e}_\pm$ towards which iterates of curves converge. As iterates of curves in X converge to $e_\pm$, the claim follows. $\square$

Lemma 8.5. *Let X be a connected surface, possibly with boundary, and let ϕ be an infinite-order mapping class on X . Suppose that for some $n > 0$ there exists a representative $g \in \phi^n$ that is a translation on X . Then ϕ is also represented by a translation. In particular, there is a hyperbolic metric on X with totally geodesic boundary for which some representative $f \in \phi$ is an isometric translation.*

Moreover, we can choose the hyperbolic metric so that the first return of f to any boundary component of X has translation length 1.

Proof. Fix a representative $f \in \phi$. Note that since $f^n \sim g$, the maps f and g commute up to isotopy. Consider the covering map

$$p : X \rightarrow Q = X/\langle g^2 \rangle.$$

Note that Q may be either of finite or of infinite type. Our goal is to show ϕ induces a well-defined mapping class $\bar{\phi}$ on Q , using the curve graph of Q . By replacing n by a positive multiple if necessary, we may assume that Q has high enough complexity so that the automorphism group of its curve graph coincides with the extended mapping class group of Q [18, 20, 21, 9]. In what follows, we write $h \sim h'$ to mean that h and h' are isotopic.

We first treat the case in which Q has no boundary. For convenience, fix a hyperbolic metric on the further quotient $Q/\langle g \rangle$ and equip X with the lifted metric, so that g acts on X by an isometry. Let $\alpha \subset Q$ be a simple closed geodesic. If α lifts, then $p^{-1}(\alpha) = \bigcup \alpha_m$ is a disjoint union of simple closed geodesics, where $\alpha_m = g^{2m}(\alpha_0)$ for some chosen lift α_0 . Let β_m be the geodesic representative of $f(\alpha_m)$. Since $fg \sim gf$, we have $\beta_m = g^{2m}(\beta_0)$. It follows that there is a unique simple closed geodesic $\beta \subset Q$ such that $\beta = p(\beta_1)$. If α does not lift, then $p^{-1}(\alpha)$ is a finite collection of lines, say $\{l_1, \dots, l_k\}$, which are cyclically permuted by the action of g^2 . Let t_i be the geodesic representative of $f(l_i)$. As for curves, the relation $fg \sim gf$ implies that g^2 cyclically permutes $\{t_1, \dots, t_k\}$. Hence there is again a unique simple closed geodesic $\beta \subset Q$ such that $\beta = p(\bigcup t_i)$. This allows us to define $f_*(\alpha) = \beta$. It is straightforward to check that f_* preserves disjointness, and hence determines an automorphism of the curve graph of Q . Hence f_* is induced by an extended mapping class¹ $\bar{\phi}$ of Q . By construction, $\bar{\phi}$ depends only on the mapping class of f .

Since $f^n \sim g$, the mapping class $\bar{\phi}$ is periodic, and in fact $\bar{\phi}^n = \bar{g}$, where $\bar{g}$ is the involution on Q induced by g . By Proposition 2.7, there is a hyperbolic metric on Q for which $\bar{\phi}$ admits an isometric representative. Lift both the metric and the isometry to X , choosing the lift $\tilde{f} : X \rightarrow X$ such that $\tilde{f}^n \sim g$. In particular, $\tilde{f}$ is an isometric translation of X .

We now claim $\tilde{f} \sim f$. Let F_0 be a fundamental domain for g , and $F_i = g^i(F_0)$. Since f and $\tilde{f}$ induce the same mapping class on Q , our choice of the lift implies that $f^{-1} \circ \tilde{f}$ is isotopic to the identity on each $F_{2i} \cup F_{2i+1}$. Moreover, since both $\tilde{f}$ and f commute with g up to isotopy, $f^{-1} \circ \tilde{f}$ is also isotopic to the identity on $F_{2i+1} \cup F_{2i+2}$. To complete the proof, we use the Alexander method (see [17] for the version on infinite-type surfaces): we choose an Alexander system such that each curve is contained in some $F_j \cup F_{j+1}$. Since $f^{-1} \circ \tilde{f}$ fixes every curve in this system up to isotopy, by the Alexander method, $f^{-1} \circ \tilde{f}$ is isotopic to the identity.

¹I.e. the homotopy class of a homeomorphism, which could be orientation-reversing.

Now suppose that Q has boundary. Double Q and X along their boundaries to obtain DQ and DX , each equipped with an involution that exchanges the two copies. Also, double the maps $Df, Dg: DX \rightarrow DX$, and we still have $(Df)^n \sim Dg$, with the homotopy commuting with the involution of DX . Since $DQ = DX/\langle Dg^2 \rangle$ has no boundary, the previous argument applies to $Df: DX \rightarrow DX$. Thus Df induces a periodic mapping class $\hat{\phi}$ on DQ . Because Df is obtained by doubling f , the mapping class $\hat{\phi}$ commutes with the involution $\tau: DQ \rightarrow DQ$. By Proposition 2.7, after choosing a suitable hyperbolic metric on DQ , we may represent $\hat{\phi}$ by an isometry that commutes with τ . It follows that this isometry preserves the fixed-point set of τ , namely the doubled boundary ∂Q , and hence restricts to an isometry $\tilde{f}$ of Q representing $\hat{\phi}$. In particular, the boundaries of Q are totally geodesic. Moreover, since any hyperbolic metric on DQ preserved by τ is the doubling of a hyperbolic metric on Q , we may prescribe, for each $\tilde{f}$ -orbit of compact boundary components of Q , the common boundary length to be any positive number L . Lift both the metric and the isometry to X , choosing the lift $\tilde{f}: X \rightarrow X$ such that $\tilde{f}^n \sim g$. By a similar argument as above, $\tilde{f} \sim f$.

Finally, let l be a boundary component of X , and suppose that $\tilde{f}^k$ is the first return to l . Then l projects to a compact boundary component of Q of length L . By choosing $L = 2n/d$, where $d = \gcd(2n, k)$, we can arrange that the first return of $\tilde{f}$ to l is a translation by length 1. $\square$

Theorem 8.6. *Let X be a connected possibly bordered surface spanned by its curves. Let f be a homeomorphism of X such that every curve is f -wandering and every boundary line is f -periodic. Then there is a hyperbolic metric on X with totally geodesic boundary and an isometric translation on X isotopic to f .*

Remark 8.7. If the surface has empty boundary, Theorem 8.6 reduces to Theorem B.

Proof. By Lemma 8.1, there are ends e_+ and e_- of X to which all curves converge under forward/backward iteration. In particular, $f(e_{\pm}) = e_{\pm}$. We now distinguish two cases.

Case 1: $e_+ \neq e_-$. Let δ be a curve or a minimal multi-arc separating e_+ and e_- as in Lemma 8.2. Thus we have $X = X_- \cup_{\delta} X_+$ where $X_{\pm}$ is a neighborhood of $e_{\pm}$. Choose $n \neq 0$ so that $f^n(X_+) \subset X_+$. Each $\delta_i = f^{ni}(\delta)^*$ is multiarc or a curve separating e_+ and e_- with complementary components nested. This shows that the dual graph is a line on which f^n acts as a translation. We may therefore isotope f^n to a translation g of X , with the region between δ_0 and δ_1 serving as a fundamental domain for the action of g . This case is finished by Lemma 8.5.

Case 2: $e_+ = e_-$. We first treat the case in which X has no boundary, and then reduce the bordered case to it by doubling.

The goal is to construct an exhaustion of X by f -invariant subsurfaces, each of which satisfies the hypothesis of Lemma 8.5. Choose a curve α_1 and then choose a finite sequence of curves

$$\alpha_1, \alpha_2, \dots, \alpha_k = f(\alpha_1)$$

such that $i(\alpha_i, \alpha_{i+1}) > 0$. Since every curve in X is wandering, the union of the $\langle f \rangle$ -orbits of the curves α_i forms a locally finite graph Γ . Therefore, they span a subsurface T_1 which is f -invariant up to isotopy. We claim f is isotopic to a translation of T_1 . Since the curves are all wandering, there exists $n_0 \geq 1$ such that α_i is disjoint from $f^n(\alpha_j)$, for all $n \geq n_0$ and $i, j \in \{1, \dots, k\}$. It follows that the induced action by $\langle f \rangle \cong \mathbb{Z}$ on Γ is proper and cocompact, so Γ is two-ended. The regular neighborhood $N(\Gamma)$ is also two-ended, and since curves in $N(\Gamma)$ are f -wandering, these ends are the attracting and repelling endpoints. Applying Case 1 to $N(\Gamma)$ we obtain a translation f_1 of $N(\Gamma)$ isotopic to $f|_{N(\Gamma)}$. The subsurface T_1 is obtained from $N(\Gamma)$ by capping off some compact boundary components with disks or once-punctured disks, and possibly attaching half-planes along noncompact boundary components. Any disk or punctured-disk attachments occur in $\mathbb{Z}$ -orbits, hence the translation extends over them. Furthermore, since T_1 is spanned by finitely many curves, it has only finitely many noncompact boundary components. So some power of f_1 fixes each of them and acts on each as a translation. It follows that some power of f_1 also extends across any attached half-plane as a translation. This shows some power of f is isotopic to a translation of T_1 , and hence the same is true of f by Lemma 8.5, proving the claim. Repeating this procedure for a larger collection of curves yields a sequence of subsurfaces $T_1 \subset T_2 \subset \dots$ exhausting X , each is f -invariant up to isotopy, and f is isotopic to a translation on each T_i . See Figure 18.




FIGURE 18. An example of the exhaustion $T_1 \subset T_2 \subset \dots$, where the map is the horizontal shift to the right, T_1 is the surface spanned by the orbit of the “horizontal” curve and T_2 the surface spanned by the orbit of the “horizontal” and the “vertical” curve.

We now construct, for each i , a hyperbolic metric on T_i with totally geodesic boundary extending the hyperbolic metric on T_{i-1} together with a map $f_i \sim f$ preserving T_i such that

- $f_i|_{T_i}$ is an isometric translation;
- the first return of f to any noncompact boundary of T_i is a translation of length 1; and

$$\bullet \quad f_i|_{T_{i-1}} = f_{i-1}|_{T_{i-1}}.$$

The base case $i = 1$ comes from Lemma 8.5. Now assume $i \geq 2$, and choose $g \sim f$ such that $g(T_i) = T_i$ and $g|_{T_{i-1}} = f_{i-1}|_{T_{i-1}}$. Let Y be the closure of a component of $T_i \setminus T_{i-1}$. Since T_i is spanned by finitely many f -orbits of curves, there is a smallest $k > 0$ such that $g^k(Y) = Y$. Let $\mathcal{Y} = \bigsqcup_{i=1}^k g^i(Y)$, regarded as a possibly disconnected surface. The restriction $g|_{\mathcal{Y}}$ is isotopic to a translation $\mathcal{Y}$, in the sense that the first return to each component is a translation. Therefore, there is a hyperbolic metric on $\mathcal{Y}$ together with an isometry $g_{\mathcal{Y}} \sim g|_{\mathcal{Y}}$, such that the first return of $g_{\mathcal{Y}}$ to any noncompact boundary of any $g^i(Y)$ is a translation by length 1. Gluing the hyperbolic metric on T_{i-1} to the one on $\mathcal{Y}$ via the identifications coming from X yields a hyperbolic metric on $T_{i-1} \cup \mathcal{Y}$. Moreover, along any common boundary between T_{i-1} and $\mathcal{Y}$, the maps f_{i-1} and $g_{\mathcal{Y}}$ have first returns that are translations of length 1 in the same direction. Hence these maps glue to an isometric translation on $T_{i-1} \cup \mathcal{Y}$ isotopic to $g|_{T_{i-1} \cup \mathcal{Y}}$. Performing this construction for each g -orbit of components of $T_i \setminus T_{i-1}$ yields a hyperbolic metric on T_i extending the one on T_{i-1} , together with an isometry on T_i extending the translation on T_{i-1} . We can now isotope f to a map f_i that agrees with this translation on T_i .

Passing to the limit, we obtain a hyperbolic metric on X with respect to which f is isotopic to an isometric translation.

Finally, in the case that X is bordered, let DX be the double of X , and let Df be the doubled homeomorphism. By Corollary 8.4, every curve in DX is Df -wandering, and the common attracting and repelling end e of X determines an end $\hat{e}$ of DX such that every curve in DX converges to e under both forward and backward iteration of Df . Thus DX with the map Df satisfies the hypotheses of the boundaryless case above. This gives a hyperbolic metric on DX and an isometric translation g isotopic to Df . All of these choices can be made equivariantly with respect to the involution exchanging the two copies of X . Hence the resulting metric is invariant under this involution, and its restriction to X is a hyperbolic metric with totally geodesic boundary. With respect to this restricted metric, $g|_X$ acts by a translation on X and is isotopic to f . $\square$

Remark 8.8. In the previous proof, in the case $e_+ = e_-$, passing to the double is necessary if X has nonempty boundary. This is because the subsurfaces T_i may only exhaust the interior of X , in which case we do not obtain a hyperbolic structure on all of X (some boundary components would lie at infinity with respect to the metric on the interior of X).

We end this section with an example showing that the non-wandering boundary-line hypothesis is necessary in Proposition 8.3, Corollary 8.4 and Theorem 8.6.

Example 8.9. Let R be the Roller compactification (see [27]) of $\mathbb{R}^2$ with its standard cube complex structure. Let X be the surface obtained by removing from R the closure of all points in $\mathbb{R}^2$ with integer coordinates. The homeomorphism f of X is the continuous extension of the map $(x, y) \mapsto (x + 1, y + 1)$ of $\mathbb{R}^2$. Then

every curve of X and every boundary line is wandering. On the other hand, there are arcs from boundary to boundary which are not wandering, such as the closure of the line $y = \frac{1}{2} \subset \mathbb{R}^2 \setminus \mathbb{Z}^2$. In particular, the map induced by f on DX is not a translation.




FIGURE 19. The surface X from Example 8.9, with a non-wandering horizontal arc

9. STRUCTURE THEOREM OF WELL-TEMPERED MAPS

The goal of this section is to prove the following classification result for well-tempered maps:

Theorem 9.1. *Let f be a well-tempered homeomorphism of a surface S of infinite type. Then, up to modifying f by an isotopy, there is a hyperbolic metric on S and a canonical decomposition of S into three f -invariant subsurfaces S_{per} , S_{∞} and S_0 with totally geodesic boundary such that, for every component X of S_{per} , S_{∞} and S_0 , the map f returns to X and the first return is isotopic to:*

- *a periodic isometry if $X \subseteq S_{\text{per}}$ or $X \subseteq S_0$;*
- *an isometric translation if $X \subseteq S_{\infty}$.*

Furthermore, each component X of S_0 is topologically trivial, i.e. X contains no essential nonperipheral curves.

We begin by developing several properties of the canonical decomposition associated to a well-tempered map f . In particular, we describe how components of S_{per} , S_{∞} , and S_0 can neighbor each other. We also show that, for each component X of the decomposition, the map f returns to X . We then combine these results with those of Section 8 and the work of Afton-Calegari-Chen-Lyman [3] to complete the proof of Theorem 9.1. Finally, we show how to deduce Theorem A from Theorem 9.1.

9.1. Properties of the canonical decomposition. For a well-tempered map, recall the almost geodesic representatives of S_{per} , S_{∞} , and S_0 defined in Section 7. An essential component of S_{per} or S_{∞} either has infinite type or has negative Euler characteristic; in particular, such a component always contains an essential curve.

Two components of the decomposition are *adjacent* if they have boundary components that are properly isotopic (equivalently, they have representatives sharing at least one boundary component).

A component X is *self-adjacent* along α if α is properly homotopic to two distinct boundary components of X . Equivalently, X is self-adjacent if there is an inessential component Y which shares two boundary components with X .

In this section we will show various properties of the canonical decomposition. The first result excludes strip components and self-adjacency (and will follow from Lemma 9.15 and Corollary 9.16).

Proposition 9.2. *No component of the canonical decomposition is a strip, nor is any component self-adjacent.*

The next proposition describes the components of S_{∞} , and it is proven over Lemmas 9.8 and 9.13:

Proposition 9.3. *Let X be a connected component of S_{∞} . Then:*

- (1) *no boundary component of X is compact;*
- (2) *every boundary component of X is periodic;*
- (3) *f returns to X and there is a hyperbolic structure on X such that the first return map is isotopic to an isometric translation.*

The following properties of S_{per} are shown in Lemmas 9.10 and 9.12:

Proposition 9.4. *No two components of S_{per} are adjacent. If X is a component of S_{per} , f returns to X and there is a hyperbolic structure on X such that the first return map is isotopic to a finite-order isometry.*

Finally, we describe the properties of S_0 :

Proposition 9.5. *The following are the topological possibilities for a component X of S_0 :*

- (1) *A closed disk with at least three points removed from the boundary. In this case, at most one neighbor of X is in S_{per} .*
- (2) *A once-punctured closed disk with at least one point removed from the boundary. In this case, all neighbors of X are in S_{∞} .*
- (3) *A cylinder with one compact boundary component and at least one point removed from the other boundary (a crown). In this case, X has exactly one neighbor in S_{per} along its compact boundary component.*
- (4) *A pair of pants. In this case, all neighbors of X are annular components of S_{per} .*

Note that all topological possibilities from Proposition 9.5 occur for components of S_0 , as shown in Figure 20. In each case, the map is a puncture shift in each shaded strip, in the direction indicated by the arrow, and it is the identity outside the strips.




FIGURE 20. Examples of well-tempered mapping classes

The rest of the section is dedicated to showing the previous propositions, whose proofs are intertwined. We start by showing that certain boundary components have to be periodic.

Lemma 9.6. *If two components of S_∞ share a common boundary component α , then α is periodic. Similarly, if a component of S_∞ is self-adjacent along α , then α is periodic.*

Proof. Let X and Y be (not necessarily distinct) components of S_∞ with a common boundary component α . Since X and Y are essential, we can find a curve $\beta \subset X \cup Y$ that intersects α essentially. Because β is not contained in S_∞ , it is not wandering, so $\mathcal{L}(\beta) \neq \emptyset$. Choose $\ell \in \mathcal{L}(\beta)$ and let γ be a curve intersecting ℓ . Since $i(\gamma, \mathcal{L}(\beta)) \neq 0$, Lemma 3.7 implies $i(\mathcal{L}(\gamma), \beta) \neq 0$, hence γ is not wandering. By Lemma 7.2, $\mathcal{L}(\gamma)$ cannot intersect any component of S_∞ (in particular, X and Y) essentially. On the other hand, $\mathcal{L}(\gamma)$ intersects β . This is only possible if $\alpha \in \mathcal{L}(\gamma)$. By f -invariance, the entire orbit of α lies in $\mathcal{L}(\gamma)$. As f is well tempered, the orbit of α is finite, i.e. α is periodic. $\square$

Lemma 9.7. *Let X be a component of S_0 . If X either*

- *shares at least one boundary component with S_∞ but contains a non-contractible curve not homotopic to that boundary component; or*

• *shares at least two boundary components with S_∞ ,*
then X is not wandering.

Proof. In the first case, let γ be a non-contractible curve in X and let Y be a component of S_∞ sharing a boundary component α with X , where γ is not homotopic to α . Since Y is essential, we can find a curve $\beta \subset X \cup Y$ crossing α , for instance by joining a curve in Y to γ along an arc and taking the boundary of a regular neighborhood of their union. In the second case, let Y and Y' be components of S_∞ (possibly $Y = Y'$) sharing boundary components α and α' , respectively, with X . We can then find a curve $\beta \subset X \cup Y \cup Y'$ intersecting α and α' , by joining a curve in Y to a curve in Y' by an arc that crosses α and α' .

We claim that if X is wandering, then β is wandering, which is a contradiction. The argument is similar to above. Indeed, suppose β is not wandering. Then $\mathcal{L}(\beta) \neq \emptyset$, but by Lemma 7.2, $\mathcal{L}(\beta)$ cannot intersect $f^n(Y)$ or $f^n(Y')$ essentially. On the other hand, $f^n(\beta)$ is contained in $f^n(X \cup Y)$ or $f^n(X \cup Y \cup Y')$, so some line $\ell \in \mathcal{L}(\beta)$ must be homotopic into $f^n(X)$ for some n . Since $\mathcal{L}(\beta)$ is f -invariant and X is wandering, infinitely many iterates of ℓ would lie in $\mathcal{L}(\beta)$, contradicting the finiteness of $\mathcal{L}(\beta)$. $\square$

We now show that all boundary components of S_∞ are lines.

Lemma 9.8. *No component X of S_∞ has a compact boundary component.*

Proof. We argue by contradiction. Suppose that α is a compact boundary component of X . Since X is spanned by curves, α is a wandering curve. Therefore the component Y adjacent to X along α cannot belong to S_{per} (since boundary curves of S_{per} are periodic) or to S_∞ (by Lemma 9.6). Hence Y is a component of S_0 .

If there is no $n > 0$ such that $f^n(Y)$ is isotopic to Y , then Y is wandering (as S_0 is an f -invariant subsurface). By Lemma 9.7, Y cannot contain a non-contractible curve different from α . If α were the only boundary component of Y , then Y is a closed disc with at most one puncture, which is impossible. Thus Y has at least two boundary components, both wandering. Hence these two boundary components belong to S_∞ , contradicting Lemma 9.6.

Therefore there exists $n > 0$ such that $f^n(Y)$ is isotopic to Y . Since α is wandering, Y contains infinitely many iterates of α , and hence contains an essential nonperipheral curve. This contradicts Proposition 7.6, completing the proof. $\square$

Using the previous results, we can deduce that no component of S_0 is wandering; moreover, for each such component the first return map has finite order.

Lemma 9.9. *For every component X of S_0 , f returns to X and there is a hyperbolic structure in which every compact boundary component has length one, such that the first return f^k is isotopic to a periodic isometry of X .*

Proof. Recall that X has no essential, nonperipheral curves, and by our construction of S_{per} and S_∞ , X cannot be a closed disk or a closed disk with one

puncture. If f does not return to X , then since S_0 is a subsurface and f -invariant, X is wandering. In particular, X cannot border S_{per} , so it cannot have a compact boundary component by Lemma 9.8. In this case, X is a disk with at most one puncture and points removed from the boundary, which yields a contradiction by Lemma 9.7. Hence f returns to X .

Up to replacing f with a power, we may assume $f(X) = X$. If X has finitely many boundary components, then f has finite order. Otherwise, X is a disk with infinitely many points removed from the boundary and possibly a puncture or an open disk removed from its interior. Consider the action of f on the noncompact boundary components of X . If the f -orbit of every component is infinite, then f is semi-conjugate to an irrational rotation of the disk, which violates the fact that f is well tempered. Thus, there exists a finite orbit; in that case, all periodic components of ∂X have the same period, and the same holds for the periodic ends. Up to passing to a power, we can assume all periodic boundary components and all periodic ends are fixed.


Suppose by contradiction that there is a wandering boundary component ℓ . By the structure of orientation-preserving homeomorphisms of the circle, there are ends $e_{\pm}$ (possibly $e_+ = e_-$) of X such that $f^n(\ell) \rightarrow e_{\pm}$ as $n \rightarrow \pm\infty$. Let Y be the component of the canonical decomposition adjacent to X along ℓ . Since one of its boundary components is wandering, $Y \subset S_{\infty}$. Let α be a curve in $\text{int}(X) \cup \text{int}(Y) \cup f(\text{int}(Y)) \cup \ell \cup f(\ell)$ which intersects ℓ essentially (see Figure 21). Note that α is not wandering, so $\mathcal{L}(\alpha) \neq \emptyset$. Without loss of generality, suppose $\mathcal{L}^+(\alpha) \neq \emptyset$. Let ℓ_0 be the line in X spanned by e_+ and the end of ℓ such that $f^i(\ell)$, $i \geq 0$, lies on one side of $Y \setminus \ell_0$, which also contains α . Let Z_0 be the subsurface corresponding to that side (see Figure 21), and let $Z_i = f^i(Z_0)$. Then Z_i converges to e_+ as $i \rightarrow \infty$. But $\mathcal{L}^+(\alpha) \subset Z_i \cup S_{\infty}$ for every $i \geq 0$, and $\mathcal{L}^+(\alpha)$ cannot intersect S_{∞} essentially, so we get a contradiction. Let k be the first return time to X . Then we obtain a hyperbolic metric on X with respect to which f^k is isotopic to a finite order isometry by looking at the quotient $X/\langle f^k \rangle$, choosing a hyperbolic structure with corners on the quotient and lifting it to X . $\square$

We can now establish the properties of the first return map for components of S_{per} .

this is already stated in
Cor 7.7 –mb
removed the statement –jt

Lemma 9.10. *Let X be a component of S_{per} . Then f returns to X and there is a hyperbolic metric on X with compact boundary components of length one such that the first return f^k is isotopic to a periodic isometry of X .*

Proof. Since X is connected and every curve in X is f -periodic, f must return to X . Up to replacing f by a power we may assume $f(X)$ is isotopic to X . If X has bounded topology (i.e. X is finite type or the double of X is of finite type), then f is isotopic to a periodic map of X . Otherwise, choose a finite collection $\mathcal{C}$ of orbits of curves in X and let $F = F(\mathcal{C})$ be the subsurface spanned by $\mathcal{C}$. After possibly enlarging $\mathcal{C}$, we can assume F is connected and has negative Euler characteristic. Since $\mathcal{C}$ is f -invariant so is F ; moreover, since all curves in $\mathcal{C}$ are f -periodic, $f|_F$

FIGURE 21. The situation in the proof of Lemma 9.9, for $e_- \neq \varepsilon_+$.

is isotopic to a periodic map of F by Lemma 2.5. Let $n > 0$ be the period of $f|_F$. For any curve α not in F , we can find a larger connected f -invariant finite-type subsurface F' containing both F and α . By the same reasoning, f is also isotopic to a periodic map on F' , but since $F' \supset F$, $f|_{F'}$ also has period n (by classical Nielsen-Thurston theory). In particular, $f^n(\alpha)$ is isotopic to α for all α . By the Alexander method ([17]), this implies that $f|_X$ is isotopic to a periodic map of X . The fact that we can realize the first return f^k to X by a periodic isometry is a direct consequence of Proposition 2.7 (see [3]). $\square$

As a consequence, we can describe more precisely the properties of pair-of-pants components of S_0 .

Corollary 9.11. *Each pair of pants component of S_0 is adjacent only to annular components of S_{per} . Furthermore, no component of S_0 is self-adjacent.*

Proof. Since a pair of pants component X of S_0 has only compact boundary components, it can only border components of S_{per} . Let Y be one such component; if Y is not an annulus, we can construct a curve $\alpha \subset X \cup Y$ intersecting the boundary of X essentially. As some power of f is periodic on both X and Y , and as by the tempered condition there is no twisting allowed along the boundary, α is periodic, a contradiction.

If a component X of S_0 were self-adjacent along a boundary component α , then α must be a curve—otherwise there is a component Y which is a strip with both boundary components in X . Such a component Y must belong to S_0 , since it has no curves, but then $X \cup Y$ would be part of the same component of S_0 . If α is a curve, then by Lemma 9.9 α is periodic, so the annulus Z about α is a component of S_{per} . Since X has at least two compact boundary components, it is a pair of pants, but then as before we could construct a curve β contained in the genus-one surface $X \cup Z$ which intersects α essentially and is periodic, a contradiction. $\square$

Our next step is to exclude (self-)adjacency of components of S_{per} .

Lemma 9.12. *No two components of S_{per} can be adjacent, nor can a component of S_{per} be self-adjacent.*

Proof. First suppose X and Y are essential components of S_{per} and α is a common boundary component. Choose a regular neighborhood $N(\alpha)$ about α contained in $X \cup Y$. There exists n such that f^n is isotopic to the identity on X and Y . Isotope f^n so that it takes $N(\alpha)$ to $N(\alpha)$ fixing $\partial N(\alpha)$ pointwise, and further isotope f^n so that it is the identity on truncated subsurfaces $X \setminus N(\alpha)$ and $Y \setminus N(\alpha)$. If $N(\alpha)$ is a strip, then we can further isotope f^n (rel boundary) inside of $N(\alpha)$ to the identity map. If α is a curve, then since f is (well) tempered, it cannot twist about α , so f^n is also isotopic (rel boundary) to the identity map on $N(\alpha)$. But this implies f^n is isotopic to the identity on $X \cup Y$, and in particular, we can find an essential curve β in $X \cup Y$ crossing α which is f -periodic.

The proof that a component X has no self-adjacency along a periodic component $N(\alpha)$ is similar. In this case, we can isotope f^n to be identity on $X \cup N(\alpha)$, which contradicts that X and $N(\alpha)$ are components of the decomposition. $\square$

With all these results at hand, we can prove the wanted properties of the first return map of components of S_∞ .

Lemma 9.13. *For every component X of S_∞ , each boundary component is periodic. In particular, f returns to X , and there is a hyperbolic metric on X such that the first return f^n is isotopic to an isometric translation on X .*

Proof. Let α be a boundary component of X . The neighbor Y of X along α cannot be in S_∞ (by Lemma 9.6). So it is in S_{per} or S_0 . In both cases, the first return map to Y is periodic, and so α must be periodic. In particular, f must return to X . By Theorem 8.6, the first return f^n up to isotopy preserves a hyperbolic metric on X on which it acts as an isometric translation. $\square$

Our next goal is to show that there is no self-adjacent component of the decomposition. We need a preliminary result.

Lemma 9.14. *A component X of S_0 which is not a pair of pants can share at most one boundary with S_{per} . In this case, if α is the common boundary between X and S_{per} , then every non-contractible curve in X is homotopic to α .*

Proof. We now know X is f -periodic (by Lemma 9.9), so the proof is similar to Lemma 9.12. The assumptions on X are to rule out the existence of a periodic curve β crossing the common boundary of X and S_{per} essentially. $\square$

We can now show the no self-adjacency result.

Lemma 9.15. *No component of the canonical decomposition is self-adjacent.*

Proof. Suppose X is a self-adjacent component. By Corollary 9.11 and Lemma 9.12, X is a component of S_∞ . Since S_∞ has no compact boundary components by Lemma 9.8, there is a strip component Y of S_0 , with core line ℓ , so that X is the only neighbor of Y . Let ℓ_1 and ℓ_2 be the boundary components of Y . By

Lemma 9.13, there is some $n \geq 1$ so that f^n fixes ℓ_1 (and therefore ℓ_2 , since they are homotopic). Since Y is a strip, an orientation on ℓ induces orientations on ℓ_1 and ℓ_2 . Now pick a curve α contained in $X \cup Y$ and intersecting both ℓ_1 and ℓ_2 once. Then $\alpha \cap X$ is an arc from ∂X to ∂X , and by Proposition 8.3 it is wandering. This implies that f^n acts as a positive translation on ℓ_1 (with respect to the chosen orientation) if and only if it acts as a positive translation on ℓ_2 . But then we can homotope f^n on the strip, leaving it invariant on its boundary, so that it is a translation, and therefore α is wandering, a contradiction. $\square$

An easy consequence of the previous lemma is the fact that there are no components of the decomposition which are strips:

Corollary 9.16. *No component of S_0 , S_{per} or S_∞ is a strip.*

Proof. Since by construction every component of S_{per} and S_∞ contains an essential (possibly peripheral) curve, if there is a strip, it must be a component of S_0 . But by construction, a strip can arise only if a component of S_{per} or S_∞ is self-adjacent, which is impossible by Lemma 9.12 and 9.15. $\square$

We can now describe the possible topological possibilities for the components of S_0 :

Proof of Proposition 9.5. If X has negative Euler characteristic, we are in case (4) by Corollaries 7.7 and 9.11.

Suppose then that the Euler characteristic is non-negative. Note that X is not a sphere, being a subsurface of S . It also cannot be a disk or a once-punctured disk, by the construction of S_∞ and S_{per} .

If $\chi(X) = 1$, then X is a disk with a closed totally disconnected set with $n \in \{0, 1, \dots, \infty\}$ points removed from its boundary. If $n = 0$ or 1 , X would be a disk or a half-plane, which is impossible. Moreover, $n \neq 2$ by Corollary 9.16, so $n \geq 3$. By Lemma 9.14, X has at most one neighbor in S_{per} .

If $\chi(X) = 0$, then the interior of X is an open annulus. We already know that X cannot be the once-punctured disk, and it cannot be a closed annulus. Thus, X is either a once-punctured disk or a closed annulus with at least one point removed from its boundary. If it is an annulus with points removed from both boundaries, then it would contain an essential nonperipheral curve, contradicting Proposition 7.6. If X is a once-punctured disk with points removed from its boundary, then X has a non-contractible curve which is nonperipheral, so it cannot border S_{per} . If X is an annulus with points removed from one of its boundary, then the core curve of X is not homotopic to the non-compact boundaries of X , so X has at most one neighbor in S_{per} , and therefore by Lemma 9.8 it has exactly one neighbor in S_{per} , joined along the compact boundary component. $\square$

Since a well-tempered map returns to each component of S_0 , S_{per} and S_∞ , one might wonder if there is a uniform return time for all components. The answer is negative, as shown by the following lemma. Moreover, given a component of S_∞ ,

there is not always a uniform power of f fixing all of its boundary components at once.

Lemma 9.17. *There is a well-tempered map such that there is no uniform return time for all components of the canonical decomposition. Furthermore, the map can be chosen so that there is a component of S_∞ with no uniform return time for its boundary components.*

Proof. Consider the bordered surface

$$P = (\mathbb{R}_{\geq 0} \times \mathbb{R}) \setminus \bigcup_{n \geq 1} B_n,$$

where B_n is the open disk of center $(n, 0)$ and radius $\frac{1}{4}$. Let a_n be the oriented segment of the x -axis from ∂B_n to ∂B_{n+1} and a_0 the oriented segment of the x -axis from the origin to B_1 . Define the homomorphism $\psi : \pi_1(P) \rightarrow \mathbb{Z}$ by

$$\psi(\alpha) = \sum_{n \geq 0} 2^n \hat{\iota}(\alpha, a_n),$$

where $\hat{\iota}(\cdot, \cdot)$ denotes the algebraic intersection form.




FIGURE 22. The surface P with the arcs a_n in the proof of Lemma 9.17.

Let $p : \Sigma \rightarrow P$ be the associated cover of the kernel of ψ , and let f be a generator of the deck group. Then Σ is a bordered surface, whose boundary is a union of lines, given by lifts of the y -axis and of the curves ∂B_n . Note that, for any n , the number of lifts of ∂B_n is $2^n - 2^{n-1}$ and they are cyclically permuted by f . So there is no uniform power of f fixing all boundary components of Σ .

To turn this into an example of a well-tempered map on a borderless surface, we simply glue to each boundary component of Σ a copy of the thrice-punctured half-plane H . We get a surface S and a homeomorphism $\hat{f}$ of S so that $\hat{f}|_\Sigma = f$ and for every copy of H , the first return map is the identity.

The map $\bar{f}$ is well tempered and:

- $S_\infty = \Sigma$ and $\hat{f}|_{S_\infty} = f$, so there is no bound on the first return time of the boundary components of S_∞ ;

- S_{per} is the union (over all attached copies of H) of components homeomorphic to a disk with three punctures, spanned by the curves in each copy of H . The first return time of a component equals the first return time of the boundary component of the corresponding copy of H , so these return times are also unbounded;
- S_0 is the union (over all attached copies of H) of components homeomorphic to an annulus with a point removed from the boundary. Again there is no uniform bound on the first return times of components of S_0 . $\square$




FIGURE 23. The topological possibilities for the components of S_0 .

9.2. Proof of the classification theorem. Theorem 9.1 is now an easy consequence of the results proved so far.

Proof of Theorem 9.1. Let S_0 , S_{per} and S_∞ be given by Proposition 7.1. The hyperbolic structure on S is given by gluing the hyperbolic structures provided by Propositions 9.3, 9.4 and 9.5. Since there are no strips (Corollary 9.16), the resulting hyperbolic structure contains no funnels or half-planes. Moreover the same results guarantee that for every component of the decomposition the first return map of f exists and is isotopic to a periodic isometry or an isometric translation, as required. $\square$

We can also deduce Theorem A:

Proof of Theorem A. Let $\lambda = \partial S_\infty$. Since S_∞ is an f -invariant subsurface and has no compact boundary components (Proposition 9.3), λ is a one-manifold without compact components. Let X be a complementary component of λ . If X is a component of S_∞ , S_{per} or S_0 , we know by Theorem 9.1 that the first return map exists and is either a translation or a finite-order map.

Suppose then that X is the union of multiple components of the canonical decomposition, which are in S_{per} or S_0 by the definition of λ . Note that there needs to be at least one component of S_{per} , since by construction no two components of S_0 can be adjacent. In particular, a first return map exists.

If X contains a unique component Y of S_{per} , then X is of the form $Y \cup \bigcup_{i \in I} Z_i$, where the Z_i are components of S_0 , each of which is adjacent to Y . Consider $n > 0$ so that f^n is isotopic to the identity on Y . Then f^n also fixes at least one boundary component of each Z_i . By the classification of the topological possibilities of the components of S_0 (Proposition 9.5), together with the fact

that f is (well) tempered and thus there is no twisting, we get that f^n is also isotopic to the identity on every Z_i . In particular the first return map to X is periodic.

If X contains multiple components of S_{per} , they all must be annuli, otherwise we could construct a periodic curve not contained in S_{per} . By Proposition 9.5, X contains then a pair of pants component Y of S_0 , the annular components of S_{per} adjacent to it, together with crown components of S_0 , one per annular components of S_{per} in X . So topologically X is a pair of pants with at least one point removed from each boundary component. The three essential curves of X are periodic, so we can choose a power n of f under which they are all sent to themselves. As there is no twisting along these curves, f^n is the identity on X , and again the first return map to X is periodic. $\square$

9.3. Constructing well-tempered maps. Theorem A shows that every well-tempered map is obtained by gluing together translations and periodic maps supported on disjoint subsurfaces, not accumulating anywhere. If we want to construct well-tempered maps by following this procedure, we need to be careful when two supporting subsurfaces share a compact boundary component: we have to make sure that the gluing does not create a Dehn twist.

On the other hand, there is no need for special care when gluing along lines. More precisely:

Lemma 9.18. *Let $\{\Sigma_i\}_{i \in I}$ be a decomposition of a surface S into essential subsurfaces without self-adjacency, not accumulating anywhere and without compact boundary components. For every i , let f_i be either a periodic map or a translation of Σ_i , and assume that for every line $\ell \subset \partial\Sigma_i \cap \partial\Sigma_j$, $f_i(\ell) = f_j(\ell)$. Then there is a well-tempered mapping class f on S such that the restriction of f on each Σ_i is properly homotopic to f_i .*

Proof. Define S' to be the surface, homeomorphic to S , given by inserting strips at each line in the boundary of the decomposition. More precisely, we take the disjoint union of the Σ_i and of a strip $S_\ell = \ell \times [0, 1]$ for every ℓ in the boundary, and if $\ell \subset \partial\Sigma_i \cap \partial\Sigma_j$, we glue $\ell \times \{0\}$ with the copy of ℓ in Σ_i and $\ell \times \{1\}$ with the copy of ℓ in Σ_j . Define then f_ℓ on S_ℓ by linearly interpolating f_i and f_j :

$$f_\ell(p, t) = (1 - t)f_i(p) + tf_j(p)$$

where we have fixed an identification of ℓ with $\mathbb{R}$. Then we obtain a homeomorphism f of S' by gluing the f_i and the f_ℓ . We claim that f is well tempered.

Indeed, let α and β be two curves in S' . By the properties of the decomposition, and up to modifying the curves by a homotopy, there are only finitely many indices $j \in I$ and lines ℓ in the boundary of the decomposition such that $\alpha \cap \Sigma_j$, $\beta \cap \Sigma_j$, $\alpha \cap S_\ell$ or $\beta \cap S_\ell$ are nonempty. Since each f_j is well tempered, the quantities

$$i(f_j^n(\alpha \cap \Sigma_j), \beta \cap \Sigma_j)$$

are uniformly bounded. Moreover,

$$i(f_\ell^n(\alpha \cap S_\ell), \beta \cap S_\ell)$$

is bounded by the number of intersections of α and β with the boundary of the decomposition, hence it is also uniformly bounded. Therefore $i(f^n(\alpha), \beta)$ is uniformly bounded, so f is tempered. A similar argument shows the finiteness of the limit set of α . Hence f is well tempered. $\square$

Finally, note that if we glue periodic maps and translations and we obtain a well-tempered map, the canonical decomposition might not coincide with the collection of supporting subsurfaces of the maps we are gluing. For instance, let $S = \mathbb{R}^2 \setminus \mathbb{Z}^2$, and

$$\Sigma_1 = S \cap \left\{ y \geq \frac{3}{4} \right\} \text{ and } \Sigma_2 = S \cap \left\{ y \leq \frac{1}{4} \right\}.$$

Let f_1 be the map $(x, y) \mapsto (x + 1, y)$ restricted to Σ_1 , and let f_2 be the map $(x, y) \mapsto (x + 2, y)$ restricted to Σ_2 . We interpolate the maps f_1 and f_2 in the intermediate strip $\{1/4 \leq y \leq 3/4\}$. Then the induced map on S is a translation; in particular, $S = S_\infty$.

REFERENCES

- [1] *Travaux de Thurston sur les surfaces*, volume 66 of *Astérisque*. Société Mathématique de France, Paris, 1979. Séminaire Orsay, With an English summary.
- [2] Carolyn Abbott, Nicholas Miller, and Priyam Patel. Infinite-type loxodromic isometries of the relative arc graph. *Algebr. Geom. Topol.*, 25(1):563–644, 2025.
- [3] Santana Afton, Danny Calegari, Lvzhou Chen, and Rylee Alanza Lyman. Nielsen realization for infinite-type surfaces. *Proc. Amer. Math. Soc.*, 149(4):1791–1799, 2021.
- [4] Venancio Álvarez and José M. Rodríguez. Structure theorems for Riemann and topological surfaces. *J. London Math. Soc. (2)*, 69(1):153–168, 2004.
- [5] Javier Aramayona, Ariadna Fossas, and Hugo Parlier. Arc and curve graphs for infinite-type surfaces. *Proc. Amer. Math. Soc.*, 145(11):4995–5006, 2017.
- [6] Javier Aramayona and Nicholas G. Vlamis. Big mapping class groups: an overview. In *In the tradition of Thurston—geometry and topology*, pages 459–496. Springer, Cham, [2020] ©2020.
- [7] Assaf Bar-Natan and Yvon Verberne. The grand arc graph. *Math. Z.*, 305(2):21, 2023. Id/No 20.
- [8] Juliette Bavard. Hyperbolicité du graphe des rayons et quasi-morphismes sur un gros groupe modulaire. *Geom. Topol.*, 20(1):491–535, 2016.
- [9] Juliette Bavard, Spencer Dowdall, and Kasra Rafi. Isomorphisms between big mapping class groups. *Int. Math. Res. Not. IMRN*, (10):3084–3099, 2020.
- [10] Lipman Bers. An extremal problem for quasiconformal mappings and a theorem by Thurston. *Acta Math.*, 141(1-2):73–98, 1978.
- [11] Mladen Bestvina and Michael Handel. Train-tracks for surface homeomorphisms. *Topology*, 34(1):109–140, 1995.
- [12] L. E. J. Brouwer. Beweis des ebenen Translationssatzes. *Math. Ann.*, 72(1):37–54, 1912.
- [13] John Cantwell, Lawrence Conlon, and Sergio R. Fenley. Endperiodic automorphisms of surfaces and foliations. *Ergodic Theory Dynam. Systems*, 41(1):66–212, 2021.
- [14] Andrew J. Casson and Steven A. Bleiler. *Automorphisms of surfaces after Nielsen and Thurston*, volume 9 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 1988.
- [15] Peter Feller, Diana Hubbard, and Hannah Turner. The Dehn twist coefficient for big and small mapping class groups. *J. Lond. Math. Soc., II. Ser.*, 112(2):29, 2025. Id/No e70251.

- [16] Michael Handel and William P. Thurston. New proofs of some results of Nielsen. *Adv. in Math.*, 56(2):173–191, 1985.
- [17] Jesús Hernández Hernández, Israel Morales, and Ferrán Valdez. The Alexander method for infinite-type surfaces. *Michigan Math. J.*, 68(4):743–753, 2019.
- [18] Nikolai V. Ivanov. Automorphism of complexes of curves and of Teichmüller spaces. *Internat. Math. Res. Notices*, (14):651–666, 1997.
- [19] Anatole Katok and Boris Hasselblatt. *Introduction to the modern theory of dynamical systems*, volume 54 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1995. With a supplementary chapter by Katok and Leonardo Mendoza.
- [20] Mustafa Korkmaz. Automorphisms of complexes of curves on punctured spheres and on punctured tori. *Topology Appl.*, 95(2):85–111, 1999.
- [21] Feng Luo. Automorphisms of the complex of curves. *Topology*, 39(2):283–298, 2000.
- [22] Richard T. Miller. Geodesic laminations from Nielsen’s viewpoint. *Adv. in Math.*, 45(2):189–212, 1982.
- [23] Jakob Nielsen. Untersuchungen zur Topologie der geschlossenen zweiseitigen Flächen. *Acta Math.*, 50(1):189–358, 1927.
- [24] Jakob Nielsen. Untersuchungen zur Topologie der geschlossenen zweiseitigen Flächen. II. *Acta Math.*, 53(1):1–76, 1929.
- [25] Jakob Nielsen. Untersuchungen zur Topologie der geschlossenen zweiseitigen Flächen. III. *Acta Math.*, 58(1):87–167, 1932.
- [26] Jakob Nielsen. Abbildungsklassen endlicher Ordnung. *Acta Math.*, 75:23–115, 1942.
- [27] Martin Roller. Poc Sets, Median Algebras and Group Actions. *arXiv e-prints*, page arXiv:1607.07747, July 2016.
- [28] William P. Thurston. On the geometry and dynamics of diffeomorphisms of surfaces. *Bull. Amer. Math. Soc. (N.S.)*, 19(2):417–431, 1988.
- [29] Dragomir Šarić. Train tracks and measured laminations on infinite surfaces. *Trans. Amer. Math. Soc.*, 374(12):8903–8947, 2021.

DEPT OF MATH, UNIV OF UTAH, SALT LAKE CITY, UT 84112, USA

Email address: `bestvina@math.utah.edu`

CNRS, UNIV PARIS EST CRETEIL, UNIV GUSTAVE EIFFEL, LAMA UMR8050, F-94010 CRETEIL, FRANCE

Email address: `federica.fanoni@u-pec.fr`

DEPT OF MATH, UNIV OF OKLAHOMA, NORMAN, OK 73019, USA

Email address: `jing@ou.edu`