

Fys3120 Oblig 11

Joseph Knutson
josephkn

May 1, 2017

1a

- Problem 1A non-relativistic particle, with electric charge q and mass m moves in a magnetic dipole field, given by the vector potential

$$\vec{A} = \frac{\mu_0}{4\pi r^3} (\vec{\mu} \times \vec{r}) \quad (1)$$

Show that the Lagrangian is

$$L = \frac{1}{2} m \vec{v}^2 + \frac{q\mu_0}{4\pi m r^3} \vec{\mu} \cdot \vec{l}, \quad (2)$$

Where $\vec{l} = m\vec{r} \times \vec{v}$

We begin with defining the Lagrangian:

$$L = T - V$$

From equation 2 it is obvious that $T = 1/2 m v^2$, which means, we need to prove that the potential is:

$$V = \frac{q\mu_0}{4\pi m r^3} \vec{\mu} \cdot \vec{l}$$

In order to prove this, we use the fact that $-q\vec{v} \cdot \vec{A} = V$:

$$\begin{aligned} V &= q\vec{v} \cdot \vec{A} \\ &= \frac{q\mu_0}{4\pi r^3} \vec{v} \cdot (\vec{\mu} \times \vec{r}) \\ &= \frac{q\mu_0}{4\pi m r^3} \vec{p} \cdot (\vec{\mu} \times \vec{r}) \\ &= -\frac{q\mu_0}{4\pi m r^3} \vec{\mu} \cdot (\vec{r} \times \vec{p}) \\ &= -\frac{q\mu_0}{4\pi m r^3} \vec{\mu} \cdot \vec{l} \end{aligned} \quad (3)$$

1b

- Particle moves in the xy-plane and ϕ is the angle between the x-axis and the particle's position vector \vec{r} . The magnetic dipole momentum, $\vec{\mu}$, has a direction along the z-axis. Express the Lagrangian as :

$$L = \frac{1}{2}m(\dot{r}^2 + r^2\dot{\phi}^2) + \lambda \frac{\dot{\phi}}{r} \quad (4)$$

where

$$\lambda = q\mu_0|\vec{\mu}|/4\pi \quad (5)$$

To express the Lagrangian in equation 4, we start by expressing the kinetic energy in polar coordinates:

$$x = r \cos \phi$$

$$y = r \sin \phi$$

$$z = 0$$

Finding velocity to express T:

$$\vec{v} = (\dot{x}, \dot{y}, 0)$$

$$\dot{x} = \dot{r} \cos \phi - r\dot{\phi} \sin \phi$$

$$\dot{y} = \dot{r} \sin \phi + r\dot{\phi} \cos \phi$$

$$\dot{z} = 0$$

$$\Rightarrow |\vec{v}| = \sqrt{\dot{r}^2 + r^2\dot{\phi}^2}$$

$$\Rightarrow T = \frac{1}{2}m(\dot{r}^2 + r^2\dot{\phi}^2)$$

We now have the kinetic term of equation 4, but we have yet to express the potential which should be: $V = \dot{\phi}\lambda/r$. To derive this, we need to substitute λ from equation 5 into the potential term of our Lagrangian found in both eq. 2 and 3.

First trick is to express the dipole momentum vector, using equation 5:

$$\vec{\mu} = |\vec{\mu}|\vec{k} \quad (6)$$

Before substituting λ into the potential, we split the magnetic dipole vector as we did in 6:

$$\begin{aligned} -V &= \frac{q\mu_0}{4\pi mr^3} \vec{\mu} \cdot \vec{l} \\ &= \frac{q\mu_0}{4\pi mr^3} |\vec{\mu}|\vec{k} \cdot \vec{l} \\ &= \frac{\lambda}{mr^3} \vec{k} \cdot \vec{l} \\ &= |\vec{l}| \frac{\lambda}{mr^3} = |\vec{r} \times \vec{v}| \frac{m\lambda}{mr^3} = |\vec{r} \times \vec{v}| \frac{\lambda}{r^3} \end{aligned} \quad (7)$$

This crossproduct requires us to look back when we derived the kinetic term:

$$\begin{aligned}\vec{r} &= (x, y, z) \\ &= (r \cos \phi, r \sin \phi, 0)\end{aligned}\tag{8}$$

$$\begin{aligned}\vec{v} &= (\dot{x}, \dot{y}, 0) \\ &= (\dot{r} \cos \phi - r \dot{\phi} \sin \phi, \dot{r} \sin \phi + r \dot{\phi} \cos \phi, 0)\end{aligned}\tag{9}$$

$$\Rightarrow \tag{10}$$

$$\vec{r} \times \vec{v} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ x & y & 0 \\ \dot{x} & \dot{y} & 0 \end{vmatrix} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ x & y & 0 \\ \dot{x} & \dot{y} & 0 \end{vmatrix}\tag{11}$$

$$\begin{aligned}&= (x\dot{y} - y\dot{x})\vec{k} \\ &= r \cos \phi (\dot{r} \sin \phi + r \dot{\phi} \cos \phi) - r \sin \phi (\dot{r} \cos \phi - r \dot{\phi} \sin \phi)\end{aligned}\tag{12}$$

$$= r^2 \dot{\phi} \cos^2 \phi + r^2 \dot{\phi} \sin^2 \phi\tag{13}$$

$$= r^2 \dot{\phi}\tag{14}$$

Now that we have the crossproduct, we can insert it into our potential, back in equation 7

$$\begin{aligned}-V &= |\vec{r} \times \vec{v}| \frac{\lambda}{r^3} \\ &= r^2 \dot{\phi} \frac{\lambda}{r^3} \\ &= \dot{\phi} \frac{\lambda}{r}\end{aligned}$$

If we look back at our kinetic term, we saw that $T = \frac{1}{2}m(\dot{r}^2 + r^2\dot{\phi}^2)$, with this we can now express the Lagrangian on the form we set out to:

$$L = T - V = \frac{1}{2}m(\dot{r}^2 + r^2\dot{\phi}^2) + \lambda \frac{\dot{\phi}}{r}\tag{15}$$

1c

- Write Lagrange's equation for the coordinate r , expressed in terms of \dot{r} , \ddot{r} and p_ϕ , and use the equation to find \ddot{r} as a function of r and p_ϕ . Compare the expression with that of the particle's kinetic energy.

From the Lagrangian in equation 15, we derive Lagrange's equation for ϕ . Notice that the second term of the Lagrange equation is zero, which gives us the constant conjugate momentum:

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{\phi}} \right) - \frac{\partial L}{\partial \phi} = 0$$

$$\frac{\partial L}{\partial \phi} = 0 \quad (16)$$

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{\phi}} \right) = \frac{d}{dt} \left(mr^2 \dot{\phi} + \frac{\lambda}{r} \right) \quad (17)$$

$$\Rightarrow p_\phi = mr^2 \dot{\phi} + \frac{\lambda}{r} \quad (18)$$

$$\Rightarrow \dot{\phi} = \frac{p_\phi r - \lambda}{mr^3} \quad (19)$$

We can use equation 19 to substitute away $\dot{\phi}$ out of Lagrange's equation of r :

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{r}} \right) - \frac{\partial L}{\partial r} = 0$$

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{r}} \right) = \frac{d}{dt} m \dot{r} = m \ddot{r} \quad (20)$$

$$\begin{aligned} \frac{\partial L}{\partial r} &= m r \dot{\phi}^2 - \lambda \frac{\dot{\phi}}{r^2} \quad \text{let's substitute } \dot{\phi} \\ &= m r \left(\frac{p_\phi r - \lambda}{mr^3} \right)^2 - \frac{p_\phi r \lambda - \lambda^2}{mr^5} \\ &= m \left(\frac{p_\phi^2 r^2 - 2p_\phi r \lambda + \lambda^2 - (p_\phi r \lambda - \lambda^2)}{m^2 r^5} \right) \\ &= m \left(\frac{p_\phi^2 r^2 - 3p_\phi r \lambda}{m^2 r^5} \right) \end{aligned}$$

Now let's set up Lagrange's equation since we have both derivative terms

figured out:

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{r}} \right) - \frac{\partial L}{\partial r} = m \left(\ddot{r} - \frac{p_\phi^2 r^2 + 3p_\phi r \lambda}{m^2 r^5} \right) \quad (21)$$

$$\implies \ddot{r} = \frac{p_\phi^2 r^2 - 3p_\phi r \lambda}{m^2 r^5} \quad (22)$$

$$\implies \dot{r} = \frac{p_\phi^2}{m^2} \int \frac{1}{r^3} dt - 3\lambda \frac{p_\phi}{m^2} \int \frac{1}{r^4} dt \quad (23)$$

Since our goal is to express \dot{r}^2 we need to integrate \ddot{r} , but first we gotta substitute dt out of the equation.

$$dt = \frac{dr}{\dot{r}} dt = \frac{dt}{\dot{r}} dr = \frac{1}{\dot{r}} dr$$

Let's substitute this into equation 23 to solve it:

$$\begin{aligned} \dot{r} &= \frac{p_\phi^2}{m^2} \int \frac{1}{r^3} dt - 3\lambda \frac{p_\phi}{m^2} \int \frac{1}{r^4} dt \\ \dot{r} &= \frac{p_\phi^2}{m^2} \int \frac{1}{\dot{r} r^3} dr - 3\lambda \frac{p_\phi}{m^2} \int \frac{1}{\dot{r} r^4} dr \end{aligned} \quad (24)$$

Let's do a cool trick and multiply with \dot{r} on both sides of equation 24:

$$\begin{aligned} \dot{r}^2 &= \frac{p_\phi^2}{m^2} \int \frac{1}{r^3} dr - 3\lambda \frac{p_\phi}{m^2} \int \frac{1}{r^4} dr \\ &= -\frac{p_\phi^2}{2m^2} \frac{1}{r^2} + \lambda \frac{p_\phi}{m^2} \frac{1}{r^3} \end{aligned} \quad (25)$$

$$= \frac{p_\phi}{2m^2 r^2} \left(\frac{2\lambda}{r} - p_\phi \right) \quad (26)$$

Comparing this mess with the kinetic energy is completely lost on me. Perhaps I made a miscalculation.

$$T = \frac{1}{2} m (\dot{r}^2 + r^2 \dot{\phi}^2) \quad (27)$$




Figure 1: Current loop ABCD of length a and width b . Zero charge density.

2 Relatively cool circuit

Figure 1 shows a rectangular current loop ABCD. In the loop's rest frame, S , the loop has length a in the x direction and width b in the y direction, the current is I and the charge density is zero. We remind you of the following general definitions of the electric dipole moment \vec{p} , and the magnetic dipole moment \vec{m} , for a given current distribution:

$$\vec{p} = \int \vec{r} \rho(\vec{r}) d^3r \quad (28)$$

$$\vec{m} = \frac{1}{2} \int (\vec{r} \times \vec{j}(\vec{r})) d^3r \quad (29)$$

2a

- Show that the electric dipole moment is zero in the loop's restframe. Show that the magnetic dipole moment is $\vec{m} = I \vec{a} \times \vec{b}$ where $I = j\Delta$. Δ is the cross sectional area of the conductor.

Because our conductor is without charge density in the S frame ($\rho = 0$), the integral in equation 28, the electric dipole momentum, equals zero.

In order to determine the magnetic dipole momentum we must express $\vec{r} \times \vec{j}$