

Path Integral Quantum Field Theory

Mathias Driesse

December 22, 2022

1 Gaussian Integrals

The general Gaussian integral, for a complex, symmetric $n \times n$ matrix A such that $\operatorname{Re} A \geq 0$ and the eigenvalues a_i of A are nonzero, is given by

$$Z_A(b) = \int d^n x \exp \left(-\frac{1}{2} \sum_{i,j=1}^n x_i A_{ij} x_j + \sum_{i=1}^n b_i x_i \right) = (2\pi)^{n/2} (\det A)^{-1/2} \exp \left(\frac{1}{2} \sum_{i,j=1}^n b_i A_{ij}^{-1} b_j \right).$$

(1.1)

Let μ be a measure in \mathbb{R}^n ; we define the expectation value

$$\begin{aligned} \langle F \rangle_\mu &= \int d\mu(x) F(x) \\ &= \int d^n x \Omega(x) F(x). \end{aligned}$$

(1.2)

The measure is normalized so that

$$\int d\mu(x) = 1.$$

(1.3)

We define the generating function

$$Z_\mu(b) = \left\langle e^{(b,x)} \right\rangle_\mu = \int d\mu(x) \exp \left(\sum_{i=1}^n b_i x_i \right),$$

(1.4)

which is a function of the n -dimensional vector b and the measure μ . The integrand can be expanded

$$\exp \left(\sum_{i=1}^n b_i x_i \right) = \sum_{\ell=0}^{\infty} \frac{1}{\ell!} \sum_{i_1 \dots i_\ell=1}^n b_{i_1} \dots b_{i_\ell} x_{i_1} \dots x_{i_\ell}.$$

(1.5)

Therefore, substituting the definition of the correlator

$$\langle x_{i_1} \dots x_{i_\ell} \rangle_\mu = \int d\mu(x) x_{i_1} \dots x_{i_\ell}$$

(1.6)

we obtain

$$Z_\mu(b) = \sum_{\ell=0}^{\infty} \frac{1}{\ell!} \sum_{i_1 \dots i_\ell=1}^n b_{i_1} \dots b_{i_\ell} \langle x_{i_1} \dots x_{i_\ell} \rangle_\mu.$$

(1.7)

Furthermore, it is useful to notice that

$$\frac{\partial}{\partial b_k} Z_\mu(b) = \int d\mu(x) x_k \exp \left(\sum_{i=1}^n b_i x_i \right),$$

(1.8)

which allows the correlators to be written as

$$\langle x_{i_1} \dots x_{i_\ell} \rangle_\mu = \frac{\partial}{\partial b_{i_1}} \dots \frac{\partial}{\partial b_{i_\ell}} Z_\mu(b) \Big|_{b=0}. \quad (1.9)$$

Let us consider the Gaussian measure

$$d\mu_0(x) = d^n x \Omega_0(x) = d^n x \mathcal{N}_0 \exp \left(-\frac{1}{2} \sum_{i,j=0}^n x_i A_{ij} x_j \right), \quad (1.10)$$

where the normalization \mathcal{N}_0 is fixed by the normalization of the measure:

$$\mathcal{N}_0 = (2\pi)^{-n/2} (\det A)^{1/2}. \quad (1.11)$$

The generating function in this case can be readily computed using the boxed equation above:

$$Z(b) = \frac{Z_A(b)}{Z_A(0)} = \exp \left(\frac{1}{2} \sum_{i,j=1}^n b_i A_{ij}^{-1} b_j \right). \quad (1.12)$$

In the future, we will drop any subscript A when referring to a Gaussian measure for simplicity. And therefore, defining $\Delta_{ij} = A_{ij}^{-1}$,

$$\langle x_{i_1} \dots x_{i_\ell} \rangle_0 = \frac{\partial}{\partial b_{i_1}} \dots \frac{\partial}{\partial b_{i_\ell}} \exp \left(\frac{1}{2} \sum_{i,j=1}^n b_i \Delta_{ij} b_j \right) \Big|_{b=0}.$$

(1.13)

1.1 Wick's Theorem

Let us start with a couple explicit examples, which we will compute in full detail in order to get familiar with the algebraic manipulations.

One-point function: Let k be an integer between 1 and n , we have

$$\begin{aligned} \langle x_k \rangle_0 &= \frac{\partial}{\partial b_k} \exp \left(\frac{1}{2} \sum_{i,j=1}^n b_i \Delta_{ij} b_j \right) \Big|_{b=0} \\ &= \left(\frac{1}{2} \sum_{j=1}^n \Delta_{kj} b_j + \frac{1}{2} \sum_{i=1}^n b_i \Delta_{ik} \right) \exp \left(\frac{1}{2} \sum_{i,j=1}^n b_i \Delta_{ij} b_j \right) \Big|_{b=0} \\ &= \left(\sum_{j=1}^n \Delta_{kj} b_j \right) \exp \left(\frac{1}{2} \sum_{i,j=1}^n b_i \Delta_{ij} b_j \right) \Big|_{b=0} \\ &= 0, \end{aligned} \quad (1.14)$$

where the last equality comes from the fact that the expression is linear in b , and we need to set $b = 0$.

Two-point function: We now consider a pair of indices k, l , and compute

$$\begin{aligned} \langle x_k x_l \rangle_0 &= \frac{\partial}{\partial b_l} \frac{\partial}{\partial b_k} \exp \left(\frac{1}{2} \sum_{i,j=1}^n b_i \Delta_{ij} b_j \right) \Big|_{b=0} \\ &= \left[\Delta_{kl} + \left(\sum_{j=1}^n \Delta_{kj} b_j \right) \left(\sum_{m=1}^n \Delta_{lm} b_m \right) \right] \exp \left(\frac{1}{2} \sum_{i,j=1}^n b_i \Delta_{ij} b_j \right) \Big|_{b=0} \\ &= \Delta_{kl} \end{aligned} \quad (1.15)$$

Thus, the generating function $Z(b)$ provides a systematic way to compute all correlators for a multi-dimensional Gaussian distribution. Having understood the general rule for such a process, we can generate a recipe to compute $\langle x_{i_1} \dots x_{i_\ell} \rangle_0$, known as *Wick's Theorem*:

- Write down each $x_{i_1} \dots x_{i_\ell}$ and organize them pairwise (i_p, i_q) . Note that if ℓ must be even for the correlator to be non-zero.
- There are $(\ell - 1) \times (\ell - 3) \times \dots \times 3 \times 1$ ways of doing this. Sum over all of these possible pairings.
- To each pair (i_p, i_q) associate a factor $\Delta_{i_p i_q}$.

Let us revisit our result for the **two-point function**: for $\langle x_{i_1} x_{i_2} \rangle_0$, there is only one possible pairing (i_1, i_2) . Therefore

$$\langle x_{i_1} x_{i_2} \rangle_0 = \Delta_{i_1 i_2}. \quad (1.16)$$

Four-point function: For $\langle x_{i_1} x_{i_2} x_{i_3} x_{i_4} \rangle_0$, there are three different pairings

$$P = \{ \{ (i_1, i_2), (i_3, i_4) \}, \{ (i_1, i_3), (i_2, i_4) \}, \{ (i_1, i_4), (i_2, i_3) \} \} \quad (1.17)$$

Wick's theorem then yields

$$\langle x_{i_1} x_{i_2} x_{i_3} x_{i_4} \rangle_0 = \Delta_{i_1 i_2} \Delta_{i_3 i_4} + \Delta_{i_1 i_3} \Delta_{i_2 i_4} + \Delta_{i_1 i_4} \Delta_{i_2 i_3}. \quad (1.18)$$

We can also represent these pairings, *Wick contractions*, as should be familiar from canonical quantum field theory:

$$\langle x_i x_j \rangle_0 = \Delta_{ij} = \overline{x_i x_j} \quad (1.19)$$

1.2 Perturbed Gaussian Measure

Let us now consider a more complicated measure,

$$\Omega(x) = \frac{1}{Z(\lambda)} e^{-S(x, \lambda)}, \quad (1.20)$$

where the normalization is given as usual by

$$Z(\lambda) = \int d^n x e^{-S(x, \lambda)} \quad (1.21)$$

and

$$\begin{aligned} S(x, \lambda) &= \frac{1}{2} \sum_{i,j=1}^n x_i A_{ij} x_j + \lambda V(x) \\ &= S_0(x) + \lambda V(x). \end{aligned} \quad (1.22)$$

We call $V(\lambda)$ the potential term, foreshadowing the physics we will be doing using the perturbed Gaussian measure. Furthermore

$$\begin{aligned} Z(\lambda) &= \sum_{k=0}^{\infty} \frac{(-\lambda)^k}{k!} \int d^n x V^k(x) \exp \left(-\frac{1}{2} \sum_{i,j=1}^n x_i A_{ij} x_j \right) \\ &= Z(0) \sum_{k=0}^{\infty} \frac{(-\lambda)^k}{k!} \langle V^k(x) \rangle_0, \end{aligned} \quad (1.23)$$

where in the first line we have expanded the exponential term containing $e^V(x)$ and in the second line used

$$\langle F(x) \rangle_0 = \frac{1}{Z(0)} \int d^n x \exp \left(-\frac{1}{2} \sum_{i,j=1}^n x_i A_{ij} x_j \right) F(x), \quad (1.24)$$

and $1/Z(0)$ referring to the normalization constant \mathcal{N}_0 from Eq. 1.10. Recalling Eq. 1.13, given that $F(x)$ admits a Taylor expansion

$$\begin{aligned} F(x) &= \sum_{\ell=0}^{\infty} \sum_{i_1 \dots i_{\ell}=1}^n F_{i_1 \dots i_{\ell}} x_{i_1} \dots x_{i_{\ell}} \\ &= F_0 + F_1 x_1 + \dots + F_n x_n \\ &\quad + F_{11} x_1^2 + F_{12} x_1 x_2 + \dots + F_{nn} x_n^2 \\ &\quad + \dots \end{aligned} \tag{1.25}$$

where $F_{i_1 \dots i_{\ell}}$ refer to the constant expansion coefficients, we can define the expectation value of F for a generic measure μ as

$$\langle F(x) \rangle_{\mu} = \sum_{\ell=0}^{\infty} \sum_{i_1 \dots i_{\ell}=1}^n F_{i_1 \dots i_{\ell}} \langle x_{i_1} \dots x_{i_{\ell}} \rangle_{\mu}; \tag{1.26}$$

for the case of a Gaussian measure we obtain

$$\begin{aligned} \langle F(x) \rangle_0 &= \sum_{\ell=0}^{\infty} \sum_{i_1 \dots i_{\ell}=1}^n F_{i_1 \dots i_{\ell}} \frac{\partial}{\partial b_{i_1}} \dots \frac{\partial}{\partial b_{i_{\ell}}} \exp \left(\frac{1}{2} \sum_{i,j=1}^n b_i \Delta_{ij} b_j \right) \Big|_{b=0} \\ &= F \left(\frac{\partial}{\partial b} \right) \exp \left(\frac{1}{2} \sum_{i,j=1}^n b_i \Delta_{ij} b_j \right) \Big|_{b=0}. \end{aligned} \tag{1.27}$$

Thus, returning to our potential term,

$$\begin{aligned} \frac{Z(\lambda)}{Z(0)} &= \left\langle e^{-\lambda V(x)} \right\rangle_0 \\ &= \exp \left[-\lambda V \left(\frac{\partial}{\partial b} \right) \right] \exp \left(\frac{1}{2} \sum_{i,j=1}^n b_i \Delta_{ij} b_j \right) \Big|_{b=0}. \end{aligned} \tag{1.28}$$

1.3 Perturbed Gaussian Correlators

The perturbative treatment discussed above can be extended to compute moments of the distribution:

$$\langle x_{i_1} \dots x_{i_{\ell}} \rangle = \frac{\int d^n x e^{-S(x,\lambda)} x_{i_1} \dots x_{i_{\ell}}}{\int d^n x e^{-S(x,\lambda)}} = \frac{1}{Z(\lambda)} \int d^n x e^{-S(x,\lambda)} x_{i_1} \dots x_{i_{\ell}} \tag{1.29}$$

which is often referred to as an ℓ -point correlator/function. Let us start our discussion with a simple example.

Two-point function: we need to evaluate

$$\begin{aligned} \int d^n x e^{-S(x,\lambda)} x_{i_1} x_{i_2} &= \int d^n x e^{-S_0(x,\lambda)} \left[\sum_{k=0}^n \frac{(-\lambda)^k}{k!} \lambda^k V^k(x) \right] x_{i_1} x_{i_2} \\ &= Z(0) \sum_{k=0}^n \frac{(-\lambda)^k}{k!} \langle V^k(x) x_{i_1} x_{i_2} \rangle, \end{aligned} \tag{1.30}$$

where in the second line we have expressed the initial correlator in terms of correlators computed in the Gaussian theory, denoted by $\langle \dots \rangle_0$. The Gaussian correlators can be computed using Wick's theorem as before. We shall consider again a quartic potential

$$V(x) = \frac{1}{4!} \sum_{i=1}^n x_i^4, \tag{1.31}$$

and compute all the terms in Eq. 1.30 order by order in λ up to order λ^2 .

- $\mathcal{O}(\lambda^0)$: For $k = 0$ we simply get the two-point Gaussian correlator

$$\langle x_{i_1} x_{i_2} \rangle = \Delta_{i_1 i_2}. \quad (1.32)$$

- $\mathcal{O}(\lambda^1)$: At first order in λ we have one insertion of V :

$$\langle V(x) x_{i_1} x_{i_2} \rangle_0 = \frac{1}{4!} \sum_{i=1}^n \langle x_i^4 x_{i_1} x_{i_2} \rangle_0. \quad (1.33)$$

This Gaussian expectation value involving six factors of x can be evaluated using Wick's theorem. There are two types of contractions.

1. x_1 is contracted with x_2 , and the four x_i are contracted amongst themselves:

$$\overbrace{x_{i_1} x_{i_2}}^{\text{contracted}} (\overbrace{x_i x_i x_i x_i}^{\text{contracted}} + \overbrace{x_i x_i x_i x_i}^{\text{contracted}} + \overbrace{x_i x_i x_i x_i}^{\text{contracted}}) = \Delta_{i_1 i_2} \langle x_i^4 \rangle_0 \quad (1.34)$$

2. x_1 and x_2 are contracted with some of the x_i ; there are 12 such contractions:

$$\overbrace{x_{i_1} x_{i_2} x_i x_i x_i x_i}^{\text{contracted}} + \dots = \Delta_{i i_1} \Delta_{i i_2} \Delta_{i i} \times 4 \times 3. \quad (1.35)$$

Collecting all the terms yields

$$\frac{1}{4!} \sum_{i=1}^n \langle x_i^4 x_{i_1} x_{i_2} \rangle_0 = \Delta_{i_1 i_2} \frac{1}{4!} \sum_{i=1}^n \langle x_i^4 \rangle_0 + \frac{1}{4!} \times 4 \times 3 \sum_{i=1}^n \Delta_{i i_1} \Delta_{i i_2} \Delta_{i i}. \quad (1.36)$$

- $\mathcal{O}(\lambda^2)$: At this order we need to evaluate

$$\frac{1}{2!} \langle V(x)^2 x_{i_1} x_{i_2} \rangle_0 = \frac{1}{2!} \frac{1}{(4!)^2} \sum_{i,j=1}^n \langle x_i^4 x_j^4 x_{i_1} x_{i_2} \rangle_0. \quad (1.37)$$

There are five different types of contractions, each of them coming with a given multiplicity. We encourage the interested reader to compute those contributions, and check carefully that the correct multiplicities are recovered. Collecting all contributions yields

$$\begin{aligned} \int d^n x e^{-S(x,\lambda)} x_{i_1} x_{i_2} = Z(0) & \left[\Delta_{i_1 i_2} - \lambda \left(\Delta_{i_1 i_2} \frac{1}{4!} \sum_{i=1}^n \langle x_i^4 \rangle_0 + \frac{1}{2} \sum_{i=1}^n \Delta_{i i_1} \Delta_{i i_2} \Delta_{i i} \right) \right. \\ & + \lambda^2 \left(\frac{1}{2!} \Delta_{i_1 i_2} \frac{1}{(4!)^2} \sum_{i,j=1}^n \langle x_i^4 x_j^4 \rangle_0 + \frac{1}{2!} \sum_{i=1}^n \Delta_{i i_1} \Delta_{i i_2} \Delta_{i i} \frac{1}{4!} \sum_{j=0}^n \langle x_j^4 \rangle_0 \right. \\ & + \frac{1}{4} \sum_{i,j=1}^n \Delta_{i i_1} \Delta_{i i_2} \Delta_{i j}^2 \Delta_{j j} + \frac{1}{6} \sum_{i,j=1}^n \Delta_{i i_1} \Delta_{j i_2} \Delta_{i j}^3 \\ & \left. \left. + \frac{1}{4} \sum_{i,j=1}^n \Delta_{i i_1} \Delta_{j i_2} \Delta_{i j} \Delta_{i i} \Delta_{j j} \right) \right]. \end{aligned} \quad (1.38)$$

A term that factorises as the product of a subdiagram with external lines and a subdiagram that is made of loops only is called a *vacuum contribution*. For instance, $\Delta_{i_1 i_2} \frac{1}{4!} \sum_{i=1}^n \langle x_i^4 \rangle_0$ is a vacuum contribution because $\langle x_i^4 \rangle_0$ is one.

Finally, we need to divide this expression by $Z(\lambda)$ in order to obtain the two-point correlator as defined in Eq. 1.29. As a result, we obtain a factor $Z(0)/Z(\lambda)$ multiplying the expression inside the square bracket in Eq. 1.38. The ratio $Z(0)/Z(\lambda)$ may be computed using Eq. 1.28 and expanding as $1/(1-x) = 1 + x + x^2 + \dots$ and sorting the terms by powers of λ to obtain

$$Z(0)/Z(\lambda) = 1 + \frac{1}{4!} \sum_{i=1}^n \langle x_i^4 \rangle_0 - \frac{1}{2!} \frac{1}{(4!)^2} \lambda^2 \sum_{i,j=1}^n \langle x_i^4 x_j^4 \rangle_0 + \frac{1}{(4!)^2} \lambda^2 \left(\sum_{i=1}^n \langle x_i^4 \rangle_0 \right)^2 + \mathcal{O}(\lambda^3) \quad (1.39)$$

This cancels all vacuum contributions.

1.4 Generating Functions for the Perturbed Gaussian Measure

We can now generalise the idea of a generating function to the case of a non-Gaussian measure. Introducing

$$Z(b, \lambda) = \int d^n x \exp[-S(x, \lambda) + b_i x_i], \quad (1.40)$$

and remembering that we can also write

$$\langle e^{b_i x_i} \rangle = Z(b, \lambda) / Z(\lambda), \quad (1.41)$$

the correlators in the perturbed measure are obtained by differentiation

$$\langle x_{i_1} \dots x_{i_\ell} \rangle = \frac{1}{Z(\lambda)} \frac{\partial}{\partial b_{i_1}} \dots \frac{\partial}{\partial b_{i_\ell}} Z(b, \lambda) \Big|_{b=0}. \quad (1.42)$$

The logarithm of $Z(b, \lambda)$ is usually denoted $W(b, \lambda)$,

$$Z(b, \lambda) = e^{W(b, \lambda)}; \quad (1.43)$$

$W(b, \lambda)$ is the generator of the connected ℓ -point correlators $W_{i_1 \dots i_\ell}$, i.e. the correlators that can be represented as a single diagram, with ℓ open ends:

$$W_{i_1 \dots i_\ell} = \frac{\partial}{\partial b_{i_1}} \dots \frac{\partial}{\partial b_{i_\ell}} W(b, \lambda) \Big|_{b=0}. \quad (1.44)$$

In statistics, the $W_{i_1 \dots i_\ell}$ are called the *cumulants* of the probability distribution $e^{-S(x, \lambda)}$.

2 Path Integral Quantum Mechanics

In this section, we will discuss the essentials of the path integral formulation of quantum mechanics, which will allow us to connect Gaussian integrals and the calculation of correlators to observables in quantum field theory. We consider a point particle in one dimension, with position operator \hat{Q} such that

$$\hat{Q} |q\rangle = q |q\rangle \quad (2.1)$$

with a completeness condition

$$\int dq |q\rangle \langle q| = 1. \quad (2.2)$$

The aim of this section is to find an expression for the quantum amplitude $\langle q' t' | q t \rangle = \langle q' | e^{-i\hat{H}(t'-t)} | q \rangle$ for a Hamiltonian \hat{H} .

2.1 Setting up the Path Integral

In order to proceed with the calculation, let us define $T = t' - t$ to be the size of the time interval, and $\epsilon = T/n$, where n is an integer. Then

$$\begin{aligned} t_0 &= t \\ t_k &= t_0 + k\epsilon, k = 1, \dots, n-1 \\ t_n &= t + n\epsilon = t' \end{aligned} \quad (2.3)$$

and

$$\begin{aligned} \langle q' | e^{-i\hat{H}(t'-t)} | q \rangle &= \langle q' | e^{-i\hat{H}T} | q \rangle \\ &= \langle q' | \left(e^{-i\hat{H}\epsilon} \right) \dots \left(e^{-i\hat{H}\epsilon} \right) | q \rangle, \end{aligned} \quad (2.4)$$

where the expression in the second line contains n factors. Inserting the completeness relation $n - 1$ times,

$$\langle q't'|qt \rangle \int \prod_{k=1}^{n-1} dq_k \langle q' | (e^{-i\hat{H}\epsilon}) | q_{n-1} \rangle \langle q_{n-1} | (e^{-i\hat{H}\epsilon}) | q_{n-2} \rangle \dots \langle q_1 | (e^{-i\hat{H}\epsilon}) | q \rangle. \quad (2.5)$$

For small ϵ we can expand the exponential to first order and evaluate the matrix elements:

$$\begin{aligned} e^{-i\hat{H}\epsilon} &= 1 - i\hat{H}\epsilon + \mathcal{O}(\epsilon^2) \\ \hat{H} &= \frac{1}{2}\hat{P}^2 + V(\hat{Q}). \end{aligned} \quad (2.6)$$

From the potential energy we get

$$\begin{aligned} \langle q_k | V(\hat{Q}) | q_{k-1} \rangle &= V(q_{k-1}) \langle q_k | q_{k-1} \rangle \\ &\approx V\left(\frac{q_k + q_{k-1}}{2}\right) \delta(q_k - q_{k-1}) \\ &= \int \hat{d}p V\left(\frac{q_k + q_{k-1}}{2}\right) e^{ip(q_k - q_{k-1})}. \end{aligned} \quad (2.7)$$

In order to evaluate the contribution of the kinetic term, we introduce eigenstates of the momentum operator $|p\rangle$

$$\hat{P} |p\rangle = p |p\rangle \quad (2.8)$$

$$\langle q | p \rangle = e^{ipq}, \quad (2.9)$$

and use the completeness of the states $|p\rangle$:

$$\langle q_k | \hat{P}^2 | q_{k-1} \rangle = \int \hat{d}p p^2 e^{ip(q_k - q_{k-1})}. \quad (2.10)$$

Hence, remembering that the exponential of the Hamiltonian has been expanded to linear order,

$$\begin{aligned} \langle q_k | e^{-i\hat{H}\epsilon} | q_{k-1} \rangle &= \langle q_k | 1 - i\epsilon \left(\frac{1}{2}\hat{P}^2 + V(\hat{Q}) \right) | q_{k-1} \rangle \\ &= \int \hat{d}p \left[1 - i\epsilon \left(\frac{1}{2}p^2 + V\left(\frac{q_k + q_{k-1}}{2}\right) \right) \right] e^{ip(q_k - q_{k-1})} \\ &= \int \hat{d}p \exp \left[i\epsilon \left(p \frac{q_k - q_{k-1}}{\epsilon} - H\left(p, \frac{q_k + q_{k-1}}{2}\right) \right) \right] + \mathcal{O}(\epsilon^2). \end{aligned} \quad (2.11)$$

Therefore

$$\langle q't'|qt \rangle = \lim_{n \rightarrow \infty} \int \prod_{k=1}^{n-1} dq_k \prod_{j=1}^n \hat{d}p_j \exp \left\{ i\epsilon \sum_{m=1}^n \left[p_m \frac{q_m - q_{m-1}}{\epsilon} - H\left(p_m, \frac{q_m + q_{m-1}}{2}\right) \right] \right\}, \quad (2.12)$$

where $q_0 = q$, and $q_n = q'$. The limit above defines the *path integral* evaluation of the quantum amplitude, which we denote

$$\boxed{\langle q't'|qt \rangle = \int \mathcal{D}q \mathcal{D}p \exp \left\{ i \int_t^{t'} dt [\dot{q}p - H(p, q)] \right\}.} \quad (2.13)$$

For Hamiltonians where the momentum dependance is **only quadratic**, the expression above can be simplified by performing the integral over the momenta p_j :

$$\int \hat{d}p \exp \left\{ i\epsilon \left[p \frac{q_k - q_{k-1}}{\epsilon} - \frac{1}{2}p^2 \right] \right\} = (2\pi i\epsilon)^{-1/2} \exp \left\{ i\epsilon \frac{1}{2} \left(\frac{q_k - q_{k-1}}{\epsilon} \right)^2 \right\}. \quad (2.14)$$

Using this result, we can rewrite the path integral in Eq. 2.13 as

$$\langle q't'|qt \rangle = \lim_{n \rightarrow \infty} (2\pi i \epsilon)^{-n/2} \int \prod_{k=1}^{n-1} dq_k \exp \left\{ i \epsilon \sum_{m=1}^n \left[\frac{1}{2} \left(\frac{q_k - q_{k-1}}{\epsilon} \right)^2 - V \left(\frac{q_m + q_{m-1}}{2} \right) \right] \right\}, \quad (2.15)$$

and, assuming that the limit exists, we have obtained the definition of the path integral as an integral over the position of the system only:

$$\boxed{\langle q't'|qt \rangle = \int \mathcal{D}q \exp \left\{ i \int_t^{t'} d\tau L(q, \dot{q}) \right\}.} \quad (2.16)$$

2.2 Correlators

We are now going to work out expressions for the matrix element of the position operator in between the initial and final state considered above. The first example we are going to consider is the matrix element

$$\langle q't'|\hat{Q}(\bar{t})|qt \rangle = \langle q'|e^{-i\hat{H}(t'-t)}\hat{Q}e^{-i\hat{H}(\bar{t}-t)}|q \rangle, \quad (2.17)$$

where we assume $t < \bar{t} < t'$, and we have used

$$\hat{Q}(t) = e^{i\hat{H}t}\hat{Q}e^{-i\hat{H}t}. \quad (2.18)$$

Proceeding as we did in the previous section, we can write

$$\langle q't'|\hat{Q}(\bar{t})|qt \rangle = \langle q'|(e^{-i\hat{H}\epsilon}) \dots (e^{-i\hat{H}\epsilon}) \hat{Q} (e^{-i\hat{H}\epsilon}) \dots (e^{-i\hat{H}\epsilon})|q \rangle \quad (2.19)$$

where we assumed that $\bar{t} = t_k$, and the first and second ellipses denote respectively $n - k$ and k factors of $(e^{-i\hat{H}\epsilon})$. Performing the same manipulations as before we obtain

$$\langle q't'|\hat{Q}(\bar{t})|qt \rangle = \lim_{n \rightarrow \infty} \int \prod_{k=1}^{n-1} dq_k \prod_{j=1}^n \hat{p}_j q_k \exp \left\{ i \epsilon \sum_{m=1}^n \left[p_m \frac{q_m - q_{m-1}}{\epsilon} - H \left(p_m, \frac{q_m + q_{m-1}}{2} \right) \right] \right\}. \quad (2.20)$$

Note the extra factor of q_k in the integrand. We write this limit as

$$\langle q't'|\hat{Q}(\bar{t})|qt \rangle = \int_{qq'} \mathcal{D}q q(\bar{t}) \exp \left\{ i \int_t^{t'} d\tau \mathcal{L}(q, \dot{q}) \right\}. \quad (2.21)$$

The two-point function is

$$\langle q't'|\hat{Q}(\bar{t}_1)\hat{Q}(\bar{t}_2)|qt \rangle, \quad (2.22)$$

where we assume $\bar{t}_1 = t_k$, $\bar{t}_2 = t_\ell$, $t < \bar{t}_2 < \bar{t}_1 < t'$. Proceeding as above yields

$$\langle q't'|\hat{Q}(\bar{t}_1)\hat{Q}(\bar{t}_2)|qt \rangle = \int_{qq'} \mathcal{D}q q(\bar{t}_1)q(\bar{t}_2) \exp \left\{ \int_t^{t'} d\tau \mathcal{L}(q, \dot{q}) \right\}. \quad (2.23)$$

There is a subtlety here that is worth noting. The ordering of times in Eq. 2.22 matters, while it does not in Eq. 2.23. Thus, it is more accurate to say that the path integral we have defined is the result of the *time-ordered product* of the operators

$$\begin{aligned} \langle q't'|T \left(\hat{Q}(\bar{t}_1)\hat{Q}(\bar{t}_2) \right) |qt \rangle &= \langle q't'| \left(\theta(\bar{t}_1 - \bar{t}_2) \hat{Q}(\bar{t}_1)\hat{Q}(\bar{t}_2) + \theta(\bar{t}_2 - \bar{t}_1) \hat{Q}(\bar{t}_2)\hat{Q}(\bar{t}_1) \right) |qt \rangle \\ &= \int_{qq'} \mathcal{D}q q(\bar{t}_1)q(\bar{t}_2) \exp \left\{ \int_t^{t'} d\tau \mathcal{L}(q, \dot{q}) \right\}, \end{aligned} \quad (2.24)$$

which can be trivially extended to an arbitrary number of insertions of the operator \hat{Q} .

2.3 Generating functional

Let us begin with a revision of properties of functional derivatives. A functional F is a mapping from a function $u(x)$ to a number $F[u]$. The functional derivative describes the change of the functional to an infinitesimal variation of the function u :

$$u(x) \mapsto u(x) + \delta u(x). \quad (2.25)$$

We define

$$\delta F = F[u + \delta u] - F[u] = \int dx \frac{\delta F}{\delta u(x)} \delta u(x). \quad (2.26)$$

In particular we have the important identity

$$\frac{\delta}{\delta f(x)} f(y) = \delta(x - y). \quad (2.27)$$

Let $f(t)$ and $h(t)$ be two functions. Then we can add so-called source terms to the path integral:

$$\langle q't'|qt \rangle_{f,h} = \int \mathcal{D}q \mathcal{D}p \exp \left\{ i \int_t^{t'} d\tau [p(\tau) \dot{q}(\tau) - H(p(\tau), q(\tau)) + f(\tau)q(\tau) + h(\tau)p(\tau)] \right\}. \quad (2.28)$$

Then we have the useful property that taking functional derivatives with respect to the source fields yields the previously discussed correlators. For example,

$$\left(\frac{1}{i} \frac{\delta}{\delta f(\bar{\tau})} \right) \langle q't'|qt \rangle_{f,h} = \int \mathcal{D}q \mathcal{D}p q(\bar{\tau}) \exp \left\{ i \int_t^{t'} d\tau [p(\tau) \dot{q}(\tau) - H(p(\tau), q(\tau)) + f(\tau)q(\tau) + h(\tau)p(\tau)] \right\}. \quad (2.29)$$

The general rule is thus

$$\boxed{\langle q't'|T(\hat{Q}(t_1) \dots \hat{Q}(t_n))|qt \rangle = \left(\frac{1}{i} \frac{\delta}{\delta f(t_1)} \right) \dots \left(\frac{1}{i} \frac{\delta}{\delta f(t_n)} \right) \langle q't'|qt \rangle_{f,h} \Big|_{f,h=0}.} \quad (2.30)$$

$$\boxed{\langle 0|0 \rangle_{f,h} = \int \mathcal{D}p \mathcal{D}q \exp \left\{ i \int_{-\infty}^{\infty} d\tau [p\dot{q} - (1 - i\epsilon)H(p, q) + f q + h p] \right\}} \quad (2.31)$$

3 Path Integral for the Scalar Field

The Lagrangian for a free real scalar field $\phi(x)$ is given by

$$\mathcal{L}_0(\phi(x)) = \frac{1}{2} \partial_\mu \phi(x) \partial^\mu \phi(x) - \frac{1}{2} m^2 \phi^2(x). \quad (3.1)$$

Note that \mathcal{L}_0 is invariant under Lorentz transformations. From the Euler-Lagrange equation, we derive the equation of motion for the field, the Klein-Gordon equation:

$$(\partial^2 + m^2)\phi(x) = 0. \quad (3.2)$$

Given the conjugate momentum

$$\Pi(x) = \frac{\partial \mathcal{L}_0}{\partial(\partial_0 \phi(x))} = \dot{\phi}(x) \quad (3.3)$$

we obtain the Hamiltonian

$$\begin{aligned} \mathcal{H}_0 &= \Pi(x) \dot{\phi}(x) - \mathcal{L}_0(\phi(x)) \\ &= \frac{1}{2} \Pi^2(x) + \frac{1}{2} \sum_{k=1}^3 (\partial_k \phi(x))^2 + \frac{1}{2} m^2 \phi^2(x). \end{aligned} \quad (3.4)$$

The quadratic Hamiltonian is the generalization of the harmonic oscillator to the case where we have an infinite number of canonical coordinates, indexed by the continuous spatial coordinate \underline{x} .

3.1 Path Integral

The vacuum-to-vacuum amplitude in the presence of a source field $J(x)$ is the straightforward generalization of the expression we have derived for the quantum mechanical system:

$$\begin{aligned} q(t) &\mapsto \phi(t, \underline{x}) \\ \hat{Q}(t) &\mapsto \hat{\phi}(t, \underline{x}) \text{ (operator)} \\ f(t) &\mapsto J(t, \underline{x}) \text{ (source)}. \end{aligned} \quad (3.5)$$

By analogy with the QM computation, we can write the expression for the path integral representation of the vacuum amplitude

$$Z_0[J] = \langle 0|0 \rangle_J = \int \mathcal{D}\phi \exp \{iS_0 + J \cdot \phi\}, \quad (3.6)$$

where

$$S_0[\phi] = \int d^D x \mathcal{L}_0(\phi(x)), \quad J \cdot \phi = \int d^D x J(x) \phi(x). \quad (3.7)$$

In momentum space, the kernel in the action is diagonal. Thus, introducing the Fourier transform

$$\phi(x) = \int d^D p e^{-ip \cdot x} \tilde{\phi}(p), \quad (3.8)$$

we can rewrite the kinetic term:

$$\begin{aligned} \int d^D x \partial_\mu \phi(x) \partial^\mu \phi(x) &= \int d^D x \int_{p,p'} (-ip_\mu) e^{-ip \cdot x} \tilde{\phi}(p) (-ip'_\mu) e^{-ip' \cdot x} \tilde{\phi}(p') \\ &= \int_p p^2 \tilde{\phi}(p) \tilde{\phi}(-p) = \int_p p^2 |\tilde{\phi}(p)|^2 \end{aligned} \quad (3.9)$$

since we have a factor of $\delta(p + p')$ from the exponential and $\tilde{\phi}(-p) = \tilde{\phi}^*(p)$. Hence the action for the field in momentum space can be written as

$$S_0[\phi] + J \cdot \phi = \frac{1}{2} \int_p \left\{ \tilde{\phi}(-p) [p^2 - m^2 + i\epsilon] \tilde{\phi}(p) + \tilde{J}(p) \tilde{\phi}(-p) + \tilde{J}(-p) \tilde{\phi}(p) \right\}. \quad (3.10)$$

Note that the contribution from the ϵ term to the exponential is

$$\exp \left\{ -\epsilon \int_p |\tilde{\phi}(p)|^2 \right\} \quad (3.11)$$

which is clearly convergent for large values of $|\tilde{\phi}(p)|$. The path integral for the free field is a simple extension of the Gaussian integrals that we have discussed previously. The Gaussian integral can be computed by performing a shift of the integration variables

$$\tilde{\chi}(p) = \tilde{\phi}(p) + \frac{\tilde{J}(p)}{p^2 - m^2 + i\epsilon}, \quad (3.12)$$

so that

$$S_0[\chi] + J \cdot \chi = \frac{1}{2} \int_p \left\{ \tilde{\chi}(-p) [p^2 - m^2 + i\epsilon] \tilde{\chi}(p) - \tilde{J}(-p) \frac{1}{p^2 - m^2 + i\epsilon} \tilde{J}(p) \right\}. \quad (3.13)$$

Then up to a normalization factor (i.e. the kinetic term)

$$\begin{aligned} Z_0[J] &\propto \exp \frac{i}{2} \int_p \tilde{J}(-p) \frac{-1}{p^2 - m^2 + i\epsilon} \tilde{J}(p) \\ &= \exp -\frac{i}{2} \int d^D x d^D x' J(x) \Delta(x, x') J(x') \end{aligned} \quad (3.14)$$

where we have introduced the *Feynman propagator* Δ .

$$\Delta(x, x') = \Delta(x - x') = \int d^{-ip \cdot (x - x')} \frac{1}{p^2 - m^2 + i\epsilon}. \quad (3.15)$$

Recalling Eq. 2.30 and the fact that $Z_0[J] = \langle 0|0 \rangle_J$, we can write correlators as

$$\begin{aligned} \langle 0|T(\phi(x_1)\phi(x_2))|0 \rangle_0 &= \left(\frac{1}{i} \frac{\delta}{\delta J(x_1)} \right) \left(\frac{1}{i} \frac{\delta}{\delta J(x_2)} \right) Z_0[J] \Big|_{J=0} \\ &= i\Delta(x_1 - x_2). \end{aligned} \quad (3.16)$$

Further correlators are obtained by taking more derivatives. They can be computed in the free theory using Wick's theorem, again following the arguments we used for Gaussian integrals, e.g.:

$$\langle 0|T(\phi(x_1)\phi(x_2)\phi(x_3)\phi(x_4))|0 \rangle_0 = i^2[\Delta(x_1 - x_2)\Delta(x_3 - x_4) + \Delta(x_1 - x_3)\Delta(x_2 - x_4) + \Delta(x_1 - x_4)\Delta(x_2 - x_3)]. \quad (3.17)$$

3.2 Interacting Theory

Let us now add an interaction term in the Lagrangian:

$$\mathcal{L}(\phi(x)) = \mathcal{L}_0(\phi(x)) + V(\phi(x)). \quad (3.18)$$

We are going to consider several examples, e.g. ϕ^3 theory:

$$V(\phi(x)) = \frac{1}{3!} g \phi^3(x). \quad (3.19)$$

Denoting by $S_0[\phi]$ the action of the free theory, we can write the path integral for the interacting theory

$$Z[J] = \langle 0|0 \rangle_J = \int \mathcal{D}\phi \exp \left\{ i \left(S_0[\phi] + \int d^D x V(\phi(x)) \right) \right\} \quad (3.20)$$

By performing the same manipulations as Eq. 1.28, we obtain

$$Z[J] = \exp \left\{ i \int d^D x V \left(\frac{1}{i} \frac{\delta}{\delta J(x)} \right) \right\} Z_0[J], \quad (3.21)$$

and by expanding both exponentials and substituting in Eq. 3.14, we obtain a perturbative expansion to our generating functional:

$$Z[J] \propto \sum_{V=0}^{\infty} \frac{1}{V!} \left[\frac{ig}{3!} \int d^D x \left(\frac{1}{i} \frac{\delta}{\delta J(x)} \right)^3 \right]^V \times \sum_{P=0}^{\infty} \frac{1}{P!} \left[-\frac{i}{2} \int d^D y d^D z J(y) \Delta(y - z) J(z) \right]^P \quad (3.22)$$

Since each power of P adds two sources and each power of V removes three sources, we are left with $E = 2P - 3V$ sources. The functional derivatives turn factors of J in the integrand into Dirac delta functions as following:

$$\frac{\delta}{\delta J(x)} \int \dots d^d y \dots J(y) \Delta(y, \dots) \dots = \int \dots \Delta(x, \dots) \dots \quad (3.23)$$

4 Scalar Field Correlators

4.1 The S -matrix

In order to compute the quantum amplitude for a physical process involving arbitrary numbers of particles in the initial and final state, we need to compute the overlap of a state prepared in the distant past (the

so-called *in* state), with the resulting final state in the distant future (the so-called *out* state). If we want to describe a $2 \rightarrow n$ process — like a pp collision at the LHC — we need to compute

$$\langle p_1, \dots, p_n; \text{out} | k_1, k_2; \text{in} \rangle. \quad (4.1)$$

The S -matrix allows us to express this scalar product between in- and out-states in terms of the states defined at any common reference time :

$$\langle p_1, \dots, p_n; \text{out} | k_1, k_2; \text{in} \rangle = \langle p_1, \dots, p_n | S | k_1, k_2 \rangle. \quad (4.2)$$

It is usual to separate the S -matrix into the identity operator, corresponding to the particles not interacting, plus a non-trivial *transition matrix* which is usually denoted T :

$$S = 1 + iT. \quad (4.3)$$

4.2 Field correlators

As discussed in the previous lecture, the field correlators are obtained from the partition function by taking functional derivatives:

$$\begin{aligned} G^{(n)}(x_1, \dots, x_n) &= \langle 0 | T(\phi(x_1) \dots \phi(x_n)) | 0 \rangle \\ &= \left(\frac{1}{i} \frac{\delta}{\delta J(x_1)} \right) \dots \left(\frac{1}{i} \frac{\delta}{\delta J(x_n)} \right) Z[J] \Big|_{J=0}. \end{aligned} \quad (4.4)$$

The perturbative definition of the path integral has led to an expansion where we can classify the terms according to the number of currents that appear, which we denoted by E above. Clearly, the only terms that contribute to an n -point function are the ones with $E = n$ ¹. Since we can express E in terms of P and V , the calculation of correlators can be expressed as a perturbative expansion in terms of V , and therefore in powers of coupling:

$$G^{(n)}(x_1, \dots, x_n) = \sum_V g^V G^{(n,V)}(x_1, \dots, x_n). \quad (4.5)$$

As we will see in future sections, the *LSZ reduction formula* provides a way to connect correlators in momentum space with S -matrix elements. This is intimately related to the structure of divergences which we can explore through *polology*. We shall now explore the calculation of two-point position space correlators and the resulting momentum-space expressions.

4.3 Polology

In this section, we shall learn some features about the analytic structure of field correlators and discuss their relevance in order to extract physical information from the correlators. These are general properties of correlators and do not rely on perturbation theory.

We begin by writing out an n -point correlator in momentum space:

$$\tilde{G}^{(n)}(p_1, \dots, p_n) = \int d^D x_1 \dots d^D x_n e^{ip_1 \cdot x_1} \dots e^{ip_n \cdot x_n} \langle T \{ \phi(x_1) \dots \phi(x_n) \} \rangle. \quad (4.6)$$

As the integral spans n time dimensions, we need to be careful regarding the ordering of the fields and the T -ordered product.

Let us focus on the contribution coming from the sector where the values x_1^0, \dots, x_r^0 are all larger than the values of x_{r+1}^0, \dots, x_n^0 for some value of r between 1 and $n-1$. We can write this condition as $\min\{x_1^0 \dots x_r^0\} > \max\{x_{r+1}^0 \dots x_n^0\}$. The contribution can then be written as

$$\begin{aligned} \tilde{G}(p_1, \dots, p_n) &\sim \int d^D x_1 \dots d^D x_n e^{ip_1 \cdot x_1} \dots e^{ip_n \cdot x_n} \\ &\quad \times \theta(\min\{x_1^0 \dots x_r^0\} - \max\{x_{r+1}^0 \dots x_n^0\}) \\ &\quad \times \langle 0 | T \{ \phi(x_1) \dots \phi(x_r) \} T \{ \phi(x_{r+1}) \dots \phi(x_n) \} | 0 \rangle. \end{aligned} \quad (4.7)$$

¹If there are less than n sources, then at least one of the functional derivatives will have nothing to act on, yielding zero. If there are more than n sources, then it will be set to zero when we evaluate the correlator, also yielding zero for the correlator.

We can insert a complete set of one-particle states between the two T -ordered products:

$$\langle 0|T\{\phi(x_1)\dots\phi(x_r)\}T\{\phi(x_{r+1})\dots\phi(x_n)\}|0\rangle = \sum_{\sigma} \int d\Omega_p \langle 0|T\{\phi(x_1)\dots\phi(x_r)\}|\mathbf{p},\sigma\rangle \langle \mathbf{p},\sigma|T\{\phi(x_{r+1})\dots\phi(x_n)\}|0\rangle \quad (4.8)$$

Let us evaluate this by doing some rewriting. Suppose that $f(z)$ is a function of a real number z , and $p = -i\frac{\partial}{\partial z}$ is the familiar quantum mechanical momentum operator. Then

$$\exp[ipa]f(z) = \exp\left[a\frac{\partial}{\partial z}\right]f(z) = \sum_{n=0}^{\infty} \frac{1}{n!} a^n \frac{\partial^n}{\partial z^n} f(z) = f(z+a). \quad (4.9)$$

Thus we can see that $\exp[ipa]$ is the generator of translations. This works in an arbitrary number of directions. However, in the case of operators, we act once on the left normally and once on the right with the inverse. Thus for an operator $F(z)$,

$$\exp[ip \cdot a]F(z)\exp[-ip \cdot a] = F(z+a). \quad (4.10)$$

By defining new integration variables,

$$x_i = x_1 + y_i, \quad \text{for } i = 2, \dots, r, \quad (4.11)$$

we can apply our knowledge of the momentum operator P as a generator of translations to write

$$\begin{aligned} \phi(x_1)\dots\phi(x_r) &= \phi(x_1)\dots\phi(x_1+y_i)\dots\phi(x_1+y_r) \\ &= e^{iP \cdot x_1}\phi(0)e^{-iP \cdot x_1}\dots e^{iP \cdot x_1}\phi(y_i)e^{-iP \cdot x_1}\dots e^{iP \cdot x_1}\phi(y_r)e^{-iP \cdot x_1} \\ &= e^{iP \cdot x_1}\phi(0)\dots\phi(y_r)e^{-iP \cdot x_1}. \end{aligned} \quad (4.12)$$

Since $e^{-iP \cdot x_1}|0\rangle = |0\rangle$ and $e^{-iP \cdot x_1}|p,\sigma\rangle = e^{-ip \cdot x_1}|p,\sigma\rangle$,

$$\langle 0|T\{\phi(x_1)\dots\phi(x_r)\}|\mathbf{p},\sigma\rangle = e^{-ip \cdot x_1}\langle 0|T\{\phi(0)\dots\phi(y_r)\}|\mathbf{p},\sigma\rangle. \quad (4.13)$$

The exact same procedure can be undergone for the other T -ordered product. The argument of the Heaviside theta function can be rewritten as

$$\min\{x_1^0\dots x_r^0\} - \max\{x_{r+1}^0\dots x_n^0\} = x_1^0 - x_{r+1}^0 + \min\{0\dots y_r^0\} - \max\{0\dots y_n^0\}. \quad (4.14)$$

Using the integral representation of the Heaviside theta function,

$$\theta(\tau) = -\frac{1}{2\pi i} \int_{-\infty}^{\infty} d\omega \frac{e^{-i\omega\tau}}{\omega + i\epsilon}, \quad (4.15)$$

yields

$$\begin{aligned} \tilde{G}(p_1, \dots, p_n) &\sim \int d^D x_1 d^D y_2 \dots d^D y_r d^D x_{r+1} d^D y_{r+2} \dots d^D y_n \\ &\times e^{ip_1 \cdot x_1} e^{ip_1 \cdot (x_1 + y_2)} \dots e^{ip_r \cdot (x_1 + y_r)} e^{ip_{r+1} \cdot x_{r+1}} e^{ip_{r+2} \cdot (x_{r+1} + y_{r+2})} \dots e^{ip_n \cdot (x_{r+1} + y_n)} \\ &\times -\frac{1}{2\pi i} \int \frac{d\omega}{\omega + i\epsilon} \exp\{-i\omega[x_1^0 - x_{r+1}^0 + \min\{0\dots y_r^0\} - \max\{0\dots y_n^0\}]\} \\ &\times \sum_{\sigma} \int d\Omega_p e^{-ip \cdot x_1} \langle 0|T\{\phi(0)\dots\phi(y_r)\}|\mathbf{p},\sigma\rangle e^{ip \cdot x_{r+1}} \langle \mathbf{p},\sigma|T\{\phi(0)\dots\phi(y_n)\}|0\rangle. \end{aligned} \quad (4.16)$$

Performing the integral over x_1 and x_{r+1} yields delta functions in terms of the arguments of the exponential: the four-momenta in the second line, the dummy variable ω in the third line and the extra exponentials due to Eq. 4.13 in the fourth line. We can split these delta functions into a product of delta functions of the

space and time components, yielding our final result

$$\begin{aligned}
\tilde{G}(p_1, \dots, p_n) &\sim \int d^D y_2 \dots d^D y_r d^D y_{r+2} \dots d^D y_n \\
&\times e^{ip_1 \cdot y_2} \dots e^{ip_r \cdot y_r} e^{ip_{r+2} \cdot y_{r+2}} \dots e^{ip_n \cdot y_n} \\
&\times -\frac{1}{2\pi i} \int \frac{d\omega}{\omega + i\epsilon} \exp\{-i\omega [\min\{0 \dots y_r^0\} - \max\{0 \dots y_n^0\}]\} \\
&\times \sum_{\sigma} \int d\Omega_p \langle 0|T\{\phi(0) \dots \phi(y_r)\}|\mathbf{p}, \sigma\rangle \langle \mathbf{p}, \sigma|T\{\phi(0) \dots \phi(y_n)\}|0\rangle \\
&\times \hat{\delta}(\mathbf{p} - \mathbf{p}_1 - \dots - \mathbf{p}_r) \hat{\delta}(E_p + \omega - p_1^0 - \dots - p_r^0) \\
&\times \hat{\delta}(\mathbf{p} - \mathbf{p}_{r+1} - \dots - \mathbf{p}_n) \hat{\delta}(E_p + \omega - p_{r+1}^0 - \dots - p_n^0).
\end{aligned} \tag{4.17}$$

Performing the Lorentz-invariant phase space integral² over the spatial components of p and ω transforms the delta functions into

$$\hat{\delta}(p_1 + \dots + p_n) \frac{1}{q^0 - E_p + i\epsilon} \frac{1}{2E_p} \tag{4.18}$$

where the factor of $1/2E_p$ comes from the integration measure $d\Omega_p$ and the four-momentum q is defined as

$$q = p_1 + \dots + p_r = -p_{r+1} - \dots - p_n. \tag{4.19}$$

To make it more clear that it is about q being on-shell, we can rewrite

$$\begin{aligned}
\lim_{q^0 \rightarrow E_p} \frac{1}{q^0 - E_p + i\epsilon} \frac{1}{2E_p} &= \lim_{q^0 \rightarrow E_p} \frac{1}{q^0 - E_p + i\epsilon} \frac{1}{E_p + E_p} \\
&= \lim_{q^0 \rightarrow E_p} \frac{1}{q^0 - E_p} \frac{1}{q^0 + E_p} \\
&= \lim_{q^0 \rightarrow E_p} \frac{1}{(q^0)^2 - E_p^2} \\
&= \lim_{q^0 \rightarrow E_p} \frac{1}{(q^0)^2 - \mathbf{q}^2 - m_{\text{phys}}^2} \\
&= \lim_{q^0 \rightarrow E_p} \frac{1}{q^2 - m_{\text{phys}}^2 + i\epsilon}.
\end{aligned} \tag{4.20}$$

The important result here is that the correlators in momentum space have a pole singularity **whenever** $q = p_1 + \dots + p_r$ **goes on-shell**. The residue at this pole will be

$$\begin{aligned}
\delta(p_1 + \dots + p_n) \frac{i}{q^2 - m_{\text{phys}}^2 + i\epsilon} \sum_{\sigma} \int d^D y_2 \dots d^D y_r e^{ip_2 \cdot y_2} \dots e^{ip_r \cdot y_r} \langle 0|T\{\phi(0) \dots \phi(y_r)\}|\mathbf{q}, \sigma\rangle \\
\times \int d^D y_{r+2} \dots d^D y_n e^{ip_{r+2} \cdot y_{r+2}} \dots e^{ip_n \cdot y_n} \langle \mathbf{q}, \sigma|T\{\phi(0) \dots \phi(y_n)\}|0\rangle
\end{aligned} \tag{4.21}$$

4.4 LSZ reduction

An important corollary of what we have just covered is obtained by setting $r = 1$. In this case, the correlator in momentum space has a pole whenever p_1 is on-shell. However, p_1 is somewhat arbitrarily chosen — we could equivalently relabel the momenta and choose a different sector of the integral — and therefore the correlator in momentum space has a pole whenever the momentum of any of the fields is on-shell. It follows that an n -point correlation function has (at least) n poles, each corresponding to one of the momenta

²You're going to have to trust me on this one.

$p_i \rightarrow m_{\text{phys}}^2$. In a real scattering event, we want each of the particles involved in the correlator to be on-shell, so somehow we must assign a value to these poles. The most natural choice is of course the residue.

$$\begin{aligned} \langle p'_1 \dots p'_{m'}; \text{out} | p_1 \dots p_m; \text{in} \rangle &= \langle p'_1 \dots p'_{m'} | S | p_1 \dots p_m \rangle \\ &= \lim_{p_j^2, p_k'^2 \rightarrow m_{\text{phys}}^2} \prod_{k=1}^{m'} (p_k'^2 - m_{\text{phys}}^2 + i\epsilon) \prod_{j=1}^m (p_j^2 - m_{\text{phys}}^2 + i\epsilon) \\ &\quad \times \tilde{G}^{(m+m')}(p_1, \dots, p_m, -p'_1, \dots, -p'_{m'}), \end{aligned} \quad (4.22)$$

where $n = m + m'$, and the fields are normalised so that

$$\langle \mathbf{p} | \phi(0) | 0 \rangle = 1. \quad (4.23)$$

Therefore, we can compute the S -matrix elements by adapting the Feynman rules given in Section 4.2:

- We associate an outgoing momentum to the external lines that correspond to particles in the final state, giving a relative minus sign compared to the normal Feynman rules when all momenta are taken to be inwards.
- We multiply each external line by a factor of $-i(p^2 - m_{\text{phys}}^2 + i\epsilon)$, this is part of computing the residue. Correlators multiplied by these factors are called *truncated correlators* or *amputated correlators*.

4.5 Källén-Lehmann representation

Let us now come back to the 2-point function, and find a representation that allows us to extract some physical information about the scalar field. We define the full propagator as

$$i\Delta_F(x-y) = \langle 0 | T \{ \phi(x) \phi(y) \} | 0 \rangle, \quad (4.24)$$

and define the field so that

$$\langle 0 | \phi(x) | 0 \rangle = 0, \text{ and } \langle \mathbf{p} | \phi(0) | 0 \rangle = 1, \quad (4.25)$$

where $|\mathbf{p}\rangle$ represents the physical one-particle state, and we have dropped the dependence on σ . As usual the full propagator in momentum space is defined by taking the Fourier transform:

$$\tilde{\Delta}_F(p) = \int d^D x e^{ip \cdot (x-y)} \Delta_F(x-y). \quad (4.26)$$

With these conventions, the free theory result for the propagator is

$$\tilde{\Delta}(p) = \frac{i}{p^2 - m^2 + i\epsilon}, \quad (4.27)$$

which has a pole at $p^2 = m^2$. For the free particle we find a pole in the propagator, at a value which coincides with the parameter in the Lagrangian.

For the interacting theory, we can derive a general expression, which again does not rely on the perturbative definition of the two-point function. Let us first consider the case where $x^0 > y^0$. We can insert a complete set of zero-, one-, and multi-particle states in between:

$$\begin{aligned} \langle 0 | T \{ \phi(x) \phi(y) \} | 0 \rangle &= \langle 0 | \phi(x) \phi(y) | 0 \rangle \\ &= \langle 0 | \phi(x) | 0 \rangle \langle 0 | \phi(y) | 0 \rangle + \\ &\quad + \int d\Omega_p \langle 0 | \phi(x) | \mathbf{p} \rangle \langle \mathbf{p} | \phi(y) | 0 \rangle + \\ &\quad + \sum_n \int d\Omega_P \langle 0 | \phi(x) | \mathbf{P}, n \rangle \langle \mathbf{P}, n | \phi(y) | 0 \rangle. \end{aligned} \quad (4.28)$$

Using the same trick as in Eq. 4.13, the conventions defined in Eq. 4.25 simplify the expression above to

$$\int d\Omega_p e^{-ip \cdot (x-y)} + \sum_n \int d\Omega_P e^{-iP \cdot (x-y)} |\langle \mathbf{P}, n | \phi(0) | 0 \rangle|^2. \quad (4.29)$$

Because we are working with a scalar field, the matrix element $\langle \mathbf{p}, n | \phi(0) | 0 \rangle$ is invariant under Lorentz transformations, and therefore can only depend on p via the invariant mass M^2 . We can therefore introduce the *spectral density*

$$\rho(s) \equiv \sum_n |\langle \mathbf{p}, n | \phi(0) | 0 \rangle|^2 \delta(s - M^2), \quad (4.30)$$

and write the two-point correlator as

$$\langle 0 | T \{ \phi(x) \phi(y) \} | 0 \rangle = \int d\Omega_p e^{-ip \cdot (x-y)} + \int_{4m_{\text{phys}}^2}^{\infty} ds \rho(s) \int d\Omega_P e^{-iP \cdot (x-y)}. \quad (4.31)$$

We start the integral at $4m_{\text{phys}}^2$ because we must have $M \geq 2m_{\text{phys}}$ for a multiparticle state, so $M^2 \geq 4m_{\text{phys}}^2$. Similar manipulations for $y^0 > x^0$ yield

$$\langle 0 | \phi(y) \phi(x) | 0 \rangle = \int d\Omega_p e^{-ip \cdot (y-x)} + \int_{4m_{\text{phys}}^2}^{\infty} ds \rho(s) \int d\Omega_P e^{-iP \cdot (y-x)}. \quad (4.32)$$

Collecting both contributions to the T -ordered product and using

$$\frac{1}{i} \int \hat{d}^D p \frac{e^{-ip \cdot (x-y)}}{p^2 - \mu^2 + i\epsilon} = \theta(x^0 - y^0) \int d_p^\Omega e^{-ip \cdot (x-y)} + \theta(y^0 - x^0) \int d_p^\Omega e^{-ip \cdot (y-x)} \Big|_{p^0 = E_{\mathbf{p}}} \quad (4.33)$$

we finally obtain

$$i \langle 0 | T \{ \phi(y) \phi(x) \} | 0 \rangle = \int \hat{d}^D p e^{-ip \cdot (x-y)} \left[\frac{i}{p^2 - m_{\text{phys}}^2 + i\epsilon} + \int_{4m_{\text{phys}}^2}^{\infty} ds \rho(s) \frac{i}{p^2 - s + i\epsilon} \right]. \quad (4.34)$$

Eq. 4.34 allows us to read the expression for the full propagator in momentum space, also known as the *Källén-Lehmann representation*:

$$\boxed{\tilde{\Delta}_F(p) = \frac{i}{p^2 - m_{\text{phys}}^2 + i\epsilon} + \int_{4m_{\text{phys}}^2}^{\infty} ds \rho(s) \frac{i}{p^2 - s + i\epsilon}} \quad (4.35)$$

We see that the two-point correlator of a field ϕ that satisfies the conditions in Eq. 4.25 has a pole for $p^2 = m_{\text{phys}}^2$, with a residue exactly equal to one. Note that the field ϕ does not need to be the field that appears in the Lagrangian. Knowledge about the multiparticle states of the theory is encoded in the spectral density part of the Källén-Lehmann representation.

4.6 Ward identities

The final example of relations between correlators that we are going to discuss are so-called *Ward identities*. Ward identities are equalities between field correlators that are obtained as a consequence of symmetries of the system. In classical mechanics, symmetries of the action translate into conserved currents according to Noether's theorem. As we will show in this section, the analogue of current conservation in quantum field theory is precisely the Ward identity.

In order to derive the identities, let us start by considering a symmetry transformation of the field, i.e. a transformation

$$\phi(x) \mapsto \phi'(x) = \phi(x) + \epsilon \delta(\phi(x)), \quad (4.36)$$

such that for constant ϵ the action is unchanged. If we introduce a dependence on the space-time coordinate, $\epsilon(x)$, then the variation of the action can be written

$$\begin{aligned}\delta S &= \int d^D x \frac{\delta S}{\delta \phi(x)} \epsilon(x) \delta \phi(x) \\ &= - \int d^D x \epsilon(x) \partial_\mu j^\mu(x),\end{aligned}\tag{4.37}$$

where $j^\mu(x)$ is precisely the Noether current that is conserved in the classical theory. In order to derive the Ward identities, we use Eq. 4.36 to perform a change of integration variables in the functional integral

$$\int \mathcal{D}\phi e^{iS[\phi]} O(\phi) = \int \mathcal{D}\phi' e^{iS[\phi']} O(\phi'),\tag{4.38}$$

where O is an arbitrary function of the field ϕ and then expand the RHS to first order in ϵ :

$$\int \mathcal{D}\phi e^{iS[\phi]} O(\phi) = \int \mathcal{D}\phi e^{iS[\phi]} (1 + i\delta S[\phi]) (O(\phi) + \delta O).\tag{4.39}$$

We can now substitute the expressions for δS and δO :

$$\int \mathcal{D}\phi e^{iS[\phi]} \left(1 - i \int d^D x \epsilon(x) \partial_\mu j^\mu(x) \right) \left(O(\phi) + \int d^D x \frac{\delta O(\phi)}{\delta \phi(x)} \epsilon(x) \delta \phi(x) \right)\tag{4.40}$$

Thus the difference between the RHS and the LHS of Eq. 4.38, ignoring terms of $\mathcal{O}(\epsilon^2)$, is

$$\int \mathcal{D}\phi e^{iS[\phi]} \left\{ -i \int d^D x \epsilon(x) \partial_\mu j^\mu(x) O(\phi) + \int d^D x \frac{\delta O(\phi)}{\delta \phi(x)} \epsilon(x) \delta \phi(x) \right\} = 0.\tag{4.41}$$

Rearranging the terms above allows us to write the identity in a way that makes its physical content more obvious

$$\int d^D x \epsilon(x) \left\{ -i \langle \partial_\mu j^\mu(x) O(\phi) \rangle + \left\langle \frac{\delta O(\phi)}{\delta \phi(x)} \delta \phi(x) \right\rangle \right\} = 0.\tag{4.42}$$

Eq. 4.42 is sometimes referred to as an *integrated Ward identity*. Since it has to be satisfied for every function $\epsilon(x)$, we can derive the *Ward identity*:

$$-i \langle \partial_\mu j^\mu(x) O(\phi) \rangle + \left\langle \frac{\delta O(\phi)}{\delta \phi(x)} \delta \phi(x) \right\rangle = 0.$$

(4.43)

There are two important physical results encoded in Eq. 4.43.

1. Symmetry in QFT translates into a relation between correlators. This is true beyond perturbation theory and is used in defining the renormalization conditions in QFT.
2. Current conservation in QFT is realised at the level of the insertion of $\partial_\mu j^\mu(x)$ in field correlators, up to the terms that come from the variation of O . If O is a product of local fields, this variation is localised in space-time, i.e. the contributions are all proportional to Dirac deltas. These terms are called *contact terms*.

Note that in deriving the Ward identity above we have assumed that the integration measure $\mathcal{D}(\phi)$ is invariant, i.e. $\mathcal{D}\phi = \mathcal{D}\phi'$. There are examples where the measure is **not invariant**, which lead to extra terms in the Ward identities. In these cases the Ward identities are called *anomalous*.

5 Fermion Fields

5.1 Fermionic Path Integral

For the case of fermion fields, we want to define the path integral following the recipe we used for the scalar field. We will treat the fields ψ and $\bar{\psi}$ as Grassmann variables, i.e.

$$\psi_\alpha(x), \psi_\beta(y) = 0.\tag{5.1}$$

In order to have a consistent implementation of the anticommuting properties of the fermion fields, the functional derivative with respect to a Grassmann variable must be a Grassmann variable itself. As a consequence

$$\frac{\delta^2 F}{\delta\psi_\alpha(x)\delta\psi_\beta(y)} = -\frac{\delta^2 F}{\delta\psi_\beta(y)\psi_\alpha(x)} \quad (5.2)$$

and

$$\frac{\delta^2 F}{\delta\psi_\alpha(x)\delta\psi_\alpha(x)} = 0. \quad (5.3)$$

In the definition of the generating functional (recall Eqs. 3.6, 3.7 for the scalar case, $\int d^D x J(x)\phi(x)$ was the source term in the action), we introduce independent sources for ψ and $\bar{\psi}$:

$$S_0 = \int d^D x \mathcal{L}_0 + \int d^D x [\bar{\eta}(x)\psi(y) + \bar{\psi}(y)\eta(y)], \quad (5.4)$$

such that

$$\frac{\delta}{\delta\eta(y)} \int d^D x [\bar{\eta}(x)\psi(y) + \bar{\psi}(y)\eta(y)] = -\bar{\psi}(x) \quad (5.5)$$

$$\frac{\delta}{\delta\bar{\eta}(x)} \int d^D x [\bar{\eta}(x)\psi(y) + \bar{\psi}(y)\eta(y)] = \psi(x). \quad (5.6)$$

Note the difference of ordering between the two terms in brackets.

The action for the free Dirac field is

$$S_0[\psi, \bar{\psi}] = \int d^D x \bar{\psi}(x)(i\not{\partial} - m)\psi(x). \quad (5.7)$$

Using the rules above for the functional derivative, we can find the classical equation of motion, i.e. Dirac's equation

$$\frac{\delta}{\delta\bar{\psi}(x)} S_0[\psi, \bar{\psi}] = 0 \implies (i\not{\partial} - m)\psi(x). \quad (5.8)$$

By analogy with the scalar case, we can write the generating functional for the free theory:

$$\begin{aligned} Z_0[\bar{\eta}, \eta] &= \int \mathcal{D}\psi \mathcal{D}\bar{\psi} \exp [i(S_0[\psi, \bar{\psi}] + \bar{\eta} \cdot \psi + \bar{\psi} \cdot \eta)] \\ &= \exp \left[- \int d^D x d^D y \bar{\eta}(x) S(x-y) \eta(y) \right]. \end{aligned} \quad (5.9)$$

By a similar method to Eq. 3.12, the Feynman propagator for the Dirac field is

$$S(x-y) = \int_p e^{-ip \cdot (x-y)} \frac{i(\not{p} + m)}{p^2 - m^2 + i\epsilon}. \quad (5.10)$$

Note that, just like in the case of the scalar field, the propagator is the inverse of the quadratic term in the action. The propagator is a 4×4 matrix in spin space, which we can write explicitly:

$$\langle 0|T \{ \psi_\alpha(x) \bar{\psi}_\beta(y) \} |0\rangle = S_{\alpha\beta}(x-y) = \int_p e^{-ip \cdot (x-y)} \frac{i((\gamma^\mu)_{\alpha\beta} p_\mu + m\delta_{\alpha\beta})}{p^2 - m^2 + i\epsilon}. \quad (5.11)$$

Because of the linear term p in the propagator the fermionic propagator is not symmetric in its arguments, and will be denoted with an arrow pointing from one end to the other:

$$S(x-y) = x \longrightarrow y \quad (5.12)$$

Correlators of fermion fields are computed by taking derivatives with respect to the source fields:

$$\langle 0|T \{ \psi_{\alpha_1} \dots \bar{\psi}_{\beta_1}(y_1) \dots \} |0\rangle_0 = \left(\frac{1}{i} \frac{\delta}{\delta\bar{\eta}_{\alpha_1}(x_1)} \right) \dots \left(i \frac{\delta}{\delta\eta_{\beta_1}(y_1)} \right) \dots Z_0[\eta, \bar{\eta}] \Big|_{\eta=\bar{\eta}=0} \quad (5.13)$$

In the interacting theory, if the interactions are specified by a potential $V(\psi, \bar{\psi})$, the generating functional is defined as

$$Z[\eta, \bar{\eta}] = \exp \left[i \int d^D x V \left(i \frac{\delta}{\delta \eta(x)}, \frac{1}{i} \frac{\delta}{\delta \bar{\eta}(x)} \right) \right], \quad (5.14)$$

and the normalization is fixed by requiring

$$Z[0, 0] = 1. \quad (5.15)$$

Thus, for a perturbative expansion to our generating functional, we require a double expansion in the powers of the interaction and in powers of the propagator.