

THE MINIMAL SUPERSYMMETRIC STANDARD MODEL

Part II: Phenomenology

Mathieu Kaltschmidt & Jonah Cedric Strauß

ITP Heidelberg

Supersymmetry Seminar

supervised by

Prof. Jörg Jäckel

Heidelberg, February 15th 2021

Outline

1. Overview
2. Phenomenology of the MSSM
3. A Top-Down Approach to Low-Energy Phenomenology: mSUGRA
4. Grand Unification and SUSY
5. Going Beyond the MSSM
6. Summary and Outlook

Outline

- 1 Overview
- 2 Phenomenology of the MSSM
- 3 A Top-Down Approach to Low-Energy Phenomenology: mSUGRA
- 4 Grand Unification and SUSY
- 5 Going Beyond the MSSM
- 6 Summary and Outlook

What have we learned so far and what needs to be discussed?


- **State of the Art:** We promoted the SM to the MSSM-124 and found a variety of new degrees of freedom that need to be taken into account in further investigations.
- Now we want to understand how to make sense of a theoretical model with $\mathcal{O}(100)$ parameters.
- **Central Question:** Is it possible to reduce this huge number of parameters to a reasonable number that may be tested in future experiments?¹
- **Phenomenology of the MSSM:** Focus on the “pMSSM” and Gauge Coupling Unification.
- Which problems are still present? Why might the “minimal” SSM not be enough?

¹Experimental details → talk next week!

Outline

- 1 Overview
- 2 Phenomenology of the MSSM**
- 3 A Top-Down Approach to Low-Energy Phenomenology: mSUGRA
- 4 Grand Unification and SUSY
- 5 Going Beyond the MSSM
- 6 Summary and Outlook


How to test the MSSM?



Some Observations

This huge additional parameter space in general comes with some severe problems :

- The respective **lepton numbers** L_e , L_μ and L_τ are not conserved \nexists
- **Flavor-changing neutral currents** (FCNCs) are unsuppressed \nexists
- New sources of **CP-violation** are present \nexists



For all of the above mentioned problems we have rather strict experimental bounds, for example:

Want to avoid contributions to **electric dipole moments** \implies complex phases of the gaugino mass parameters, A -parameters and $|\mu| \lesssim 10^{-2} - 10^{-3}$ for TeV-ish sfermion and gaugino masses [12].

General Requirements

We use these phenomenological constraints to define the **phenomenological MSSM** (pMSSM) [11]:

- No additional sources of CP-violation \implies all phases in the soft-SUSY-breaking potential zero
- No FCNCs \implies simple, diagonal structure of the sfermion and trilinear coupling matrices
- First and second generation universality \implies soft-SUSY-breaking scalar masses + trilinear couplings² A^u , A^d and A^ℓ are the same for the first two generations

In addition, for different regions of the parameter space, one can use even more severe constraints³!

²Usually they are just set to zero for the first two generations

³In actual experiments only very simplified models with two or three parameters are tested.

Which Parameters do we need?

This already reduces the number of relevant parameters by a large amount, a summary is given below:

Which Parameters do we need?

This already reduces the number of relevant parameters by a large amount, a summary is given below:

- $\tan \beta = v_u/v_d$, the ratio of the VEVs in the two-Higgs doublet model

Which Parameters do we need?

This already reduces the number of relevant parameters by a large amount, a summary is given below:

- $\tan \beta = v_u/v_d$, the ratio of the VEVs in the two-Higgs doublet model
- $m_A^2 = 2m_{12}^2/\sin 2\beta$, the mass of the pseudoscalar Higgs boson

Which Parameters do we need?

This already reduces the number of relevant parameters by a large amount, a summary is given below:

- $\tan \beta = v_u/v_d$, the ratio of the VEVs in the two-Higgs doublet model
- $m_A^2 = 2m_{12}^2/\sin 2\beta$, the mass of the pseudoscalar Higgs boson
- M_1 , M_2 and M_3 , the bino, wino and gluino masses

Which Parameters do we need?

This already reduces the number of relevant parameters by a large amount, a summary is given below:

- $\tan \beta = v_u/v_d$, the ratio of the VEVs in the two-Higgs doublet model
- $m_A^2 = 2m_{12}^2/\sin 2\beta$, the mass of the pseudoscalar Higgs boson
- M_1 , M_2 and M_3 , the bino, wino and gluino masses
- μ , the Higgs-higgsino mass parameter

Which Parameters do we need?

This already reduces the number of relevant parameters by a large amount, a summary is given below:

- $\tan \beta = v_u/v_d$, the ratio of the VEVs in the two-Higgs doublet model
- $m_A^2 = 2m_{12}^2/\sin 2\beta$, the mass of the pseudoscalar Higgs boson
- M_1 , M_2 and M_3 , the bino, wino and gluino masses
- μ , the Higgs-higgsino mass parameter
- $m_{\tilde{q}}$, $m_{\tilde{u}_R}$, $m_{\tilde{d}_R}$, $m_{\tilde{\ell}}$ and $m_{\tilde{e}_R}$, the first/second generation sfermion masses

Which Parameters do we need?

This already reduces the number of relevant parameters by a large amount, a summary is given below:

- $\tan \beta = v_u/v_d$, the ratio of the VEVs in the two-Higgs doublet model
- $m_A^2 = 2m_{12}^2/\sin 2\beta$, the mass of the pseudoscalar Higgs boson
- M_1 , M_2 and M_3 , the bino, wino and gluino masses
- μ , the Higgs-higgsino mass parameter
- $m_{\tilde{q}}$, $m_{\tilde{u}_R}$, $m_{\tilde{d}_R}$, $m_{\tilde{\ell}}$ and $m_{\tilde{e}_R}$, the first/second generation sfermion masses
- $m_{\tilde{Q}}$, $m_{\tilde{t}_R}$, $m_{\tilde{b}_R}$, $m_{\tilde{L}}$ and $m_{\tilde{\tau}_R}$, the third generation sfermion masses

Which Parameters do we need?

This already reduces the number of relevant parameters by a large amount, a summary is given below:

- $\tan \beta = v_u/v_d$, the ratio of the VEVs in the two-Higgs doublet model
- $m_A^2 = 2m_{12}^2/\sin 2\beta$, the mass of the pseudoscalar Higgs boson
- M_1 , M_2 and M_3 , the bino, wino and gluino masses
- μ , the Higgs-higgsino mass parameter
- $m_{\tilde{q}}$, $m_{\tilde{u}_R}$, $m_{\tilde{d}_R}$, $m_{\tilde{\ell}}$ and $m_{\tilde{e}_R}$, the first/second generation sfermion masses
- $m_{\tilde{Q}}$, $m_{\tilde{t}_R}$, $m_{\tilde{b}_R}$, $m_{\tilde{L}}$ and $m_{\tilde{\tau}_R}$, the third generation sfermion masses
- A^t , A^b and A^τ , the third generation trilinear couplings

In total we are left with 19 additional parameters in the pMSSM.

What about SUSY-breaking?

- The MSSM-124 fails to explain the fundamental origin of the SUSY-breaking parameters.
- **Last talks:** Gauge- and gravity-mediated SUSY breaking may provide a way around!
- The phenomenon of **gauge coupling unification** hints at a simpler structure of the MSSM at high energies.





Figure: Gravity-mediated SUSY-breaking in the hidden sector⁴.

⁴Figure inspired by the visualization in: https://www.thphys.uni-heidelberg.de/~plehn/includes/bad_honnef_12/kribs_2.pdf (13.02.15)

Outline

- 1 Overview
- 2 Phenomenology of the MSSM
- 3 A Top-Down Approach to Low-Energy Phenomenology: mSUGRA**
- 4 Grand Unification and SUSY
- 5 Going Beyond the MSSM
- 6 Summary and Outlook

The mSUGRA Framework I



The mSUGRA framework II

- Want to naturally include **gravity**, or more precisely **local SUGRA**.
- SUGRA is broken at some high energy scale, usually at around $M_{\text{Planck}} \sim 10^{19} \text{ GeV}$.
- Assume **gravity-mediated SUSY breaking**, i. e. $m_{\text{soft}} \sim \frac{\langle F \rangle}{M_{\text{Planck}}}$

The general idea is now to assume a “**minimal**” **normalization** of the kinetic terms in the SUGRA Lagrangian, i. e.

$$\mathcal{L}_{\text{SUGRA}} \supset -\frac{1}{M_{\text{Planck}}} F \left(\frac{1}{2} \alpha \lambda \lambda + \frac{1}{6} \beta \phi \phi \phi + \frac{1}{2} \gamma \phi \phi \right) + \text{h.c.} - \frac{1}{M_{\text{Planck}}^2} F F^* \delta \phi \phi^*$$

Comparison with initial $\mathcal{L}_{\text{soft}} \implies$ Simple relations for the input parameters!

Finding the correct initial Conditions

The relatively simple form of the kinetic terms allows to simplify the structure of the relevant couplings at the initial scale as follows:

- scalar squared-masses are **flavor-diagonal** and **universal**:

$$m_q^2(M_X) = m_u^2(M_X) = m_d^2(M_X) = m_0^2 \mathbb{1}$$

$$m_\ell^2(M_X) = m_e^2(M_X) = m_0^2 \mathbb{1}$$

$$m_1^2(M_X) = m_2^2(M_X) = m_0^2$$

- the same is true for the A -parameters:

$$A^U(M_X) = A^D(M_X) = A^E(M_X) = A_0^2 \mathbb{1}$$

- Additionally we assume unification of the (tree level) gaugino masses:

$$M_1(M_X) = M_2(M_X) = M_3(M_X) = m_{1/2}$$

Renormalization Group Equations

This allows us to compute the low-energy MSSM parameters with the help of standard **renormalization group techniques**!

Reminder: The **beta function** $\beta(g) = \frac{\partial g}{\partial \ln M}$ is the rate of change of the renormalized coupling at the scale M , where the bare coupling is fixed [17].

For example, we get the following result for the low-energy gaugino mass parameters:

$$M_i = \frac{\alpha_i(M_Z)}{\alpha_{\text{GUT}}} m_{1/2} \longrightarrow M_3(M_Z) = \frac{\alpha_3(M_Z)}{\alpha_2(M_Z)} M_2(M_Z) = \frac{\alpha_3(M_Z)}{\alpha_1(M_Z)} M_1(M_Z)$$

This results in the well known relation:

$$M_1 = \frac{5}{3} \tan^2 \theta_W M_2 \sim \frac{1}{2} M_2$$

Solution of the RGEs for mSUGRA




Figure: Running of the mSUGRA parameters⁵.

⁵Figure taken from: <https://www.physi.uni-heidelberg.de/~uwer/lectures/ParticlePhysics/Vorlesung/Lect-10b.pdf> (10.02.21)

Parameter Count in the mSUGRA Framework

This means in total we are left with only the following **four continuous** and **one discrete** free parameters in the mSUGRA model:

- $\tan\beta$, the ratio of the VEVs in the two-Higgs doublet model
- $m_{1/2}$, the universal gaugino mass
- m_0 , the universal scalar (sfermion/Higgs) mass
- A_0 , the universal trilinear coupling
- $\text{sign}(\mu)$, the sign of the Higgs-higgsino mass parameter

The relations for $\tan\beta$ and $|\mu|$ come from the **two minimum conditions** for the Higgs potential.

Remark: Additional requirements such as the unification of the top, bottom and tau Yukawa couplings at the GUT scale further restricts the possible values of $\tan\beta$ and A_0 .

Map of the mSUGRA Parameter Space




Figure: Map of the mSUGRA parameter space for different values of the universal mass parameters, slightly adapted from [10].

Outline

- 1 Overview
- 2 Phenomenology of the MSSM
- 3 A Top-Down Approach to Low-Energy Phenomenology: mSUGRA
- 4 Grand Unification and SUSY**
- 5 Going Beyond the MSSM
- 6 Summary and Outlook

Gauge Couplings in the Standard Model

The Standard Model is described as the most general renormalizable field theory with gauge group

$$\mathcal{G}_{\text{SM}} = \text{SU}(3) \times \text{SU}(2) \times \text{U}(1),$$

with associated gauge couplings α_3 , α_2 and α_1 , three generations of fermions and a scalar [4].

- The couplings are larger for the larger component of the gauge group, i. e.

$$\alpha_3(m_Z) > \alpha_2(m_Z) > \alpha_1(m_Z)$$

- Interesting observation: Values of the running couplings come close together at some high energy scale $\Lambda_{\text{GUT}} \sim 10^{16} \text{ GeV}$ (cf. next slide).
- **GEORGI-GLASHOW** [13]: Embed \mathcal{G}_{SM} in larger gauge group, i. e. $\text{SU}(5) \implies \text{GUT?}$

One Loop Running of the SM Gauge Couplings I





Figure: Running of the (inverse) SM gauge couplings, plot inspired by [15].

One Loop Running of the SM Gauge Couplings II

- In the Georgi-Glashow model, we need $-\mu^2 \sim -(100 \text{ GeV})^2$ to reproduce the correct W and Z masses \Rightarrow Gauge hierarchy problem! [17]
- SUSY provides way out: If SUSY breaking works such that the mass differences between the superpartners are large enough, one can reproduce the correct Higgs mass \Rightarrow superpartners influence the running of the (MS)SM gauge couplings (cf. next slide)



- In the end it remains a complicated fine tuning task!

Comparison: SM vs. MSSM





Figure: Running of the (inverse) gauge couplings in the SM and the MSSM, plots inspired by [\[15\]](#).

Outline

- 1 Overview
- 2 Phenomenology of the MSSM
- 3 A Top-Down Approach to Low-Energy Phenomenology: mSUGRA
- 4 Grand Unification and SUSY
- 5 Going Beyond the MSSM**
- 6 Summary and Outlook

Are all Problems solved now?

The MSSM as introduced in the first part of our talk still fails to answer some of the key questions:

- How do we explain the value of the μ parameter in the MSSM?
- What about the “total” particle content?
- Is there a “natural” way to implement SUSY-breaking?
- What is $\mathcal{G}_{\text{MSSM}}$?
- and (many) more ...

To conclude our talk we want to have a look at some ideas concerning the first mentioned problem, the value of the μ parameter.

Different Approaches to explain the Value of the μ Parameter

- **General Problem:** μ is a SUSY-preserving parameter, but from phenomenology we know that it must be of the order of the SUSY-breaking scale.
- “Natural” solution: Symmetry enforcing $\mu = 0$, and a small SUSY-breaking parameter that generates a value of μ which is not parametrically larger than the SUSY-breaking scale.

In proposed extensions of the MSSM, some other approaches have been presented, for example:

Different Approaches to explain the Value of the μ Parameter

- **General Problem:** μ is a SUSY-preserving parameter, but from phenomenology we know that it must be of the order of the SUSY-breaking scale.
- “Natural” solution: Symmetry enforcing $\mu = 0$, and a small SUSY-breaking parameter that generates a value of μ which is not parametrically larger than the SUSY-breaking scale.

In proposed extensions of the MSSM, some other approaches have been presented, for example:

- 1 Replace μ by the VEV of a new $SU(3) \times SU(2) \times U(1)$ scalar singlet \Rightarrow NMSSM

Different Approaches to explain the Value of the μ Parameter

- **General Problem:** μ is a SUSY-preserving parameter, but from phenomenology we know that it must be of the order of the SUSY-breaking scale.
- “Natural” solution: Symmetry enforcing $\mu = 0$, and a small SUSY-breaking parameter that generates a value of μ which is not parametrically larger than the SUSY-breaking scale.

In proposed extensions of the MSSM, some other approaches have been presented, for example:

- ① Replace μ by the VEV of a new $SU(3) \times SU(2) \times U(1)$ scalar singlet \Rightarrow NMSSM
- ② Add a new broken $U(1)$ gauge symmetry to the NMSSM \Rightarrow USSM

Different Approaches to explain the Value of the μ Parameter

- **General Problem:** μ is a SUSY-preserving parameter, but from phenomenology we know that it must be of the order of the SUSY-breaking scale.
- “Natural” solution: Symmetry enforcing $\mu = 0$, and a small SUSY-breaking parameter that generates a value of μ which is not parametrically larger than the SUSY-breaking scale.

In proposed extensions of the MSSM, some other approaches have been presented, for example:

- 1 Replace μ by the VEV of a new $SU(3) \times SU(2) \times U(1)$ scalar singlet \Rightarrow NMSSM
- 2 Add a new broken $U(1)$ gauge symmetry to the NMSSM \Rightarrow USSM
- 3 Allow all possible renormalizable terms in the superpotential \Rightarrow new mass terms (GNMSSM)

Different Approaches to explain the Value of the μ Parameter

- **General Problem:** μ is a SUSY-preserving parameter, but from phenomenology we know that it must be of the order of the SUSY-breaking scale.
- “Natural” solution: Symmetry enforcing $\mu = 0$, and a small SUSY-breaking parameter that generates a value of μ which is not parametrically larger than the SUSY-breaking scale.

In proposed extensions of the MSSM, some other approaches have been presented, for example:

- ① Replace μ by the VEV of a new $SU(3) \times SU(2) \times U(1)$ scalar singlet \Rightarrow NMSSM
- ② Add a new broken $U(1)$ gauge symmetry to the NMSSM \Rightarrow USSM
- ③ Allow all possible renormalizable terms in the superpotential \Rightarrow new mass terms (GNMSSM)
- ④ Possible connection to the strong CP problem? \Rightarrow PQ symmetry \Rightarrow Axion Physics?

Different Approaches to explain the Value of the μ Parameter

- **General Problem:** μ is a SUSY-preserving parameter, but from phenomenology we know that it must be of the order of the SUSY-breaking scale.
- “Natural” solution: Symmetry enforcing $\mu = 0$, and a small SUSY-breaking parameter that generates a value of μ which is not parametrically larger than the SUSY-breaking scale.

In proposed extensions of the MSSM, some other approaches have been presented, for example:

- 1 Replace μ by the VEV of a new $SU(3) \times SU(2) \times U(1)$ scalar singlet \Rightarrow NMSSM
- 2 Add a new broken $U(1)$ gauge symmetry to the NMSSM \Rightarrow USSM
- 3 Allow all possible renormalizable terms in the superpotential \Rightarrow new mass terms (GNMSSM)
- 4 Possible connection to the strong CP problem? \Rightarrow PQ symmetry \Rightarrow Axion Physics?
- 5 Higher-dimensional Higgs multiplets?

Outline

- 1 Overview
- 2 Phenomenology of the MSSM
- 3 A Top-Down Approach to Low-Energy Phenomenology: mSUGRA
- 4 Grand Unification and SUSY
- 5 Going Beyond the MSSM
- 6 Summary and Outlook

What have we learned today?

- The SM can be promoted to the MSSM using the concepts and methods introduced in the scope of this seminar.
- In addition to the 19 parameters of the SM we get $\mathcal{O}(100)$ new ones, which complicates the experimental access to the MSSM a lot!
- Phenomenological models of the MSSM allow for a drastic reduction of the number of independent parameters that have to be measured experimentally.
- SUSY may lead to a better understanding of the connection between Gravity and the SM. Additionally SUGRA helps us to further reduce the number of dof's in the MSSM.
- In the MSSM the unification of the SM gauge couplings at some high-energy scale M_X can be realized due to the effect of loop-corrections arising from the additional superpartners.
- There are still plenty of problems where the MSSM fails to explain the experimental results.
⇒ The MSSM is not the “ultimate” model to describe our Universe.

References Part I

- [1] DJH Chung, LL Everett, GL Kane, SF King, J Lykken, and Lian-Tao Wang.
The soft supersymmetry-breaking Lagrangian: Theory and applications.
Physics Reports 407.1-3 (2005), pp. 1–203.
- [2] Patrick Draper and Heidi Rzehak.
A review of Higgs mass calculations in supersymmetric models.
Physics Reports 619 (2016), pp. 1–24.
- [3] Howard E Haber.
The status of the minimal supersymmetric standard model and beyond.
Nuclear Physics B-Proceedings Supplements 62.1-3 (1998), pp. 469–484.
- [4] Arthur Hebecker.
Lectures on Beyond the Standard Model and the String Theory Landscape.
Heidelberg University, 2020 (cit. on p. 25).
- [5] Stephen P. Martin.
A Supersymmetry Primer.
Advanced Series on Directions in High Energy Physics (1998), pp. 1–98.
- [6] Hans - Peter Nilles.
Supersymmetry, Supergravity and Particle Physics.
Phys. Rept. 110 (1984), pp. 1–162.
- [7] Adrian Signer.
ABC of SUSY.
Journal of Physics G: Nuclear and Particle Physics 36.7 (2009), p. 073002.
- [8] P.A. Zyla et al.
Review of Particle Physics.
PTEP 2020.8 (2020), p. 083C01.

References Part II

- [9] Ugo Amaldi, Wim de Boer, and Hermann Fürstenau.
Comparison of grand unified theories with electroweak and strong coupling constants measured at LEP.
Phys. Lett. B 260.CERN-PPE-91-44. IEKP-KA-91-01 (1991), 447–455. 20 p.
- [10] G. L. Bayatian et al.
CMS technical design report, volume II: Physics performance.
J. Phys. G 34.6 (2007), pp. 995–1579 (cit. on p. 23).
- [11] A. Djouadi et al.
The Minimal supersymmetric standard model: Group summary report.
GDR (Groupement De Recherche) - Supersymetrie. 1998. arXiv: [hep-ph/9901246](#) (cit. on p. 6).
- [12] Willy Fischler, Sonia Paban, and Scott D. Thomas.
Bounds on microscopic physics from P and T violation in atoms and molecules.
Phys. Lett. B 289 (1992), pp. 373–380. arXiv: [hep-ph/9205233](#) (cit. on p. 5).
- [13] H. Georgi and S. L. Glashow.
Unity of All Elementary Particle Forces.
Phys. Rev. Lett. 32 (1974), pp. 438–441 (cit. on p. 25).
- [3] Howard E Haber.
The status of the minimal supersymmetric standard model and beyond.
Nuclear Physics B-Proceedings Supplements 62.1-3 (1998), pp. 469–484.
- [4] Arthur Hebecker.
Lectures on Beyond the Standard Model and the String Theory Landscape.
Heidelberg University, 2020 (cit. on p. 25).
- [14] G.L. Kane.
Perspectives on Supersymmetry II.
Advanced series on directions in high energy physics. World Scientific, 2010.
- [15] D. I. Kazakov.
Beyond the Standard Model: In Search of Supersymmetry.
2000 European School of High-Energy Physics. 2000, pp. 125–199. arXiv: [hep-ph/0012288](#) (cit. on pp. 26, 28).
- [16] Stephen P. Martin and Michael T. Vaughn.
Two loop renormalization group equations for soft supersymmetry breaking couplings.
Phys. Rev. D 50 (1994). [Erratum: *Phys.Rev.D* 78, 039903 (2008)], p. 2282. arXiv: [hep-ph/9311340](#).
- [17] Michael E. Peskin and Daniel V. Schroeder.
An Introduction to quantum field theory.
Reading, USA: Addison-Wesley, 1995 (cit. on pp. 20, 27).
- [18] M. Tanabashi et al.
Review of Particle Physics 2018 and 2019 Update.
Phys. Rev. D 98.3 (2018), p. 030001.