

Matt study, Nuclear transparency

Mathieu Ouillon

2024
September

0.1 Strategy

0.1.1 Carbon target

Carbon runs : 018339, 018340, 018341, 018342, 018343, 018344, 018346, 018440, 018441, 018442, 018443, 018444, 018445, 018475, 018498, 018524, 018756, 018850

path: /cache/hallb/scratch/rg-d/production/Bspot/v5dstCxC/dst/recon/

- Find event with a trigger electron(`REC::Particle::pid == 11` and `status<0`), at least a π^+ and at least a π^- .
- Apply cut on electron: $-3 < \chi_{pid}^2 < 3$ and $-12 < v_z < 5$
- Find all π^+ in event: `REC::Particle::pid == 211`
- Apply cut on π^+ : $-10 < \chi_{pid}^2 < 10$
- Find all π^- in event: `REC::Particle::pid == -211`
- Apply cut on π^- : $-10 < \chi_{pid}^2 < 10$
- Find all combinaison of π^+ and π^-
- Cut to select reaction :
 - $W = (p_i + \gamma^*)^2 > 2GeV$, with $p_i = (0, 0, 0, M_p)$, $M_p = 0.938GeV$
 - $z_h = \frac{E_{\rho^0}}{v} > 0.9$
 - $0.1 < -t = (\gamma^* - p_{\rho^0}) < 0.5GeV^2$
 - $l_c \leq 0.5fm$
- Fill invariant mass of ρ^0 for the Q^2 bins : $1 \leq Q^2 < 2$, $2 \leq Q^2 < 2.5$, $2.5 \leq Q^2 < 3$, $3 \leq Q^2 < 3.5$, $3.5 \leq Q^2 < 4.5$, $4.5 \leq Q^2 < 6$
- Fit the distribution with a Breit–Wigner and a 3rd order polynom :

$$BW(x; x_0, \Gamma, \alpha) = \alpha \times \frac{1}{\pi} \times \frac{\frac{1}{2}\Gamma}{(x - x_0)^2 + \frac{1}{2}\Gamma} \quad (1)$$

$$pol3(x; a, b, c, d) = a + b \times x + c \times x^2 + d \times x^3 \quad (2)$$






where x_0 is the location parameter, specifying the location of the peak of the distribution, γ is full width at half maximum (FWHM).

0.2 Carbon data






0.2.1 Numbers

- Total number of event : 354888652
- Number of event with a trigger electron (`REC::Particle::pid == 11` and `status<0`) and at least a π^+ and at least a π^- : 22585713; ratio between this one and the previous number : 0.0636417
- Number of event with a good electron : 19319798; ratio : 0.855399
- Number of π^+ after electron cuts: 26473357
- Number of good π^+ after electron and π^+ cuts : 17525249; ratio : 0.661996
- Number of bad π^+ after electron and opposite π^+ cuts: 8948108; ratio : 0.338004
- Number of π^- after electron cuts: 24167298
- Number of good π^- after electron and π^- cuts: 19747026; ratio : 0.817097
- Number of bad π^- after electron and opposite π^- cuts: 4420272; ratio : 0.182903
- Number of ρ^0 : 19678575;
- Number of ρ^0 that pass the W cut : 19034880; ratio : 0.96729
- Number of ρ^0 that pass the W and z_h cuts : 343967; ratio : 0.0180704
- Number of ρ^0 that pass the W , z_h and t cuts : 158735; ratio : 0.461483
- Number of ρ^0 that pass the W , z_h , t and l_c cuts : 9719; ratio : 0.0612278






0.2.2 Electron









0.2.3 π^+




0.2.4 π^-



0.2.5 Event selection



0.2.6 ρ^0 invariant mass



0.2.7 LD2

LD2 runs : 018419, 018421, 018424, 018427, 018428, 018429, 018431, 018432, 018433, 018439, 018528, 018559, 018644, 018656, 018851, 018873, 019021, 019058

path: /cache/hallb/scratch/rg-d/production/Bspot/v5dstLD2/dst/recon/

- Select the trigger electron: `REC::Particle::pid == 11` and `status<0`
- Apply cut on electron: $-3 < \chi^2_{pid} < 3$ and $-12 < v_z < 5$

- Find all π^+ in event: `REC::Particle::pid == 211`
- Apply cut on π^+ : $-10 < \chi_{pid}^2 < 10$
- Find all π^- in event: `REC::Particle::pid == -211`
- Apply cut on π^- : $-10 < \chi_{pid}^2 < 10$
- Find all combinaison of π^+ and π^-
- Cut to select reaction :
 - $W = (p_i + \gamma^*)^2 > 2GeV$
 - $z_h = \frac{E_{\rho^0}}{v} > 0.9$
 - $0.1 < -t < 0.5GeV^2$
 - $l_c \leq 0.5fm$