

INF5620 - First compulsory project

Arnfinn Mihle Paulsrud

October 9, 2012

1 mathematical problem

$$\frac{\partial^2 u}{\partial t^2} + b \frac{\partial u}{\partial t} = \frac{\partial}{\partial x} \left(q(x, y) \frac{\partial u}{\partial x} \right) + \frac{\partial}{\partial y} \left(q(x, y) \frac{\partial u}{\partial y} \right) + f(x, y, t) \quad (1)$$

The way we solve this numerically is by discretization

$$\frac{\partial^2 u}{\partial t^2} \rightarrow \frac{u_{i,j}^{n+1} - 2u_{i,j}^n + u_{i,j}^{n-1}}{\Delta t^2} \quad (2)$$

$$b \frac{\partial u}{\partial t} \rightarrow b \frac{u_{i,j}^{n+1} - u_{i,j}^{n-1}}{2\Delta t} \quad (3)$$

$$\begin{aligned} \frac{\partial}{\partial x} \left(q(x, y) \frac{\partial u}{\partial x} \right) &\rightarrow \frac{(q(x, y) \frac{\partial u}{\partial x})_{i+\frac{1}{2},j} - (q(x, y) \frac{\partial u}{\partial x})_{i-\frac{1}{2},j}}{\Delta x} \\ &= \frac{q(x, y)_{i+\frac{1}{2},j}^n \frac{u_{i+1,j}^n - u_{i,j}^n}{\Delta x} - q(x, y)_{i-\frac{1}{2},j}^n \frac{u_{i,j}^n - u_{i-1,j}^n}{\Delta x}}{\Delta x} \\ &= \frac{1}{\Delta x^2} \left(q(x, y)_{i+\frac{1}{2},j}^n (u_{i+1,j}^n - u_{i,j}^n) - q(x, y)_{i-\frac{1}{2},j}^n (u_{i,j}^n - u_{i-1,j}^n) \right) \end{aligned} \quad (4)$$

We can find a solution for $q(x, y)_{i+\frac{1}{2},j}^n$ and $q(x, y)_{i-\frac{1}{2},j}^n$ by taking the average between $q(x, y)_{i+1,j}^n$ and $q(x, y)_{i,j}^n$ and between $q(x, y)_{i,j}^n$ and $q(x, y)_{i-1,j}^n$.

$$q(x, y)_{i+\frac{1}{2},j}^n \approx \frac{q(x, y)_{i+1,j}^n + q(x, y)_{i,j}^n}{2} \quad (5a)$$

and

$$q(x, y)_{i-\frac{1}{2},j}^n \approx \frac{q(x, y)_{i,j}^n + q(x, y)_{i-1,j}^n}{2} \quad (6a)$$

This gives us

$$\begin{aligned} \frac{\partial}{\partial x} \left(q(x, y) \frac{\partial u}{\partial x} \right) &\rightarrow \frac{1}{\Delta x^2} \left(\frac{q(x, y)_{i+1,j}^n + q(x, y)_{i,j}^n}{2} (u_{i+1,j}^n - u_{i,j}^n) - \frac{q(x, y)_{i,j}^n + q(x, y)_{i-1,j}^n}{2} (u_{i,j}^n - u_{i-1,j}^n) \right) \\ &= \frac{1}{2\Delta x^2} \left((q(x, y)_{i+1,j}^n + q(x, y)_{i,j}^n) (u_{i+1,j}^n - u_{i,j}^n) - (q(x, y)_{i,j}^n + q(x, y)_{i-1,j}^n) (u_{i,j}^n - u_{i-1,j}^n) \right) \end{aligned} \quad (7)$$

By doing the same for $\left(q(x, y) \frac{\partial u}{\partial y} \right)$, we get

$$\begin{aligned}
& \frac{\partial}{\partial y} \left(q(x, y) \frac{\partial u}{\partial y} \right) \\
& \rightarrow \frac{1}{2\Delta y^2} \left((q(x, y)_{i,j+1}^n + q(x, y)_{i,j}^n) (u_{i,j+1}^n - u_{i,j}^n) - (q(x, y)_{i,j}^n + q(x, y)_{i,j-1}^n) (u_{i,j}^n - u_{i,j-1}^n) \right)
\end{aligned} \tag{8}$$

The last part of eq.(1) becomes

$$f(x, y, t) \rightarrow f(x, y, t)_{i,j}^n \tag{9}$$

If we now combine our discretized equations we get a numerically solvable wave equation

$$\begin{aligned}
& \frac{u_{i,j}^{n+1} - 2u_{i,j}^n + u_{i,j}^{n-1}}{\Delta t^2} + b \frac{u_{i,j}^{n+1} - u_{i,j}^{n-1}}{2\Delta t} = \\
& \frac{1}{2\Delta x^2} \left((q(x, y)_{i+1,j}^n + q(x, y)_{i,j}^n) (u_{i+1,j}^n - u_{i,j}^n) - (q(x, y)_{i,j}^n + q(x, y)_{i-1,j}^n) (u_{i,j}^n - u_{i-1,j}^n) \right) \\
& + \frac{1}{2\Delta y^2} \left((q(x, y)_{i,j+1}^n + q(x, y)_{i,j}^n) (u_{i,j+1}^n - u_{i,j}^n) - (q(x, y)_{i,j}^n + q(x, y)_{i,j-1}^n) (u_{i,j}^n - u_{i,j-1}^n) \right) \\
& + f(x, y, t)_{i,j}^n
\end{aligned}$$

We would now like to solve this equation for $u_{i,j}^{n+1}$

$$\begin{aligned}
& u_{i,j}^{n+1} - 2u_{i,j}^n + u_{i,j}^{n-1} + \frac{b\Delta t}{2} (u_{i,j}^{n+1} - u_{i,j}^{n-1}) = \\
& \frac{\Delta t^2}{2\Delta x^2} \left((q(x, y)_{i+1,j}^n + q(x, y)_{i,j}^n) (u_{i+1,j}^n - u_{i,j}^n) - (q(x, y)_{i,j}^n + q(x, y)_{i-1,j}^n) (u_{i,j}^n - u_{i-1,j}^n) \right) \\
& + \frac{\Delta t^2}{2\Delta y^2} \left((q(x, y)_{i,j+1}^n + q(x, y)_{i,j}^n) (u_{i,j+1}^n - u_{i,j}^n) - (q(x, y)_{i,j}^n + q(x, y)_{i,j-1}^n) (u_{i,j}^n - u_{i,j-1}^n) \right) \\
& + \Delta t^2 f(x, y, t)_{i,j}^n
\end{aligned}$$

$$\begin{aligned}
& u_{i,j}^{n+1} \left(1 + \frac{b\Delta t}{2} \right) = \\
& \frac{\Delta t^2}{2\Delta x^2} \left((q(x, y)_{i+1,j}^n + q(x, y)_{i,j}^n) (u_{i+1,j}^n - u_{i,j}^n) - (q(x, y)_{i,j}^n + q(x, y)_{i-1,j}^n) (u_{i,j}^n - u_{i-1,j}^n) \right) \\
& + \frac{\Delta t^2}{2\Delta y^2} \left((q(x, y)_{i,j+1}^n + q(x, y)_{i,j}^n) (u_{i,j+1}^n - u_{i,j}^n) - (q(x, y)_{i,j}^n + q(x, y)_{i,j-1}^n) (u_{i,j}^n - u_{i,j-1}^n) \right) \\
& + \Delta t^2 f(x, y, t)_{i,j}^n + 2u_{i,j}^n + u_{i,j}^{n-1} \left(\frac{b\Delta t}{2} - 1 \right)
\end{aligned}$$

We end up with the discretized equation

$$\begin{aligned}
& u_{i,j}^{n+1} = \\
& \frac{\Delta t^2}{2\Delta x^2 \left(1 + \frac{b\Delta t}{2} \right)} \left((q(x, y)_{i+1,j}^n + q(x, y)_{i,j}^n) (u_{i+1,j}^n - u_{i,j}^n) - (q(x, y)_{i,j}^n + q(x, y)_{i-1,j}^n) (u_{i,j}^n - u_{i-1,j}^n) \right) \\
& + \frac{\Delta t^2}{2\Delta y^2 \left(1 + \frac{b\Delta t}{2} \right)} \left((q(x, y)_{i,j+1}^n + q(x, y)_{i,j}^n) (u_{i,j+1}^n - u_{i,j}^n) - (q(x, y)_{i,j}^n + q(x, y)_{i,j-1}^n) (u_{i,j}^n - u_{i,j-1}^n) \right) \\
& + \frac{\Delta t^2}{1 + \frac{b\Delta t}{2}} f(x, y, t)_{i,j}^n + 2u_{i,j}^n + u_{i,j}^{n-1} \frac{1 + \frac{b\Delta t}{2}}{\frac{b\Delta t}{2} - 1}
\end{aligned} \tag{10}$$

2 Boundary condition

If we take a look at our numerical equation, we can see that we'll get a problem when we get to the boundary $i = 0$, $i = L_x$, $j = 0$ and $j = L_y$. We have been given the boundary condition $\frac{\partial u}{\partial \eta} = 0$

$$\frac{\partial u}{\partial \eta} = 0 \rightarrow [D_{2x}u]_{i,j}^n = 0 \quad (11)$$

When we are on the boundary $x = 0 \Rightarrow i = 0$ we get

$$[D_{2x}u]_{i,j}^n = \frac{u_{i+1,j}^n - u_{i-1,j}^n}{2\Delta x} = 0 \quad (12)$$

$$\begin{aligned} \frac{u_{1,j}^n - u_{-1,j}^n}{2\Delta x} &= 0 \\ \rightarrow u_{1,j}^n &= u_{-1,j}^n \end{aligned}$$

When we are on the boundary $y = 0 \Rightarrow j = 0$ we get

$$[D_{2x}u]_{i,j}^n = \frac{u_{i+1,j}^n - u_{i-1,j}^n}{2\Delta x} = 0 \quad (13)$$

$$\begin{aligned} \frac{u_{1,j}^n - u_{-1,j}^n}{2\Delta x} &= 0 \\ \rightarrow u_{1,j}^n &= u_{-1,j}^n \end{aligned}$$

3 Initial condition

If we take a look at eq.(??) we can see that we are going to get a problem with $u_{i,j}^{n-1}$ in our first time step. We can solve this by looking at our initial condition. If we assume that the velocity at $t = 0$ is zero, $u'(x, y, 0) = 0$ then we get

$$\frac{\partial u}{\partial t} = 0 \rightarrow [D_t u]_{i,j}^n = 0 \quad (14)$$

$$\frac{u_{i,j}^{n+1} - u_{i,j}^{n-1}}{2\Delta t} = 0 \quad (15)$$

For $t = 0 \rightarrow n = 0$ we get

$$\frac{u_{i,j}^1 - u_{i,j}^{-1}}{2\Delta t} = 0 \quad (16)$$

$$u_{i,j}^1 = u_{i,j}^{-1} \quad (17)$$

This solves our problem. If we now use $n = 0$ in eq.(??) we get

$$\begin{aligned} u_{i,j}^1 \left(1 + \frac{b\Delta t}{2} \right) &= \\ &= \frac{\Delta t^2}{2\Delta x^2} \left((q(x, y)_{i+1,j}^0 + q(x, y)_{i,j}^0) (u_{i+1,j}^0 - u_{i,j}^0) - (q(x, y)_{i,j}^0 + q(x, y)_{i-1,j}^0) (u_{i,j}^0 - u_{i-1,j}^0) \right) \\ &+ \frac{\Delta t^2}{2\Delta y^2} \left((q(x, y)_{i,j+1}^0 + q(x, y)_{i,j}^0) (u_{i,j+1}^0 - u_{i,j}^0) - (q(x, y)_{i,j}^0 + q(x, y)_{i,j-1}^0) (u_{i,j}^0 - u_{i,j-1}^0) \right) \\ &+ \Delta t^2 f(x, y, t)_{i,j}^0 + 2u_{i,j}^0 + u_{i,j}^{-1} \left(\frac{b\Delta t}{2} - 1 \right) \end{aligned}$$

If we now use $u_{i,j}^1 = u_{i,j}^{-1}$ we get

$$\begin{aligned}
u_{i,j}^1 = & \frac{\Delta t^2}{4\Delta x^2} ((q(x,y)_{i+1,j}^0 + q(x,y)_{i,j}^0) (u_{i+1,j}^0 - u_{i,j}^0) - (q(x,y)_{i,j}^0 + q(x,y)_{i-1,j}^0) (u_{i,j}^0 - u_{i-1,j}^0)) \\
& + \frac{\Delta t^2}{4\Delta y^2} ((q(x,y)_{i,j+1}^0 + q(x,y)_{i,j}^0) (u_{i,j+1}^0 - u_{i,j}^0) - (q(x,y)_{i,j}^0 + q(x,y)_{i,j-1}^0) (u_{i,j}^0 - u_{i,j-1}^0)) \\
& + \frac{\Delta t^2}{2} f(x,y,t)_{i,j}^0 + u_{i,j}^0
\end{aligned}$$

4 Stability criterion