

Interior Point Algorithms III

Yinyu Ye

Department of Management Science and Engineering

Stanford University

Stanford, CA 94305, U.S.A.

<http://www.stanford.edu/~yyye>

(LY: Chapter 5)

Termination with Exact Optimizers

- The first is a “cross-over” procedure to find a **basic feasible solution** (BFS, corner point) whose objective value is at least as good as the current interior point. Let $A, \mathbf{b}, \mathbf{c}$ be integers and L be their **bit length**, and let a second best BFS solution be \mathbf{x}^{2nd} and the **optimal objective value** be z^* . Then

$$\mathbf{c}^T \mathbf{x}^{2nd} - z^* > 2^{-L}.$$

Thus, one can terminate interior-point algorithm when

$$\mathbf{c}^T \mathbf{x}^k - \mathbf{b}^T \mathbf{y}^k \leq 2^{-L}.$$

- The second approach is to compute a **strictly complementary solution pair**. The method uses the primal-dual interior-point pair to identify the **strict complementarity partition** (P^*, Z^*) and then “purify or project” the primal interior solution onto the **primal optimal face** and the dual interior solution onto

the **dual optimal face**, based on the following theorem:

Theorem 1 *Given an interior solution \mathbf{x}^k and \mathbf{s}^k in the solution sequence generated by an interior-point algorithm, define*

$$P^k = \{j : x_j^k \geq s_j^k, \forall j\} \quad \text{and} \quad Z^k = \{1, \dots, n\} \setminus P^k.$$

Then, we have $P^k = P^$ whenever*

$$\mathbf{c}^T \mathbf{x}^k - \mathbf{b}^T \mathbf{y}^k \leq 2^{-L}.$$

Thus, the **worst-case iteration bound** for interior-point algorithms is $O(\sqrt{n}L)$ if the initial point pair $(\mathbf{x}^0)^T \mathbf{s}^0 \leq 2^L$.

Initialization

- Combining the primal and dual into a single **linear feasibility** problem, then applying LP algorithms to find a feasible point of the problem. Theoretically, this approach can retain the currently best complexity result.
- The **big M** method, i.e., add one or more artificial column(s) and/or row(s) and a huge penalty parameter M to force solutions to become feasible during the algorithm.
- **Phase I-then-Phase II method**, i.e., first try to find a feasible point (and possibly one for the dual problem), and then start to look for an optimal solution if the problem is feasible and bounded.
- **Combined Phase I-Phase II method**, i.e., approach feasibility and optimality simultaneously. To our knowledge, the “best” complexity of this approach is $O(n \log(R/\epsilon))$.

Homogeneous and Self-Dual Algorithm

- It solves the linear programming problem without any regularity assumption concerning the existence of **optimal, feasible, or interior feasible** solutions, while it retains the currently best complexity result
- It can start at any positive primal-dual pair, **feasible or infeasible**, near the central ray of the positive orthant (cone), and it does not use any big M penalty parameter or lower bound.
- Each iteration solves a system of linear equations whose dimension is almost the **same** as that solved in the standard (primal-dual) interior-point algorithms.
- If the LP problem has a solution, the algorithm generates a sequence that approaches **feasibility and optimality** simultaneously; if the problem is infeasible or unbounded, the algorithm will produce an **infeasibility certificate** for at least one of the primal and dual problems.

Primal-Dual Alternative Systems

A pair of LP has **two alternatives**

(Solvable)	$A\mathbf{x} - \mathbf{b} = \mathbf{0}$	(Infeasible)	$A\mathbf{x} = \mathbf{0}$
	$-A^T\mathbf{y} + \mathbf{c} \geq \mathbf{0},$		$-A^T\mathbf{y} \geq \mathbf{0},$
	$\mathbf{b}^T\mathbf{y} - \mathbf{c}^T\mathbf{x} = 0,$	or	$\mathbf{b}^T\mathbf{y} - \mathbf{c}^T\mathbf{x} > 0,$
	$\mathbf{y} \text{ free, } \mathbf{x} \geq \mathbf{0}$		$\mathbf{y} \text{ free, } \mathbf{x} \geq \mathbf{0}$

An Integrated Homogeneous System

The two alternative systems can be **homogenized** as one:

$$\begin{aligned} (HP) \quad & A\mathbf{x} - \mathbf{b}\tau = \mathbf{0} \\ & -A^T\mathbf{y} + \mathbf{c}\tau = \mathbf{s} \geq \mathbf{0}, \\ & \mathbf{b}^T\mathbf{y} - \mathbf{c}^T\mathbf{x} = \kappa \geq 0, \\ & \mathbf{y} \text{ free}, (\mathbf{x}; \tau) \geq \mathbf{0} \end{aligned}$$

where the **two alternatives** are

$$(\text{Solvable}) : (\tau > 0, \kappa = 0) \quad \text{or} \quad (\text{Infeasible}) : (\tau = 0, \kappa > 0)$$

The Homogeneous System is Self-Dual

$$\begin{array}{ll}
 (HP) & \begin{array}{l} A\mathbf{x} - \mathbf{b}\tau = \mathbf{0}, (\mathbf{y}') \\ -A^T\mathbf{y} + \mathbf{c}\tau = \mathbf{s} \geq \mathbf{0}, (\mathbf{x}') \\ \mathbf{b}^T\mathbf{y} - \mathbf{c}^T\mathbf{x} = \kappa \geq 0, (\tau') \\ \mathbf{y} \text{ free}, (\mathbf{x}; \tau) \geq \mathbf{0} \end{array} \\
 (HD) & \begin{array}{l} A\mathbf{x}' - \mathbf{b}\tau' = \mathbf{0}, \\ A^T\mathbf{y}' - \mathbf{c}\tau' \leq \mathbf{0}, \\ -\mathbf{b}^T\mathbf{y}' + \mathbf{c}^T\mathbf{x}' \leq 0, \\ \mathbf{y}' \text{ free}, (\mathbf{x}'; \tau') \geq \mathbf{0} \end{array}
 \end{array}$$

Theorem 2 System (HP) is feasible (e.g. all zeros) and any feasible solution $(\mathbf{y}, \mathbf{x}, \tau, \mathbf{s}, \kappa)$ is *self-complementary*:

$$\mathbf{x}^T \mathbf{s} + \tau \kappa = 0.$$

Furthermore, it has a *strictly* self-complementary feasible solution

$$\begin{pmatrix} \mathbf{x} + \mathbf{s} \\ \tau + \kappa \end{pmatrix} > \mathbf{0},$$

Let's Find Such a Feasible Solution

Given $\mathbf{x}^0 = \mathbf{e} > \mathbf{0}$, $\mathbf{s}^0 = \mathbf{e} > \mathbf{0}$, and $\mathbf{y}^0 = \mathbf{0}$, we formulate

$$\begin{aligned}
 (HSDP) \quad & \min \quad \theta \\
 \text{s.t.} \quad & A\mathbf{x} - \mathbf{b}\tau + \bar{\mathbf{b}}\theta = \mathbf{0}, \\
 & -A^T\mathbf{y} + \mathbf{c}\tau - \bar{\mathbf{c}}\theta \geq \mathbf{0}, \\
 & \mathbf{b}^T\mathbf{y} - \mathbf{c}^T\mathbf{x} + \bar{z}\theta \geq 0, \\
 & \mathbf{y} \text{ free}, \quad \mathbf{x} \geq \mathbf{0}, \quad \tau \geq 0, \quad \theta \text{ free},
 \end{aligned}$$

where

$$\bar{\mathbf{b}} = \mathbf{b} - A\mathbf{e}, \quad \bar{\mathbf{c}} = \mathbf{c} - \mathbf{e}, \quad \bar{z} = \mathbf{c}^T\mathbf{e} + 1.$$

But it may just give us the **all-zero solution**.

A HSD linear program

Let's try to add one more constraint to **prevent the all-zero solution**

$$\begin{aligned}
 (HSDP) \quad & \min && (n+1)\theta \\
 & \text{s.t.} && Ax - b\tau + \bar{b}\theta = 0, \\
 & && -A^T y + c\tau - \bar{c}\theta \geq 0, \\
 & && b^T y - c^T x + \bar{z}\theta \geq 0, \\
 & && -\bar{b}^T y + \bar{c}^T x - \bar{z}\tau = -(n+1), \\
 & && y \text{ free, } x \geq 0, \tau \geq 0, \theta \text{ free.}
 \end{aligned}$$

Note that the constraints of (HSDP) form a **skew-symmetric system** and the objective coefficient vector is the negative of the right-hand-side vector, so that it remains a **self-dual** linear program.

$(y = 0, x = e, \tau = 1, \theta = 1)$ is a **strictly** feasible point for (HSDP).

$$\begin{aligned}
 (HSDP) \quad & \min && (n+1)\theta \\
 \text{s.t.} \quad & & Ax & - \mathbf{b}\tau & + \bar{\mathbf{b}}\theta & = \mathbf{0}, \\
 & -A^T \mathbf{y} & & + \mathbf{c}\tau & - \bar{\mathbf{c}}\theta & = \mathbf{s} \geq \mathbf{0}, \\
 & \mathbf{b}^T \mathbf{y} & - \mathbf{c}^T \mathbf{x} & & + \bar{\mathbf{z}}\theta & = \kappa \geq 0, \\
 & -\bar{\mathbf{b}}^T \mathbf{y} & + \bar{\mathbf{c}}^T \mathbf{x} & - \bar{\mathbf{z}}\tau & & = -(n+1), \\
 & \mathbf{y} \text{ free, } & \mathbf{x} \geq \mathbf{0}, & \tau \geq 0, & \theta \text{ free.}
 \end{aligned}$$

Denote by \mathcal{F}_h the set of all points $(\mathbf{y}, \mathbf{x}, \tau, \theta, \mathbf{s}, \kappa)$ that are feasible for (HSDP). Denote by \mathcal{F}_h^0 the set of interior feasible points with $(\mathbf{x}, \tau, \mathbf{s}, \kappa) > \mathbf{0}$ in \mathcal{F}_h . By combining the constraints, we can derive the last (equality) constraint as

$$\mathbf{e}^T x + \mathbf{e}^T s + \tau + \kappa - (n+1)\theta = (n+1),$$

which serves indeed as a **normalizing constraint** for (HSDP) to prevent the all-zero solution.

Theorem 3 Consider problems (HSDP) and (HSDD).

i) (HSDD) has the same form as (HSDP), i.e., (HSDD) is simply (HSDP) with $(\mathbf{y}, \mathbf{x}, \tau, \theta)$ being replaced by $(\mathbf{y}', \mathbf{x}', \tau', \theta')$.

ii) (HSDP) has a *strictly* feasible point

$$\mathbf{y} = \mathbf{0}, \quad \mathbf{x} = \mathbf{e} > \mathbf{0}, \quad \tau = 1, \quad \theta = 1, \quad \mathbf{s} = \mathbf{e} > \mathbf{0}, \quad \kappa = 1.$$

iii) (HSDP) has an optimal solution and its optimal solution set is *bounded*.

iv) The optimal value of (HSDP) is zero, and

$$(\mathbf{y}, \mathbf{x}, \tau, \theta, \mathbf{s}, \kappa) \in \mathcal{F}_h \quad \text{implies that} \quad (n+1)\theta = \mathbf{x}^T \mathbf{s} + \tau \kappa.$$

v) There is an optimal solution $(\mathbf{y}^*, \mathbf{x}^*, \tau^*, \theta^* = 0, \mathbf{s}^*, \kappa^*) \in \mathcal{F}_h$ such that

$$\begin{pmatrix} \mathbf{x}^* + \mathbf{s}^* \\ \tau^* + \kappa^* \end{pmatrix} > \mathbf{0},$$

which we call a *strictly self-complementary solution*. (Similarly, we sometimes call an optimal solution to (HSDP) a self-complementary solution; the strict inequalities above need not hold.)

Theorem 4 Let $(\mathbf{y}^*, \mathbf{x}^*, \tau^*, \theta^* = 0, \mathbf{s}^*, \kappa^*)$ be a strictly self complementary solution for (HSDP).

- i) (LP) has a solution (*feasible and bounded*) if and only if $\tau^* > 0$. In this case, \mathbf{x}^* / τ^* is an optimal solution for (LP) and $(\mathbf{y}^* / \tau^*, \mathbf{s}^* / \tau^*)$ is an optimal solution for (LD).
- ii) (LP) has *no solution* if and only if $\kappa^* > 0$. In this case, \mathbf{x}^* / κ^* or \mathbf{s}^* / κ^* or both are certificates for proving *infeasibility*: if $\mathbf{c}^T \mathbf{x}^* < 0$ then (LD) is infeasible; if $-\mathbf{b}^T \mathbf{y}^* < 0$ then (LP) is infeasible; and if both $\mathbf{c}^T \mathbf{x}^* < 0$ and $-\mathbf{b}^T \mathbf{y}^* < 0$ then both (LP) and (LD) are infeasible.

Theorem 5 i) For any $\mu > 0$, there is a unique $(\mathbf{y}, \mathbf{x}, \tau, \theta, \mathbf{s}, \kappa)$ in \mathcal{F}_h^0 , such that

$$\begin{pmatrix} X\mathbf{s} \\ \tau\kappa \end{pmatrix} = \mu \mathbf{e}.$$

ii) Let $(\mathbf{d}_y, \mathbf{d}_x, d_\tau, d_\theta, \mathbf{d}_s, d_\kappa)$ be in the null space of the constraint matrix of (HSDP) after adding surplus variables \mathbf{s} and κ , i.e.,

$$\begin{aligned} A\mathbf{d}_x - \mathbf{b}d_\tau + \bar{\mathbf{b}}d_\theta &= \mathbf{0}, \\ -A^T\mathbf{d}_y + \mathbf{c}d_\tau - \bar{\mathbf{c}}d_\theta - \mathbf{d}_s &= \mathbf{0}, \\ \mathbf{b}^T\mathbf{d}_y - \mathbf{c}^T\mathbf{d}_x + \bar{\mathbf{z}}d_\theta - d_\kappa &= 0, \\ -\bar{\mathbf{b}}^T\mathbf{d}_y + \bar{\mathbf{c}}^T\mathbf{d}_x - \bar{\mathbf{z}}d_\tau &= 0. \end{aligned} \tag{1}$$

$$(\mathbf{d}_x)^T \mathbf{d}_s + d_\tau d_\kappa = 0.$$

Endogenous Potential Function and Central Path

$$\psi_{n+\rho}(\mathbf{x}, \mathbf{s}, \tau, \kappa) := (n+1+\rho) \log(\mathbf{x}^T \mathbf{s} + \tau \kappa) - \sum_{j=1}^n \log(x_j s_j) - \log(\tau \kappa),$$

and

$$\mathcal{C} = \left\{ (\mathbf{y}, \mathbf{x}, \tau, \theta, \mathbf{s}, \kappa) \in \mathcal{F}_h^0 : \begin{pmatrix} X\mathbf{s} \\ \tau\kappa \end{pmatrix} = \frac{\mathbf{x}^T \mathbf{s} + \tau\kappa}{n+1} \mathbf{e} \right\}.$$

Obviously, the initial interior feasible point proposed in Theorem 3 is on the path with $\mu = 1$ or $(\mathbf{x}^0)^T \mathbf{s}^0 + \tau^0 \kappa^0 = n+1$.

Solving (HSDP)

Consider solving the following **system of linear equations** for $(\mathbf{d}_y, \mathbf{d}_x, d_\tau, d_\theta, \mathbf{d}_s, d_\kappa)$ that satisfies (1) and

$$\begin{pmatrix} X\mathbf{d}_s + S\mathbf{d}_x \\ \tau^k d_\kappa + \kappa^k d_\tau \end{pmatrix} = \gamma\mu\mathbf{e} - \begin{pmatrix} X\mathbf{s} \\ \tau\kappa \end{pmatrix}.$$

Theorem 6 The $O(\sqrt{n} \log((\mathbf{x}^0)^T \mathbf{s}^0 / \epsilon))$ interior-point algorithm, coupled with a termination technique described above, generates a *strictly self-complementary solution* for (HSDP) in $O(\sqrt{n}(\log(c(A, \mathbf{b}, \mathbf{c})) + \log n))$ iterations and $O(n^3(\log(c(A, \mathbf{b}, \mathbf{c})) + \log n))$ operations, where $c(A, \mathbf{b}, \mathbf{c})$ is a positive number depending on the data $(A, \mathbf{b}, \mathbf{c})$. If (LP) and (LD) have integer data with *bit length* L , then by the construction, the data of (HSDP) remains integral and its length is $O(L)$. Moreover, $c(A, \mathbf{b}, \mathbf{c}) \leq 2^L$. Thus, the algorithm terminates in $O(\sqrt{n}L)$ iterations and $O(n^3L)$ operations.

Example

Consider the example where

$$A = \begin{pmatrix} -1 & 0 & 0 \end{pmatrix}, \quad b = 1, \quad \text{and} \quad \mathbf{c} = \begin{pmatrix} 0 & 1 & -1 \end{pmatrix}.$$

Then,

$$y^* = 2, \quad \mathbf{x}^* = (0, 2, 1)^T, \quad \tau^* = 0, \quad \theta^* = 0, \quad \mathbf{s}^* = (2, 0, 0)^T, \quad \kappa^* = 1$$

could be a strictly self-complementary solution generated for (HSDP) with

$$\mathbf{c}^T \mathbf{x}^* = 1 > 0, \quad by^* = 2 > 0.$$

Thus (y^*, \mathbf{s}^*) demonstrates the infeasibility of (LP), but \mathbf{x}^* doesn't show the infeasibility of (LD). Of course, if the algorithm generates instead $\mathbf{x}^* = (0, 1, 2)^T$, then we get demonstrated infeasibility of both.

Primal-Dual Interior-Point Algorithm for Monotone LCP

Find $(\mathbf{x}, \mathbf{s}) \geq 0$ such that

$$\mathbf{s} = M\mathbf{x} + \mathbf{q}, \mathbf{x}^T \mathbf{s} = 0, (\mathbf{x}, \mathbf{s}) \geq 0.$$

Once have a pair $(\mathbf{x}, \mathbf{s}) \in \text{int } \mathcal{F}$, we compute **direction vectors** \mathbf{d}_x and \mathbf{d}_s from the Newton system equations:

$$\begin{aligned} S\mathbf{d}_x + X\mathbf{d}_s &= \frac{\mathbf{x}^T \mathbf{s}}{n+\rho} \mathbf{e} - XSe, \\ M\mathbf{d}_x - \mathbf{d}_s &= \mathbf{0}. \end{aligned} \tag{2}$$

Note that $\mathbf{d}_x^T \mathbf{d}_s = \mathbf{d}_x^T M\mathbf{d}_x \geq 0$ if M is monotone.

Lemma 1 Let the direction vector $\mathbf{d} = (\mathbf{d}_x, \mathbf{d}_s)$ be generated by equation (2), and let

$$\theta = \frac{\alpha \sqrt{\min(XS\mathbf{e})}}{\|(XS)^{-1/2}(\frac{\mathbf{x}^T \mathbf{s}}{n+\rho} \mathbf{e} - XS\mathbf{e})\|}, \quad (3)$$

where α is a *positive constant* less than 1. Let

$$\mathbf{x}^+ = \mathbf{x} + \theta \mathbf{d}_x \quad \text{and} \quad \mathbf{s}^+ = \mathbf{s} + \theta \mathbf{d}_s.$$

Then, we have $(\mathbf{x}^+, \mathbf{y}^+, \mathbf{s}^+) \in \text{int } \mathcal{F}$ and

$$\begin{aligned} \psi_{n+\rho}(\mathbf{x}^+, \mathbf{s}^+) - \psi_{n+\rho}(\mathbf{x}, \mathbf{s}) &\leq -\alpha \sqrt{\min(XS\mathbf{e})} \|(XS)^{-1/2}(\mathbf{e} - \frac{(n+\rho)}{\mathbf{x}^T \mathbf{s}} XS\mathbf{s})\| \\ &\quad + \alpha^2 \max\left\{ \frac{(n+\rho)\mathbf{d}_x^T \mathbf{d}_s}{\mathbf{x}^T \mathbf{s}}, \frac{\|X^{-1}\mathbf{d}_s\|^2 + \|S^{-1}\mathbf{d}_x\|^2}{2(1-\alpha)} \right\} \\ &\leq -\frac{\sqrt{3}}{2}\alpha + \alpha^2 \max\left\{ \frac{n+\rho}{2n}, \frac{1}{2(1-\alpha)} \right\}. \end{aligned}$$

Software Implementation

Cplex

SEDUMI: <http://sedumi.mcmaster.ca/>

MOSEK: http://www.mosek.com/products_mosek.html

IPOPT: <https://projects.coin-or.org/Ipopt>

sphslfbf: Sparse Linear Programming Solver (Matlab .m file).