

# Orthogonal Projection

Alvin Kim

July 17, 2024



## 1 Introduction

- Definitions
- Example

## 2 Formulas

- Projection onto a Vector
- Projection onto a Subspace
- The Projection Matrix

## 3 Properties

- Projection Minimizes Error
- $P^T$
- $P^2$

## 4 Special Cases

- Sanity Check:  $A$  Has One Column
- Projection from  $R^3$  onto y-axis
- Projection from  $R^3$  onto xz-plane
- $A$  is an Orthonormal Matrix
- $\vec{b} \perp S$
- $\vec{b} \in S$

## 5 Conclusion

- Formulas Recapped
- Properties Recapped
- Conclusion



# What is a projection?

$\text{proj}_S \vec{b}$ , the projection of vector  $\vec{b}$  onto subspace  $S$ , is the vector inside of  $S$  closest to  $\vec{b}$ .





# What is a projection?

$\text{proj}_S \vec{b}$ , the projection of vector  $\vec{b}$  onto subspace  $S$ , is the vector inside of  $S$  closest to  $\vec{b}$ .

## Note

In this presentation, projection refers to specifically orthogonal projection,  $\vec{p}$  refers to  $\text{proj}_S \vec{b}$ , and  $\vec{e}$  refers to the error vector  $\vec{e} = \vec{b} - \vec{p}$ .



# What is a projection?

$\text{proj}_S \vec{b}$ , the projection of vector  $\vec{b}$  onto subspace  $S$ , is the vector inside of  $S$  closest to  $\vec{b}$ .

Closest means that the error is orthogonal to  $S$ .



# What is a projection?

$\text{proj}_S \vec{b}$ , the projection of vector  $\vec{b}$  onto subspace  $S$ , is the vector inside of  $S$  closest to  $\vec{b}$ .


Closest means that the error is orthogonal to  $S$ .

## Note

The dimension of subspace  $S$  can be anything from 0 to the dimension of the space we are working inside.




# Example



## Example

Draw  $\text{proj}_a b$ .

# Example




## Note


When projecting a vector onto another vector  $\vec{a}$ , the subspace  $S$  is the span of  $\vec{a}$ .



# Example



# Example





# Projection onto a Vector

Consider projecting  $\vec{b}$  onto  $\vec{a}$ .



# Projection onto a Vector

Consider projecting  $\vec{b}$  onto  $\vec{a}$ .

What we know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{p} = c\vec{a}$$

$$\vec{e} \cdot \vec{a} = 0$$



# Projection onto a Vector

From that:

Consider projecting  $\vec{b}$  onto  $\vec{a}$ .

What we know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{p} = c\vec{a}$$

$$\vec{e} \cdot \vec{a} = 0$$

$$\vec{e} = \vec{b} - c\vec{a}$$

$$(\vec{b} - c\vec{a}) \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} - c\vec{a} \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} = c\vec{a} \cdot \vec{a}$$

$$c = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}}$$

$$\boxed{\vec{p} = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}} \vec{a}}$$

# Projection onto a Vector

From that:

Consider projecting  $\vec{b}$  onto  $\vec{a}$ .

What we know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{p} = c\vec{a}$$

$$\vec{e} \cdot \vec{a} = 0$$

$$\vec{e} = \vec{b} - c\vec{a}$$

$$(\vec{b} - c\vec{a}) \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} - c\vec{a} \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} = c\vec{a} \cdot \vec{a}$$

$$c = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}}$$

$$\boxed{\vec{p} = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}} \vec{a}}$$

# Projection onto a Vector

From that:

Consider projecting  $\vec{b}$  onto  $\vec{a}$ .

What we know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{p} = c\vec{a}$$

$$\vec{e} \cdot \vec{a} = 0$$

$$\vec{e} = \vec{b} - c\vec{a}$$

$$(\vec{b} - c\vec{a}) \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} - c\vec{a} \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} = c\vec{a} \cdot \vec{a}$$

$$c = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}}$$

$$\boxed{\vec{p} = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}} \vec{a}}$$

# Projection onto a Vector

From that:

Consider projecting  $\vec{b}$  onto  $\vec{a}$ .

What we know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{p} = c\vec{a}$$

$$\vec{e} \cdot \vec{a} = 0$$

$$\vec{e} = \vec{b} - c\vec{a}$$

$$(\vec{b} - c\vec{a}) \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} - c\vec{a} \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} = c\vec{a} \cdot \vec{a}$$

$$c = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}}$$

$$\boxed{\vec{p} = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}} \vec{a}}$$



# Projection onto a Vector

From that:

Consider projecting  $\vec{b}$  onto  $\vec{a}$ .

What we know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{p} = c\vec{a}$$

$$\vec{e} \cdot \vec{a} = 0$$

$$\vec{e} = \vec{b} - c\vec{a}$$

$$(\vec{b} - c\vec{a}) \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} - c\vec{a} \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} = c\vec{a} \cdot \vec{a}$$

$$c = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}}$$

$$\boxed{\vec{p} = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}} \vec{a}}$$

# Projection onto a Vector

From that:

Consider projecting  $\vec{b}$  onto  $\vec{a}$ .

What we know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{p} = c\vec{a}$$

$$\vec{e} \cdot \vec{a} = 0$$

$$\vec{e} = \vec{b} - c\vec{a}$$

$$(\vec{b} - c\vec{a}) \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} - c\vec{a} \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} = c\vec{a} \cdot \vec{a}$$

$$c = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}}$$

$$\vec{p} = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}} \vec{a}$$



# Project $\vec{b}$ onto $S$





# Project $\vec{b}$ onto $S$

Suppose that  $\vec{a}_1, \vec{a}_2, \dots, \vec{a}_n$  are a basis of  $S$ , let

$$A = \begin{bmatrix} | & | & \dots & | \\ \vec{a}_1 & \vec{a}_2 & \dots & \vec{a}_n \\ | & | & \dots & | \end{bmatrix}$$

and let  $\vec{p} = A\vec{c}$ . We also know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$C(A)^\perp = N(A^T)$$



# Project $\vec{b}$ onto $S$

Suppose that  $\vec{a}_1, \vec{a}_2, \dots, \vec{a}_n$  are a basis of  $S$ , let

$$A = \begin{bmatrix} | & | & \dots & | \\ \vec{a}_1 & \vec{a}_2 & \dots & \vec{a}_n \\ | & | & \dots & | \end{bmatrix}$$

and let  $\vec{p} = A\vec{c}$ . We also know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$A^T \vec{e} = \vec{0}$$



# Project $\vec{b}$ onto $S$

Suppose that  $\vec{a}_1, \vec{a}_2, \dots, \vec{a}_n$  are a basis of  $S$ , let

$$A = \begin{bmatrix} | & | & \dots & | \\ \vec{a}_1 & \vec{a}_2 & \dots & \vec{a}_n \\ | & | & \dots & | \end{bmatrix}$$

and let  $\vec{p} = A\vec{c}$ . We also know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$A^T \vec{e} = \vec{0}$$

Then,

$$\vec{e} = \vec{b} - A\vec{c}$$

$$A^T(\vec{b} - A\vec{c}) = \vec{0}$$

$$A^T \vec{b} - A^T A \vec{c} = \vec{0}$$

$$A^T \vec{b} = A^T A \vec{c}$$

$$(A^T A)^{-1} A^T \vec{b} = \vec{c}$$

$$\boxed{\vec{c} = (A^T A)^{-1} A^T \vec{b}}$$

$$\boxed{\vec{p} = A(A^T A)^{-1} A^T \vec{b}}$$

# The Projection Matrix

Recall that for some matrix  $A$ , the projection  
$$\vec{p} = A(A^T A)^{-1} A^T \vec{b}.$$

## Observation

For a specific  $A$ ,  $\vec{p}$  is an unchanging linear transformation of  $\vec{b}$ . So, we can represent the linear transformation with a matrix  $P$ .



# The Projection Matrix

Recall that for some matrix  $A$ , the projection

$$\vec{p} = A(A^T A)^{-1} A^T \vec{b}.$$




# The Projection Matrix

Recall that for some matrix  $A$ , the projection  
 $\vec{p} = A(A^T A)^{-1} A^T \vec{b}$ .

Let  $\boxed{P = A(A^T A)^{-1} A^T}$ . Then,  $\vec{p} = P\vec{b}$ .

## Note

$A$  must have linearly independent columns.

- Applications
  - Graphics
  - Least Squares Regression
- Further Learning
  - Practice
  - *Introduction to Linear Algebra* 6th Edition Chapter 4.2 by Gilbert Strang