

Introduction

Formulas

Properties

Special Cases

Conclusion

Orthogonal Projection

Alvin Kim

July 19, 2024

MATH
KIMCHI

Alvin Kim

Orthogonal Projection

2024-07-19

Orthogonal Projection

Alvin Kim

July 19, 2024

1 Introduction

- Definitions
- Example

2 Formulas

- Projection onto a Vector
- Projection onto a Subspace
- The Projection Matrix
- The Error Matrix

3 Properties

- Projection Minimizes Error
- $P = P^T$
- $P = P^2$
- Sanity Check: $E = E^2$

4 Special Cases

- Sanity Check: A has One Column
- A is an Orthonormal Matrix
- Projection from \mathbb{R}^3 onto the xz -plane
- More Special Cases

5 Conclusion

- Key Ideas
- Next Steps



2024-07-19

Orthogonal Projection

1 Introduction	1 Special Cases
■ Definitions	■ Sanity Check: A has One Column
■ Example	■ A is an Orthonormal Matrix
2 Formulas	■ Projection from \mathbb{R}^3 onto the xz -plane
■ Projection onto a Vector	■ More Special Cases
■ Projection onto a Subspace	3 Conclusion
■ The Projection Matrix	■ Key Ideas
■ The Error Matrix	■ Next Steps
3 Properties	
■ Projection Minimizes Error	
■ $P = P^T$	
■ $P = P^2$	
■ Sanity Check: $E = E^2$	

What is a projection?

$\text{proj}_S \vec{b}$, the projection of vector \vec{b} onto subspace S , is the vector inside of S closest to \vec{b} .



Orthogonal Projection

└ Introduction

└ Definitions

└ What is a projection?

What is a projection?

$\text{proj}_S \vec{b}$, the projection of vector \vec{b} onto subspace S , is the vector inside of S closest to \vec{b} .

Note

In this presentation, projection refers to specifically orthogonal projection, \vec{p} refers to $\text{proj}_S \vec{b}$, and \vec{e} refers to the error vector $\vec{e} = \vec{b} - \vec{p}$. The error vector is also called the rejection.



Orthogonal Projection

└ Introduction

└ Definitions

└ What is a projection?

What is a projection?

$\text{proj}_S \vec{b}$, the projection of vector \vec{b} onto subspace S , is the vector inside of S closest to \vec{b} .

Note

In this presentation, projection refers to specifically orthogonal projection, \vec{p} refers to $\text{proj}_S \vec{b}$, and \vec{e} refers to the error vector $\vec{e} = \vec{b} - \vec{p}$. The error vector is also called the rejection.

What is a projection?

$\text{proj}_S \vec{b}$, the projection of vector \vec{b} onto subspace S , is the vector inside of S closest to \vec{b} .

Closest means that the error is orthogonal to S .



Orthogonal Projection

└ Introduction

└ Definitions

└ What is a projection?

The reason why I am using this meaning rather than the minimizing error meaning is because this meaning is simpler to work with. Ultimately, it can be proven that both meanings are the same.

What is a projection?

$\text{proj}_S \vec{b}$, the projection of vector \vec{b} onto subspace S , is the vector inside of S closest to \vec{b} .

Closest means that the error is orthogonal to S .

Note

The dimension of subspace S can be anything from 0 to the dimension of the space we are working inside.



Orthogonal Projection

└ Introduction

└ Definitions

└ What is a projection?

What is a projection?


$\text{proj}_S \vec{b}$, the projection of vector \vec{b} onto subspace S , is the vector inside of S closest to \vec{b} .
Closest means that the error is orthogonal to S .

Note

The dimension of subspace S can be anything from 0 to the dimension of the space we are working inside.

The reason why I am using this meaning rather than the minimizing error meaning is because this meaning is simpler to work with. Ultimately, it can be proven that both meanings are the same.

Example



Example

Draw $\text{proj}_a b$.



2024-07-19

Orthogonal Projection

└ Introduction

└ Example

└ Example


Example



Example

Draw $\text{proj}_a b$.

Example



Note

When projecting a vector onto another vector \vec{a} , the subspace S is the span of \vec{a} .



Orthogonal Projection

└ Introduction

└ Example


└ Example



Note

When projecting a vector onto another vector \vec{a} , the subspace S is the span of \vec{a} .

Example



2024-07-19

Orthogonal Projection

└ Introduction


└ Example

└ Example

Example



Example



2024-07-19

Orthogonal Projection

└ Introduction

└ Example

└ Example



Projection onto a Vector

Consider projecting \vec{b} onto \vec{a} .



Orthogonal Projection

└ Formulas

└ Projection onto a Vector

└ Projection onto a Vector

Projection onto a Vector

Consider projecting \vec{b} onto \vec{a} .

What we know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{p} = c\vec{a}$$

$$\vec{e} \cdot \vec{a} = 0$$



Orthogonal Projection

└ Formulas

└ Projection onto a Vector

└ Projection onto a Vector

Consider projecting \vec{b} onto \vec{a} .
What we know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{p} = c\vec{a}$$

$$\vec{e} \cdot \vec{a} = 0$$

If $\vec{a} = \vec{0}$, then p will be 0 because p must be in the zero subspace.

If $\vec{a} \neq 0$, then we are projecting b onto the line that spans \vec{a} . tell the audience to try this

Projection onto a Vector

From that:

Consider projecting \vec{b} onto \vec{a} .

What we know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{p} = c\vec{a}$$

$$\vec{e} \cdot \vec{a} = 0$$

$$\vec{e} = \vec{b} - c\vec{a}$$

$$(\vec{b} - c\vec{a}) \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} - c\vec{a} \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} = c\vec{a} \cdot \vec{a}$$

$$c = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}}$$

$$\vec{p} = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}} \vec{a}$$



2024-07-19

Orthogonal Projection

└ Formulas

└ Projection onto a Vector

└ Projection onto a Vector

From that:

Consider projecting \vec{b} onto \vec{a} .

What we know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{p} = c\vec{a}$$

$$\vec{e} \cdot \vec{a} = 0$$

$$\vec{e} = \vec{b} - c\vec{a}$$

$$(\vec{b} - c\vec{a}) \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} - c\vec{a} \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} = c\vec{a} \cdot \vec{a}$$

$$c = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}}$$

$$\vec{p} = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}} \vec{a}$$

If $\vec{a} = \vec{0}$, then p will be 0 because p must be in the zero subspace.

If $\vec{a} \neq \vec{0}$, then we are projecting b onto the line that spans \vec{a} . tell the audience to try this

Projection onto a Vector

From that:

Consider projecting \vec{b} onto \vec{a} .

What we know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{p} = c\vec{a}$$

$$\vec{e} \cdot \vec{a} = 0$$

$$\vec{e} = \vec{b} - c\vec{a}$$

$$(\vec{b} - c\vec{a}) \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} - c\vec{a} \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} = c\vec{a} \cdot \vec{a}$$

$$c = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}}$$

$$\vec{p} = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}} \vec{a}$$



2024-07-19

Orthogonal Projection

└ Formulas

└ Projection onto a Vector

└ Projection onto a Vector

Projection onto a Vector

From that:

Consider projecting \vec{b} onto \vec{a} .
What we know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{p} = c\vec{a}$$

$$\vec{e} \cdot \vec{a} = 0$$

$$\vec{e} = \vec{b} - c\vec{a}$$

$$(\vec{b} - c\vec{a}) \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} - c\vec{a} \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} = c\vec{a} \cdot \vec{a}$$

$$c = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}}$$

$$\vec{p} = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}} \vec{a}$$

If $\vec{a} = \vec{0}$, then p will be 0 because p must be in the zero subspace.

If $\vec{a} \neq \vec{0}$, then we are projecting b onto the line that spans \vec{a} . tell the audience to try this

Projection onto a Vector

From that:

Consider projecting \vec{b} onto \vec{a} .

What we know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{p} = c\vec{a}$$

$$\vec{e} \cdot \vec{a} = 0$$

$$\vec{e} = \vec{b} - c\vec{a}$$

$$(\vec{b} - c\vec{a}) \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} - c\vec{a} \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} = c\vec{a} \cdot \vec{a}$$

$$c = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}}$$

$$\vec{p} = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}} \vec{a}$$



2024-07-19

Orthogonal Projection

└ Formulas

└ Projection onto a Vector

└ Projection onto a Vector

Projection onto a Vector

From that:

Consider projecting \vec{b} onto \vec{a} .
What we know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{p} = c\vec{a}$$

$$\vec{e} \cdot \vec{a} = 0$$

$$\vec{e} = \vec{b} - c\vec{a}$$

$$(\vec{b} - c\vec{a}) \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} - c\vec{a} \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} = c\vec{a} \cdot \vec{a}$$

$$c = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}}$$

$$\vec{p} = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}} \vec{a}$$

If $\vec{a} = \vec{0}$, then p will be 0 because p must be in the zero subspace.

If $\vec{a} \neq \vec{0}$, then we are projecting b onto the line that spans \vec{a} . tell the audience to try this

Projection onto a Vector

From that:

Consider projecting \vec{b} onto \vec{a} .

What we know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{p} = c\vec{a}$$

$$\vec{e} \cdot \vec{a} = 0$$

$$\vec{e} = \vec{b} - c\vec{a}$$

$$(\vec{b} - c\vec{a}) \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} - c\vec{a} \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} = c\vec{a} \cdot \vec{a}$$

$$c = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}}$$

$$\vec{p} = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}} \vec{a}$$



2024-07-19

Orthogonal Projection

└ Formulas

└ Projection onto a Vector

└ Projection onto a Vector

Projection onto a Vector

From that:

Consider projecting \vec{b} onto \vec{a} .
What we know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{p} = c\vec{a}$$

$$\vec{e} \cdot \vec{a} = 0$$

$$\vec{e} = \vec{b} - c\vec{a}$$

$$(\vec{b} - c\vec{a}) \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} - c\vec{a} \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} = c\vec{a} \cdot \vec{a}$$

$$c = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}}$$

$$\vec{p} = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}} \vec{a}$$

If $\vec{a} = \vec{0}$, then p will be 0 because p must be in the zero subspace.

If $\vec{a} \neq \vec{0}$, then we are projecting b onto the line that spans \vec{a} . tell the audience to try this

Projection onto a Vector

From that:

Consider projecting \vec{b} onto \vec{a} .

What we know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{p} = c\vec{a}$$

$$\vec{e} \cdot \vec{a} = 0$$

$$\vec{e} = \vec{b} - c\vec{a}$$

$$(\vec{b} - c\vec{a}) \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} - c\vec{a} \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} = c\vec{a} \cdot \vec{a}$$

$$c = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}}$$

$$\vec{p} = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}} \vec{a}$$



2024-07-19

Orthogonal Projection

└ Formulas

└ Projection onto a Vector

└ Projection onto a Vector

Projection onto a Vector

From that:

Consider projecting \vec{b} onto \vec{a} .

What we know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{p} = c\vec{a}$$

$$\vec{e} \cdot \vec{a} = 0$$

$$\vec{e} = \vec{b} - c\vec{a}$$

$$(\vec{b} - c\vec{a}) \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} - c\vec{a} \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} = c\vec{a} \cdot \vec{a}$$

$$c = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}}$$

$$\vec{p} = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}} \vec{a}$$

If $\vec{a} = \vec{0}$, then p will be 0 because p must be in the zero subspace.

If $\vec{a} \neq \vec{0}$, then we are projecting b onto the line that spans \vec{a} . tell the audience to try this

Projection onto a Vector

From that:

Consider projecting \vec{b} onto \vec{a} .

What we know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{p} = c\vec{a}$$

$$\vec{e} \cdot \vec{a} = 0$$

$$\vec{e} = \vec{b} - c\vec{a}$$

$$(\vec{b} - c\vec{a}) \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} - c\vec{a} \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} = c\vec{a} \cdot \vec{a}$$

$$c = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}}$$

$$\vec{p} = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}} \vec{a}$$



2024-07-19

Orthogonal Projection

└ Formulas

└ Projection onto a Vector

└ Projection onto a Vector

Projection onto a Vector

From that:

Consider projecting \vec{b} onto \vec{a} .

What we know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{p} = c\vec{a}$$

$$\vec{e} \cdot \vec{a} = 0$$

$$\vec{e} = \vec{b} - c\vec{a}$$

$$(\vec{b} - c\vec{a}) \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} - c\vec{a} \cdot \vec{a} = 0$$

$$\vec{b} \cdot \vec{a} = c\vec{a} \cdot \vec{a}$$

$$c = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}}$$

$$\vec{p} = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}} \vec{a}$$

If $\vec{a} = \vec{0}$, then p will be 0 because p must be in the zero subspace.

If $\vec{a} \neq \vec{0}$, then we are projecting b onto the line that spans \vec{a} . tell the audience to try this

Project \vec{b} onto S

2024-07-19

Orthogonal Projection

└ Formulas

└ Projection onto a Subspace

└ Project \vec{b} onto S Project \vec{b} onto S

the a vecs are LI, but the A matrix might be rectangular so it isn't invertible there is guaranteed to be exactly one c for a specific p and A bc p is in S and A has LI colsthe orthogonal complement of col space is the left null spacethe orthogonal complement of col space is the left null space tell them to try this proof Hint: ATA is invertible bc A has LI columnsthe coefficients vector is also good to remember, because it is sometimes more useful than the projectionas mentioned before, we can't distribute the inverse sign because A might be rectangular



Project \vec{b} onto S

Suppose that $\vec{a}_1, \vec{a}_2, \dots, \vec{a}_n$ are a basis of S , let

$$A = \begin{bmatrix} | & | & & | \\ \vec{a}_1 & \vec{a}_2 & \dots & \vec{a}_n \\ | & | & & | \end{bmatrix}$$

and let $\vec{p} = A\vec{c}$. We also know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$C(A)^\perp = N(A^T)$$



Orthogonal Projection

└ Formulas

└ Projection onto a Subspace

└ Project \vec{b} onto S

$$A = \begin{bmatrix} | & | & & | \\ \vec{a}_1 & \vec{a}_2 & \dots & \vec{a}_n \\ | & | & & | \end{bmatrix}$$

$$\vec{e} = \vec{b} - \vec{p}$$

$$C(A)^\perp = N(A^T)$$

the \vec{a} vecs are LI, but the A matrix might be rectangular so it isn't invertible there is guaranteed to be exactly one \vec{c} for a specific \vec{p} and A bc \vec{p} is in S and A has LI cols the orthogonal complement of col space is the left null space the orthogonal complement of col space is the left null space tell them to try this proof Hint: ATA is invertible bc A has LI columns the coefficients vector is also good to remember, because it is sometimes more useful than the projection as mentioned before, we can't distribute the inverse sign because A might be rectangular

Project \vec{b} onto S

Suppose that $\vec{a}_1, \vec{a}_2, \dots, \vec{a}_n$ are a basis of S , let

$$A = \begin{bmatrix} | & | & & | \\ \vec{a}_1 & \vec{a}_2 & \dots & \vec{a}_n \\ | & | & & | \end{bmatrix}$$

and let $\vec{p} = A\vec{c}$. We also know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$A^T \vec{e} = \vec{0}$$



2024-07-19

Orthogonal Projection

└ Formulas

└ Projection onto a Subspace

└ Project \vec{b} onto S Project \vec{b} onto S

Suppose that $\vec{a}_1, \vec{a}_2, \dots, \vec{a}_n$ are a basis of S , let

$$A = \begin{bmatrix} | & | & & | \\ \vec{a}_1 & \vec{a}_2 & \dots & \vec{a}_n \\ | & | & & | \end{bmatrix}$$

and let $\vec{p} = A\vec{c}$. We also know:

$$\vec{e} = \vec{b} - \vec{p}$$

$$A^T \vec{e} = \vec{0}$$

the a vecs are LI, but the A matrix might be rectangular so it isn't invertible there is guaranteed to be exactly one c for a specific p and A bc p is in S and A has LI colsthe orthogonal complement of col space is the left null spacethe orthogonal complement of col space is the left null space tell them to try this proof Hint: ATA is invertible bc A has LI columnsthe coefficients vector is also good to remember, because it is sometimes more useful than the projectionas mentioned before, we can't distribute the inverse sign because A might be rectangular

Project \vec{b} onto S

Suppose that $\vec{a}_1, \vec{a}_2, \dots, \vec{a}_n$ are a basis of S , let

$$A = \begin{bmatrix} | & | & & | \\ \vec{a}_1 & \vec{a}_2 & \dots & \vec{a}_n \\ | & | & & | \end{bmatrix}$$

and let $\vec{p} = A\vec{c}$. We also know:

$$\begin{aligned}\vec{e} &= \vec{b} - \vec{p} \\ A^T \vec{e} &= \vec{0}\end{aligned}$$

Then,

$$\begin{aligned}\vec{e} &= \vec{b} - A\vec{c} \\ A^T(\vec{b} - A\vec{c}) &= \vec{0} \\ A^T\vec{b} - A^T A\vec{c} &= \vec{0} \\ A^T\vec{b} &= A^T A\vec{c} \\ (A^T A)^{-1} A^T\vec{b} &= \vec{c}\end{aligned}$$

$$\boxed{\vec{c} = (A^T A)^{-1} A^T \vec{b}}$$

$$\boxed{\vec{p} = A(A^T A)^{-1} A^T \vec{b}}$$



2024-07-19

Orthogonal Projection

└ Formulas

└ Projection onto a Subspace

└ Project \vec{b} onto S Project \vec{b} onto S

Suppose that $\vec{a}_1, \vec{a}_2, \dots, \vec{a}_n$ are a basis of S , let

$$A = \begin{bmatrix} | & | & & | \\ \vec{a}_1 & \vec{a}_2 & \dots & \vec{a}_n \\ | & | & & | \end{bmatrix}$$

and let $\vec{p} = A\vec{c}$. We also know:

$$\begin{aligned}\vec{e} &= \vec{b} - \vec{p} \\ A^T \vec{e} &= \vec{0}\end{aligned}$$

Then,

$$\vec{e} = \vec{b} - A\vec{c}$$

$$A^T(\vec{b} - A\vec{c}) = \vec{0}$$

$$A^T\vec{b} - A^T A\vec{c} = \vec{0}$$

$$A^T\vec{b} = A^T A\vec{c}$$

$$(A^T A)^{-1} A^T\vec{b} = \vec{c}$$

$$\boxed{\vec{c} = (A^T A)^{-1} A^T \vec{b}}$$

$$\boxed{\vec{p} = A(A^T A)^{-1} A^T \vec{b}}$$

the a vecs are LI, but the A matrix might be rectangular so it isn't invertible there is guaranteed to be exactly one c for a specific p and A bc p is in S and A has LI colsthe orthogonal complement of col space is the left null space the orthogonal complement of col space is the left null space tell them to try this proof Hint: ATA is invertible bc A has LI columnsthe coefficients vector is also good to remember, because it is sometimes more useful than the projectionas mentioned before, we can't distribute the inverse sign because A might be rectangular

The Projection Matrix

Recall that for some matrix A , the projection
 $\vec{p} = A(A^T A)^{-1} A^T \vec{b}$.

Observation

For a specific A , \vec{p} is an unchanging linear transformation of \vec{b} . So, we can represent the linear transformation with a matrix P .



Orthogonal Projection

└ Formulas

└ The Projection Matrix

└ The Projection Matrix

Recall that for some matrix A , the projection
 $\vec{p} = A(A^T A)^{-1} A^T \vec{b}$.

Observation

For a specific A , \vec{p} is an unchanging linear transformation of \vec{b} . So, we can represent the linear transformation with a matrix P .

To multiply some vector by a matrix is to apply a linear transformation to the vector.

The Projection Matrix

Recall that for some matrix A , the projection
 $\vec{p} = A(A^T A)^{-1} A^T \vec{b}$.



Orthogonal Projection

└ Formulas

└ The Projection Matrix

└ The Projection Matrix

The Projection Matrix

Recall that for some matrix A , the projection

$$\vec{p} = A(A^T A)^{-1} A^T \vec{b}.$$

Let $\boxed{P = A(A^T A)^{-1} A^T}$. Then, $\vec{p} = P\vec{b}$.

Note

A must have linearly independent columns.



Orthogonal Projection

└ Formulas

└ The Projection Matrix

└ The Projection Matrix

The Projection Matrix

Recall that for some matrix A , the projection

$$\vec{p} = A(A^T A)^{-1} A^T \vec{b}.$$

Let $\boxed{P = A(A^T A)^{-1} A^T}$. Then, $\vec{p} = P\vec{b}$.

Note

A must have linearly independent columns.

The Error Matrix

The error vector is \vec{b} projected onto S^\perp .



The error vector is \vec{b} projected onto S^\perp .

The Error Matrix

The error vector is \vec{b} projected onto S^\perp . Since finding the error vector is also a projection, there should be an error matrix E for some P such that $\vec{e} = E\vec{b}$.



The Error Matrix

The error vector is \vec{b} projected onto S^\perp . Since finding the error vector is also a projection, there should be an error matrix E for some P such that $\vec{e} = E\vec{b}$.

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{e} = I\vec{b} - P\vec{b}$$

$$\vec{e} = (I - P)\vec{b}$$

$$\boxed{E = I - P}$$



Orthogonal Projection

└ Formulas

└ The Error Matrix

└ The Error Matrix

The error vector is \vec{b} projected onto S^\perp . Since finding the error vector is also a projection, there should be an error matrix E for some P such that $\vec{e} = E\vec{b}$.

$$\vec{e} = \vec{b} - \vec{p}$$

$$\vec{e} = I\vec{b} - P\vec{b}$$

$$\vec{e} = (I - P)\vec{b}$$

$$\boxed{E = I - P}$$

Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

Hint

$$\|\vec{d}\| > \|\vec{e}\| \iff d^2 > e^2$$



Orthogonal Projection

└ Properties

└ Projection Minimizes Error

└ Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

Hint

$$\|\vec{d}\| > \|\vec{e}\| \iff d^2 > e^2$$

Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$\vec{d}^2 = (\vec{b} - \vec{x})^2$$



Orthogonal Projection

└ Properties

└ Projection Minimizes Error

└ Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$\vec{d}^2 = (\vec{b} - \vec{x})^2$$

magnitudes must be non-negative
vector squared = magnitude of vector squared = vector dot prod with itself
thus, projection minimizes the error's magnitude as well as the error's magnitude squared (foreshadowing LSR)

Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$\vec{d}^2 = (\vec{b} + \vec{0} - \vec{x})^2$$



Orthogonal Projection

└ Properties

└ Projection Minimizes Error

└ Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$\vec{d}^2 = (\vec{b} + \vec{0} - \vec{x})^2$$

magnitudes must be non-negative
vector squared = magnitude of vector squared = vector dot prod with itself
thus, projection minimizes the error's magnitude as well as the error's magnitude squared (foreshadowing LSR)

Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$d^2 = (\vec{b} - \vec{p} + \vec{p} - \vec{x})^2$$



2024-07-19

Orthogonal Projection

└ Properties

└ Projection Minimizes Error

└ Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$d^2 = (\vec{b} - \vec{p} + \vec{p} - \vec{x})^2$$

magnitudes must be non-negative
vector squared = magnitude
of vector squared = vector dot prod with itself
thus, projection
minimizes the error's magnitude as well as the error's magnitude squared (foreshadowing LSR)

Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$d^2 = (\vec{b} - \vec{p} + \vec{p} - \vec{x})^2$$



Orthogonal Projection

└ Properties

└ Projection Minimizes Error

└ Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$d^2 = (\vec{b} - \vec{p} + \vec{p} - \vec{x})^2$$

magnitudes must be non-negative
vector squared = magnitude
of vector squared = vector dot prod with itself
thus, projection
minimizes the error's magnitude as well as the error's magnitude squared (foreshadowing LSR)

Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$\vec{d}^2 = (\vec{e} + \vec{p} - \vec{x})^2$$



Orthogonal Projection

└ Properties

└ Projection Minimizes Error

└ Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$\vec{d}^2 = (\vec{e} + \vec{p} - \vec{x})^2$$

magnitudes must be non-negative
vector squared = magnitude of vector squared = vector dot prod with itself
thus, projection minimizes the error's magnitude as well as the error's magnitude squared (foreshadowing LSR)

Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$\vec{d}^2 = \vec{e}^2 + 2\vec{e} \cdot (\vec{p} - \vec{x}) + (\vec{p} - \vec{x})^2$$



Orthogonal Projection

└ Properties

└ Projection Minimizes Error

└ Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$\vec{d}^2 = \vec{e}^2 + 2\vec{e} \cdot (\vec{p} - \vec{x}) + (\vec{p} - \vec{x})^2$$

magnitudes must be non-negative
vector squared = magnitude of vector squared = vector dot prod with itself
thus, projection minimizes the error's magnitude as well as the error's magnitude squared (foreshadowing LSR)

Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$\vec{d}^2 = \vec{e}^2 + 2\vec{e} \cdot (\vec{p} - \vec{x}) + (\vec{p} - \vec{x})^2$$



Orthogonal Projection

└ Properties

└ Projection Minimizes Error

└ Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$\vec{d}^2 = \vec{e}^2 + 2\vec{e} \cdot (\vec{p} - \vec{x}) + (\vec{p} - \vec{x})^2$$

magnitudes must be non-negative
vector squared = magnitude of vector squared = vector dot prod with itself
thus, projection minimizes the error's magnitude as well as the error's magnitude squared (foreshadowing LSR)

Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$\vec{d}^2 = \vec{e}^2 + (\vec{p} - \vec{x})^2$$



Orthogonal Projection

└ Properties

└ Projection Minimizes Error

└ Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$\vec{d}^2 = \vec{e}^2 + (\vec{p} - \vec{x})^2$$

magnitudes must be non-negative
vector squared = magnitude of vector squared = vector dot prod with itself
thus, projection minimizes the error's magnitude as well as the error's magnitude squared (foreshadowing LSR)

Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$\vec{d}^2 = \vec{e}^2 + (\vec{p} - \vec{x})^2$$

$$\vec{d}^2 - \vec{e}^2 = (\vec{p} - \vec{x})^2$$



Orthogonal Projection

└ Properties

└ Projection Minimizes Error

└ Projection Minimizes Error

Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$\vec{d}^2 = \vec{e}^2 + (\vec{p} - \vec{x})^2$$

$$\vec{d}^2 - \vec{e}^2 = (\vec{p} - \vec{x})^2$$

magnitudes must be non-negative
vector squared = magnitude
of vector squared = vector dot prod with itself
thus, projection
minimizes the error's magnitude as well as the error's magnitude squared (foreshadowing LSR)

Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$\vec{d}^2 = \vec{e}^2 + (\vec{p} - \vec{x})^2$$

$$\vec{d}^2 - \vec{e}^2 = (\vec{p} - \vec{x})^2 > 0$$



Orthogonal Projection

└ Properties

└ Projection Minimizes Error

└ Projection Minimizes Error

Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$\vec{d}^2 = \vec{e}^2 + (\vec{p} - \vec{x})^2$$

$$\vec{d}^2 - \vec{e}^2 = (\vec{p} - \vec{x})^2 > 0$$

magnitudes must be non-negative
vector squared = magnitude of vector squared = vector dot prod with itself
thus, projection minimizes the error's magnitude as well as the error's magnitude squared (foreshadowing LSR)

Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$\begin{aligned}\vec{d}^2 &= \vec{e}^2 + (\vec{p} - \vec{x})^2 \\ \vec{d}^2 - \vec{e}^2 &= (\vec{p} - \vec{x})^2 > 0 \\ \vec{d}^2 &> \vec{e}^2\end{aligned}$$



Orthogonal Projection

└ Properties

└ Projection Minimizes Error

└ Projection Minimizes Error

Projection Minimizes Error

Prove that for a vector $\vec{x} \in S$ where $\vec{x} \neq \vec{p}$ and $\vec{d} = \vec{b} - \vec{x}$, that $\|\vec{d}\| > \|\vec{e}\|$.

$$\begin{aligned}\vec{d}^2 &= \vec{e}^2 + (\vec{p} - \vec{x})^2 \\ \vec{d}^2 - \vec{e}^2 &= (\vec{p} - \vec{x})^2 > 0 \\ \vec{d}^2 &> \vec{e}^2\end{aligned}$$

magnitudes must be non-negative
vector squared = magnitude of vector squared = vector dot prod with itself
thus, projection minimizes the error's magnitude as well as the error's magnitude squared (foreshadowing LSR)

1

9.

Prov



$$P = P^T$$

Prove that $P = P^T$.

Hint

Start with $P = A(A^T A)^{-1} A^T$.



2024-07-19

Orthogonal Projection

└ Properties

└ $P = P^T$

└ Prove that $P = P^T$.

Prove that $P = P^T$.

Hint

Start with $P = A(A^T A)^{-1} A^T$.

recommend them to try this

$$P = P^T$$

Prove that $P = P^T$.

$$\begin{aligned} P &= A(A^T A)^{-1} A^T \\ P^T &= (A(A^T A)^{-1} A^T)^T \\ P^T &= (A^T)^T ((A^T A)^{-1})^T A^T \\ P^T &= A((A^T A)^T)^{-1} A^T \\ P^T &= A(A^T (A^T)^T)^{-1} A^T \\ P^T &= A(A^T A)^{-1} A^T \\ P^T &= P \end{aligned}$$



2024-07-19

Orthogonal Projection

└ Properties

└ $P = P^T$

└ Prove that $P = P^T$.

Prove that $P = P^T$.

$$\begin{aligned} P &= A(A^T A)^{-1} A^T \\ P^T &= (A(A^T A)^{-1} A^T)^T \\ P^T &= (A^T)^T ((A^T A)^{-1})^T A^T \\ P^T &= A((A^T A)^T)^{-1} A^T \\ P^T &= A(A^T (A^T)^T)^{-1} A^T \\ P^T &= A(A^T A)^{-1} A^T \\ P^T &= P \end{aligned}$$

recommend them to try this

$P = P^2$

Prove that $P = P^2$.



2024-07-19

Orthogonal Projection

└ Properties

└ $P = P^2$

└ Prove that $P = P^2$.

Prove that $P = P^2$.

explain why this logically makes sense
recommend them to try this

$$P = P^2$$

Prove that $P = P^2$.

$$P = A(A^T A)^{-1} A^T$$

$$P^2 = (A(A^T A)^{-1} A^T)^2$$

$$P^2 = (A(A^T A)^{-1} A^T)(A(A^T A)^{-1} A^T)$$

$$P^2 = A(A^T A)^{-1} (A^T A) (A^T A)^{-1} A^T$$

$$P^2 = A(A^T A)^{-1} I A^T$$

$$P^2 = A(A^T A)^{-1} A^T$$

$$P^2 = P$$



2024-07-19

Orthogonal Projection

└ Properties

└ $P = P^2$

└ Prove that $P = P^2$.

Prove that $P = P^2$.

$$\begin{aligned} P &= A(A^T A)^{-1} A^T \\ P^2 &= (A(A^T A)^{-1} A^T)^2 \\ P^2 &= (A(A^T A)^{-1} A^T)(A(A^T A)^{-1} A^T) \\ P^2 &= A(A^T A)^{-1} (A^T A) (A^T A)^{-1} A^T \\ P^2 &= A(A^T A)^{-1} I A^T \\ P^2 &= A(A^T A)^{-1} A^T \\ P^2 &= P \end{aligned}$$

explain why this logically makes sense
recommend them to try this

Sanity Check: $E = E^2$ Prove that $E = E^2$.

Reminder

 E is also a projection matrix and $E = I - P$.

$$E = I - P$$

$$E^2 = (I - P)^2$$

$$E^2 = I - 2IP + P^2$$

$$E^2 = I - 2P + P$$

$$E^2 = I - P$$



2024-07-19

Orthogonal Projection

└ Properties

└ Sanity Check: $E = E^2$ └ Prove that $E = E^2$.Prove that $E = E^2$.

Reminder

 E is also a projection matrix and $E = I - P$.

$$E = I - P$$

$$E^2 = (I - P)^2$$

$$E^2 = I - 2IP + P^2$$

$$E^2 = I - 2P + P$$

$$E^2 = I - P$$

recommend them to try this

Sanity Check: A has One Column

Suppose $A = [\vec{a}]$. Then, $\vec{p} = A(A^T A)^{-1} A^T b$ and $\vec{p} = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}} \vec{a}$.



Orthogonal Projection

└ Special Cases

└┐ Sanity Check: A has One Column└┐┐ Sanity Check: A has One Column

This is an informal sanity check mention inner product = dot product
 It is easy to see that $A^T A$ in the former equation corresponds to $\vec{a} \cdot \vec{a}$ and since it is inverted, it is the divisor in the latter equation. $A^T b$ corresponds to $\vec{a} \cdot \vec{b}$. the orders do seem off, but since many of the terms in the latter equation are scalars, the order is flexible.

A is an Orthonormal Matrix

A is an Orthonormal Matrix

Note

An orthonormal matrix has unit columns all orthogonal to each other. In other words, an orthonormal matrix Q will satisfy $Q^T Q = I$.



2024-07-19

Orthogonal Projection

└ Special Cases

└ A is an Orthonormal Matrix

└ A is an Orthonormal Matrix

 A is an Orthonormal Matrix

Note

An orthonormal matrix has unit columns all orthogonal to each other. In other words, an orthonormal matrix Q will satisfy $Q^T Q = I$.

A is an Orthonormal Matrix

A is an Orthonormal Matrix

Note

An orthonormal matrix has unit columns all orthogonal to each other. In other words, an orthonormal matrix Q will satisfy $Q^T Q = I$.

Then, $P = A(A^T A)^{-1} A^T = A I A^T = A A^T$.



2024-07-19

Orthogonal Projection

└ Special Cases

└ A is an Orthonormal Matrix

└ A is an Orthonormal Matrix

 A is an Orthonormal Matrix

Note

An orthonormal matrix has unit columns all orthogonal to each other. In other words, an orthonormal matrix Q will satisfy $Q^T Q = I$.

Then, $P = A(A^T A)^{-1} A^T = A I A^T = A A^T$.

A is an Orthonormal Matrix

A is an Orthonormal Matrix

Note

An orthonormal matrix has unit columns all orthogonal to each other. In other words, an orthonormal matrix Q will satisfy $Q^T Q = I$.

Then, $P = A(A^T A)^{-1} A^T = A I A^T = A A^T$.

Additionally, the coefficients vector, $\vec{c} = (A^T A)^{-1} A^T \vec{b}$, will just be $\vec{c} = A^T \vec{b}$.



2024-07-19

Orthogonal Projection

└ Special Cases

└ A is an Orthonormal Matrix

└ A is an Orthonormal Matrix

 A is an Orthonormal Matrix

Note

An orthonormal matrix has unit columns all orthogonal to each other. In other words, an orthonormal matrix Q will satisfy $Q^T Q = I$.

Then, $P = A(A^T A)^{-1} A^T = A I A^T = A A^T$.

Additionally, the coefficients vector, $\vec{c} = (A^T A)^{-1} A^T \vec{b}$, will just be $\vec{c} = A^T \vec{b}$.

Projection from \mathbb{R}^3 onto the xz-plane



2024-07-19

- Orthogonal Projection
 - └ Special Cases
 - └ Projection from \mathbb{R}^3 onto the xz-plane
 - └ Projection from \mathbb{R}^3 onto the xz-plane

Projection from \mathbb{R}^3 onto the xz-plane

Hint

What are a basis of the xz-plane?



2024-07-19

Orthogonal Projection

└ Special Cases

└ Projection from \mathbb{R}^3 onto the xz-plane└ Projection from \mathbb{R}^3 onto the xz-plane

Hint

What are a basis of the xz-plane?

Projection from \mathbb{R}^3 onto the xz-plane

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix}$$



Orthogonal Projection

└ Special Cases

└ Projection from \mathbb{R}^3 onto the xz-plane

└ Projection from \mathbb{R}^3 onto the xz-plane

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix}$$

Projection from \mathbb{R}^3 onto the xz-plane

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix}$$

Note

A is orthonormal.



Orthogonal Projection

└ Special Cases

└ Projection from \mathbb{R}^3 onto the xz-plane

└ Projection from \mathbb{R}^3 onto the xz-plane

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix}$$

Note

A is orthonormal.

Projection from \mathbb{R}^3 onto the xz-plane

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix}$$

$$P = AA^T = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$



Orthogonal Projection

└ Special Cases

└ Projection from \mathbb{R}^3 onto the xz-plane

└ Projection from \mathbb{R}^3 onto the xz-plane

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix}$$
$$P = AA^T = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Projection from \mathbb{R}^3 onto the xz-plane

$$P = AA^T = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Example

$$P \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} x \\ 0 \\ z \end{bmatrix}$$



Orthogonal Projection

└ Special Cases

└ Projection from \mathbb{R}^3 onto the xz-plane

└ Projection from \mathbb{R}^3 onto the xz-plane

$$P = AA^T = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Example

$$P \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} x \\ 0 \\ z \end{bmatrix}$$

More Special Cases (Practice)

- Find the projection matrix from \mathbb{R}^3 onto the y-axis.
- What would \vec{p} be if $\vec{b} \perp S$? What if $\vec{b} \in S$?



2024-07-19

Orthogonal Projection

- └ Special Cases
 - └ More Special Cases
 - └ More Special Cases (Practice)

- Find the projection matrix from \mathbb{R}^3 onto the y-axis.
- What would \vec{p} be if $\vec{b} \perp S$? What if $\vec{b} \in S$?

emphasize that these are just the special, interesting cases, and that these problems don't represent most problems.

Key Ideas

- The orthogonal projection of \vec{b} onto subspace S , $\vec{p} = \text{proj}_S \vec{b} \in S$, makes the error $\vec{e} = \vec{b} - \vec{p}$ orthogonal to S .
- The projection also minimizes the error.
- $$P = A(A^T A)A^T$$
- $P = P^2$
- Other useful equations: $\vec{p} = Pb$, $E = I - P$,
 $\vec{e} = (A^T A)A^T b$, $P = P^T$, $c = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}}$, $\vec{p} = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}} \vec{a}$, $\vec{p} = c\vec{a}$,
 $\vec{p} = A\vec{c}$



Orthogonal Projection

└ Conclusion

└ Key Ideas

└ Key Ideas

- The orthogonal projection of \vec{b} onto subspace S , $\vec{p} = \text{proj}_S \vec{b} \in S$, makes the error $\vec{e} = \vec{b} - \vec{p}$ orthogonal to S .
- The projection also minimizes the error.
- $$P = A(A^T A)A^T$$
- $P = P^2$
- Other useful equations: $\vec{p} = Pb$, $E = I - P$,
 $\vec{e} = (A^T A)A^T b$, $P = P^T$, $c = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}}$, $\vec{p} = \frac{\vec{a} \cdot \vec{b}}{\vec{a} \cdot \vec{a}} \vec{a}$, $\vec{p} = c\vec{a}$,
 $\vec{p} = A\vec{c}$

A must have LI cols

Next Steps

- Applications
 - Graphics
 - Least Squares Regression
 - Orthonormal and Orthogonal Matrices
- Further Learning
 - Practice
 - *Introduction to Linear Algebra* 6th Edition Chapter 4.2 by Gilbert Strang



2024-07-19

Orthogonal Projection

└ Conclusion

└ Next Steps

└ Next Steps

- Applications
 - Graphics
 - Least Squares Regression
 - Orthonormal and Orthogonal Matrices
- Further Learning
 - Practice
 - *Introduction to Linear Algebra* 6th Edition Chapter 4.2 by Gilbert Strang