

Subset-sum landscapes under deformed and random measures: the biased invariant $\Gamma^{(q)}$ and unconditional rigidity

(working title — scaffolding, round 100)

Kentaro Amauchi*

August 9, 2026

Abstract

[**TODO** written last.] Papers 1–3 fixed both the sequence and the measure. This paper moves the measure along two axes — bias the coin, and randomise the sequence — and collects the two open questions recorded in paper 3.

1 Introduction

[**TODO** full introduction.] The two deformation axes, and why each is worth taking:

- *Bias.* The classification theorem of paper 1 is a statement about which elements lie on which side of a target; it says nothing about how configurations are weighted. So the whole combinatorial layer survives verbatim when the uniform measure is replaced by the Bernoulli(q) product measure, and the gap series deforms into an explicit one-parameter family. §2 does this, and the answer has a shape worth stating in the introduction: **the fair coin gives the least rugged landscape**.
- *Randomness.* For a random odd sequence the minor-arc input that cost paper 2 fifty rounds is replaced by a second-moment bound. §4 is therefore expected to carry *no ineffective constant at all* — the first such statement in the series. [**TODO** write; see §4.]

2 The biased invariant

Let P_q be the product measure under which each $a_i \in A$ is present independently with probability $q \in (0, 1)$, and write $\text{lm}_q(n)$, $\text{deg}_q(n)$ for the P_q -measures of the sets of strict local minima and of ground states at the target n . The target is taken at the q -tilted centre $n_q = \lfloor qT \rfloor$.

Definition 2.1 (biased gap series).

$$\Gamma^{(q)}(A) = 1 + \sum_{d \geq 1} \left[q^{N_d} + (1 - q)^{N_d} \right].$$

At $q = \frac{1}{2}$ every bracket is $2 \cdot 2^{-N_d}$ and $\Gamma^{(1/2)}$ is the window series W_D of paper 1, hence $\Gamma(A) + (2D + 1)2^{-k}$.

*Independent researcher.

Remark 2.2 (why the classification survives the deformation). Paper 1's classification is measure-free: at error d , an overshooting configuration is a strict local minimum exactly when every $a \leq 2d$ is *excluded*, and an undershooting one exactly when every $a \leq 2d$ is *included*. Under P_q those two events have probability $(1-q)^{N_d}$ and q^{N_d} , which is Definition 2.1. Writing $r_B^q(m) = \sum_{U \subseteq B, \sigma(U)=m} q^{|U|}(1-q)^{|B|-|U|}$ and factorising P_q across I_d and B_d ,

$$\text{over}_q = (1-q)^{N_d} r_{B_d}^q(n+d), \quad \text{under}_q = q^{N_d} r_{B_d}^q(n-d-\sigma_d), \quad \deg_q = \sum_{J \subseteq I_d} q^{|J|} (1-q)^{N_d-|J|} r_{B_d}^q(n-\sigma(J)),$$

and the binomial sum over J is exactly 1, so a flat r^q gives the d -th term of Definition 2.1. [\[derived; the assignment verified per stratum, see Remark 2.7\]](#)

2.1 The fair coin is the flattest

Lemma 2.3 (termwise minimality). *For every integer $N \geq 0$ the function $f_N(q) = q^N + (1-q)^N$ on $(0,1)$ is symmetric about $q = \frac{1}{2}$ and convex, and $f_N(q) \geq f_N(\frac{1}{2}) = 2^{1-N}$, with strict inequality for $q \neq \frac{1}{2}$ as soon as $N \geq 2$.*

Proof. $f_N(1-q) = f_N(q)$ is immediate. For $N \geq 2$, $f'_N(q) = N[q^{N-1} - (1-q)^{N-1}]$ and $f''_N(q) = N(N-1)[q^{N-2} + (1-q)^{N-2}] > 0$, so f_N is strictly convex; and $t \mapsto t^{N-1}$ is strictly increasing, so $f'_N < 0$ on $(0, \frac{1}{2})$ and $f'_N > 0$ on $(\frac{1}{2}, 1)$. For $N \in \{0,1\}$, f_N is the constant 2 or 1. $\square$

Theorem 2.4 (the fair coin minimises the invariant). *For every finite increasing sequence A of odd positive integers with $|A| \geq 2$ and every $q \in (0,1)$,*

$$\Gamma^{(q)}(A) \geq \Gamma^{(1/2)}(A),$$

with equality if and only if $q = \frac{1}{2}$.

Proof. Sum Lemma 2.3 over d . For strictness it suffices that some $d \leq D$ has $N_d \geq 2$: take any $d \geq (a_2 - 1)/2$, which is at most $D = (a_k - 1)/2$ because $a_2 \leq a_k$. $\square$

Remark 2.5 (what the theorem says). $q(1-q)$ is the per-element variance of the coin. Biasing the coin in either direction lowers that variance and raises the invariant: **the more predictable the coin, the more rugged the landscape**. The effect is exactly symmetric in $q \leftrightarrow 1-q$, as it must be, since complementing every configuration exchanges q with $1-q$ and n with $T-n$.

Corollary 2.6 (closed form for the odd numbers). *For $A = (1, 3, \dots, 2k-1)$ one has $N_d = d$ for $d \leq k$, so in the limit $k \rightarrow \infty$*

$$\Gamma^{(q)} = 1 + \frac{q}{1-q} + \frac{1-q}{q} = \frac{1}{q(1-q)} - 1 \geq 3,$$

with equality only at $q = \frac{1}{2}$: the invariant of the odd numbers is the reciprocal of the coin's variance, minus one. [\[verified against the summed series to \$5.3 \cdot 10^{-15}\$ over \$q = 0.10, \dots, 0.80\$; e5_r098\]](#)

2.2 Verification

[\[experimentally confirmed\]](#) Measured $\text{lm}_q / \deg_q$ at n_q against the point prediction of Definition 2.1, relative error:

profile	q	$k = 60$	$k = 100$	$k = 140$
odds	0.20	$1.00 \cdot 10^{-2}$	$3.16 \cdot 10^{-3}$	$1.49 \cdot 10^{-3}$
odds	0.30	$2.63 \cdot 10^{-3}$	$8.02 \cdot 10^{-4}$	$3.75 \cdot 10^{-4}$
odds	0.50	$2.73 \cdot 10^{-4}$	$5.99 \cdot 10^{-5}$	$2.19 \cdot 10^{-5}$
primes	0.30	$2.18 \cdot 10^{-3}$	$5.02 \cdot 10^{-4}$	$2.20 \cdot 10^{-4}$
primes	0.50	$1.73 \cdot 10^{-4}$	$2.26 \cdot 10^{-5}$	$7.32 \cdot 10^{-6}$

There are no free parameters. Two features: the error decays faster than $1/k$ (fitted exponent 2.25 at $q = 0.2$ to 3.0 at $q = 0.5$), and it grows monotonically as q leaves $\frac{1}{2}$, by a factor ≈ 40 from $q = 0.5$ to $q = 0.2$ at fixed k . The second is what the biased flatness hypothesis will have to quantify. **[TODO state the biased (H); design after §2.3.]**

Remark 2.7 (the assignment, and a test that could not decide it). Definition 2.1 is symmetric in $q \leftrightarrow 1 - q$, and so is the exact complementation identity $\text{lm}_q(n) = \text{lm}_{1-q}(T - n)$ — at the q -tilted centre, $T - qT = (1 - q)T$. Neither can therefore distinguish the assignment of q^{N_d} and $(1 - q)^{N_d}$ to the two strata: measured at $k = 100$, the ratio at $q = 0.4$ and at $q = 0.6$ agrees to every printed digit. The discriminating test is per stratum, $\text{over}_q/\text{deg}_q$ against $(1 - q)^{N_d}$ and $\text{under}_q/\text{deg}_q$ against q^{N_d} ; measured, both ratios lie in $[0.9978, 1.0018]$ over $d \leq 6$, whereas a swap would disagree by a factor of 161 at $d = 6$. **[experimentally confirmed; e5_r098]**

2.3 The bias breaks the reflection symmetry

[experimentally confirmed, shape] Off the q -tilted centre the deviation $\text{dev}_q = (\text{lm}_q/\text{deg}_q)/\Gamma^{(q)} - 1$ behaves differently at $q = \frac{1}{2}$ and away from it, and the reason is structural. Paper 3's transfer function is even in λ because the two strata sit at $\pm\delta_d$ about a common centre — the analytic form of $\text{lm}(n) = \text{lm}(T - n)$. Under bias that symmetry maps q to $1 - q$ rather than to itself, so the odd orders need not cancel, and the first-order coefficient is

$$L^{(q)}(A) = \sum_{d \geq 1} \delta_d [q^{N_d} - (1 - q)^{N_d}],$$

which is antisymmetric in $q \leftrightarrow 1 - q$ and vanishes identically at $q = \frac{1}{2}$. Measured for the odd numbers at $k = 80$: $L^{(q)} = -28.634, -8.843, 0, +8.843, +28.634$ at $q = 0.3, 0.4, 0.5, 0.6, 0.7$, and for the primes $-83.717, -23.677, 0, +23.677, +83.717$.

The data agree. Over the same range of targets, the spread (max over min) of $|\text{dev}_q - \text{dev}_q(0)|$ divided by λ_q and by λ_q^2 :

profile	q	divided by λ_q	divided by λ_q^2
odds	0.30	1.329	9.644
odds	0.50	5.091	1.001
primes	0.30	1.284	8.858

So the response is linear in λ under bias and quadratic at $q = \frac{1}{2}$, where $(\text{dev} - \text{dev}_0)/\lambda^2$ is constant to four digits (19.8186 to 19.8352 across a fivefold range of the target offset). **[experimentally confirmed; e6_r100]** **[TODO the primes at $q = \frac{1}{2}$ are not a clean test in this range: $\text{dev} - \text{dev}_0$ crosses zero inside it, and dropping the single point where it is $2.7 \cdot 10^{-7}$ leaves the quadratic ratio in $[45.06, 48.86]$. Re-measure on a target range that does not straddle the crossing.]**

3 The biased transfer function

[**TODO** fable-5: derive $\Phi^{(q)}$ by the partition-function method of paper 3’s rem:mushift, with the biased normaliser. The two strata carry unequal weights q^{N_d} and $(1-q)^{N_d}$ and a q -shifted μ_d , so the cosh becomes a general combination of two exponentials; §2.3 says the surviving odd part has coefficient $L^{(q)}$, which is the first thing the derivation must reproduce.]

4 Random sequences, unconditionally

[**TODO** fable-5: for random odd A the minor-arc input is a second-moment bound $\mathbb{E}|S_A|^2 = b$ plus Borel–Cantelli on a θ -net — no Kumchev, no Siegel–Walfisz, no Siegel constant. If it lands it is the trilogy’s first zero-ineffective-constant rigidity theorem. The modulus-4 floor of papers 1–2, which is universal for odd sequences, is then the exact leading ripple.]

5 Two questions inherited from paper 3

[**TODO** (i) the c_d fold-in: paper 3 keeps Φ as printed and counts $c_d = e^{-\delta_d^2/2K_d''}$ in E_k , on the measurement that folding it in accounts for at most 3% of the residual. Extend that measurement across profiles and k and decide. (ii) the mechanism behind $c_A = c'_A$, conjectured in paper 3’s rem:onek.]

6 Outlook

[**TODO** the algorithmic reading — local search on structured instances — which is where this programme started. Outlook only; no claims.]

7 Honest scope

- *Proved:* Lemma 2.3, Theorem 2.4, Corollary 2.6; and Remark 2.2’s derivation of Definition 2.1 modulo the biased flatness hypothesis, which is not yet stated.
- *Experimentally confirmed, with the range stated:* the point prediction (§2.2, two profiles, five q , three k), the stratum assignment (Remark 2.7), the linear response under bias (§2.3).
- *Not yet written:* §§3–6, and the biased hypothesis that §2 needs to become a theorem.

References

- [1] K. Amauchi, *The gap series of an integer sequence*. Manuscript, 2026.
- [2] K. Amauchi, *Asymptotic flatness of subset-sum landscapes of primes*. Manuscript, 2026.
- [3] K. Amauchi, *The transfer function of subset-sum landscapes*. Manuscript, 2026.