

HW: Custom Problem, Section 5A #19, 21

Sam Fleischer

March 17, 2015

Custom Problem

Let $T \in \mathcal{L}(\mathbb{C}^2)$ by $T(z, w) = (2w, -8z)$.

(a) Find a basis for \mathbb{C}^2 consisting of eigenvectors of T .

To find the eigenvalues, set $T(z, w) = \lambda(z, w)$.

$$\begin{aligned} T(z, w) &= \lambda(z, w) \\ \implies (2w, -8z) &= (\lambda z, \lambda w) \\ \implies 2w &= \lambda z, \quad \text{and} \quad -8z = \lambda w \\ \implies -8z &= \frac{\lambda^2}{2}z \end{aligned}$$

If $z = 0$, then $w = 0$. Then λ would not be an eigenvalue. So we suppose $z \neq 0$, and thus

$$\begin{aligned} -8 &= \frac{\lambda^2}{2} \\ \implies -16 &= \lambda^2 \\ \implies \lambda &= \pm 4i \end{aligned}$$

Thus the two eigenvalues for T are $4i$ and $-4i$. To find their eigenspaces, set $T(z, w) = 4i(z, w)$ and $T(z, w) = -4i(z, w)$, respectively. So,

$$\begin{aligned} T(z, w) &= 4i(z, w) \\ \implies (2w, -8z) &= (4iz, 4iw) \\ \implies w &= 2iz \end{aligned}$$

Thus $(1, 2i)$ is an eigenvector corresponding to $4i$. Furthermore, $(1, 2i)$ is a basis for $E(T, 4i)$. Also,

$$\begin{aligned} T(z, w) &= -4i(z, w) \\ \implies (2w, -8z) &= (-4iz, -4iw) \\ \implies w &= -2iz \end{aligned}$$

Thus $(1, -2i)$ is an eigenvector corresponding to $-4i$. Furthermore, $(1, -2i)$ is a basis for $E(T, -4i)$. Then $\pi = ((1, 2i), (1, -2i))$ is a linearly independent set since eigenvectors corresponding to different eigenvalues are linearly independent. Since $\dim(\mathbb{C}^2) = 2$ and $\text{len}(\pi) = 2$, π is a basis for \mathbb{C}^2 consisting of eigenvectors of T .

(b) Find $\mathcal{M}(T)$ with respect to this basis.

Since

$$\begin{aligned} T(1, 2i) &= (2(2i), -8(1)) \\ &= (4i, -8) \\ &= 4i(1, 2i) + 0(1, -2i) \end{aligned}$$

and

$$\begin{aligned} T(1, -2i) &= (2(-2i), -8(1)) \\ &= (-4i, -8) \\ &= 0(1, 2i) - 4i(1, -2i) \end{aligned}$$

then

$$\mathcal{M}(T, \pi) = \begin{pmatrix} 4i & 0 \\ 0 & -4i \end{pmatrix}$$

5A

#19

Suppose n is a positive integer and $T \in \mathcal{L}(\mathbb{F}^n)$ is defined by

$$T(x_1, \dots, x_n) = (x_1 + \dots + x_n, \dots, x_1 + \dots + x_n)$$

in other words, T is the operator whose matrix (with respect to the standard basis) consists of all 1's. Find the eigenvalues and eigenvectors of T .

To find the eigenvalues, set $T(x_1, \dots, x_n) = \lambda(x_1, \dots, x_n)$.

$$\begin{aligned} T(x_1, \dots, x_n) &= \lambda(x_1, \dots, x_n) \\ \implies (x_1 + \dots + x_n, \dots, x_1 + \dots + x_n) &= (\lambda x_1, \dots, \lambda x_n) \\ \implies x_1 + \dots + x_n &= \lambda x_1 = \dots = \lambda x_n \\ \implies \lambda &= \frac{x_1 + \dots + x_n}{x_1} = \dots = \frac{x_1 + \dots + x_n}{x_n} \end{aligned}$$

If $\lambda \neq 0$, then $x_1 = \dots = x_n$, and thus $(1, \dots, 1)$ is a basis for $E(T, \lambda)$. However, if $\lambda = 0$, then $x_1 + \dots + x_n = 0$. Then $E(T, 0) = \{(x_1, \dots, x_n) \in \mathbb{F}^n \mid x_1 + \dots + x_n = 0\}$. Note $\dim(E(T, 0)) = n - 1$ and a basis for $E(T, 0)$ is

$$\pi = ((1, -1, 0, \dots, 0), (1, 0, -1, 0, \dots, 0), \dots, (1, 0, \dots, 0, -1))$$

#21

Suppose $T \in \mathcal{L}(V)$ is invertible.

- (a) Suppose $\lambda \in \mathbb{F}$ with $\lambda \neq 0$. Prove that λ is an eigenvalue of T if and only if $\frac{1}{\lambda}$ is an eigenvalue of T^{-1} .

$$\begin{aligned} & \lambda \text{ is an eigenvalue of } T \\ \iff & \exists v \in V \text{ such that } T(v) = \lambda v \\ \iff & T^{-1}(T(v)) = T^{-1}(\lambda v) \\ \iff & v = \lambda T^{-1}(v) \\ \iff & T^{-1}(v) = \frac{1}{\lambda} v \text{ for some } v \in V \\ \iff & \frac{1}{\lambda} \text{ is an eigenvalue of } T^{-1} \end{aligned}$$

□

- (b) Prove that T and T^{-1} have the same eigenvectors.

Let \hat{v} be an eigenvector of T corresponding to an arbitrary eigenvalue of T 's, say $\hat{\lambda}$. Then

$$\begin{aligned} & T(\hat{v}) = \hat{\lambda}\hat{v} \\ \implies & T^{-1}(T(\hat{v})) = T^{-1}(\hat{\lambda}\hat{v}) \\ \implies & \hat{v} = \hat{\lambda}T^{-1}(\hat{v}) \\ \implies & T^{-1}(\hat{v}) = \frac{1}{\hat{\lambda}}\hat{v} \end{aligned}$$

Thus \hat{v} is an eigenvector of T^{-1} corresponding to $\frac{1}{\hat{\lambda}}$. Thus any eigenvector of T is an eigenvector of T^{-1} . However, the same argument and $(T^{-1})^{-1} = T$ implies any eigenvector of T^{-1} is an eigenvector of T . Thus T and T^{-1} have the same eigenvectors. □