

HW #6

Sam Fleischer

November 16, 2015

Problem 1

Let $k : [0, 1] \times [0, 1] \rightarrow \mathbb{R}$ be a continuous function. Define the map $T : C([0, 1]) \rightarrow C([0, 1])$ by

$$(Tf)(x) = \int_0^1 k(x, y)f(y)dy, \quad \text{for all } f \in C([0, 1])$$

a)

Let $\|T\|$ denote the operator norm of T . Prove

$$\|T\| = \sup_{x \in [0, 1]} \int_0^1 |k(x, y)|dy \quad (1)$$

First, we show $\|T\| \leq \sup_{x \in [0, 1]} \int_0^1 |k(x, y)|dy$.

$$\begin{aligned} \|T\| &= \sup_{\|f\|=1} \|Tf\| = \sup_{\|f\|=1} \left\| \int_0^1 k(\cdot, y)f(y)dy \right\| \\ &= \sup_{x \in [0, 1]} \left(\sup_{\|f\|=1} \left| \int_0^1 k(x, y)f(y)dy \right| \right) \\ &\leq \sup_{x \in [0, 1]} \left(\sup_{\|f\|=1} \int_0^1 |k(x, y)||f(y)|dy \right) \\ &\leq \sup_{x \in [0, 1]} \left(\sup_{\|f\|=1} \int_0^1 |k(x, y)||f|dy \right) \\ &= \sup_{x \in [0, 1]} \int_0^1 |k(x, y)|dy \end{aligned}$$

Note that since $\int_0^1 |k(x, y)|dy$ is a continuous functions of x , then $\exists x^* \in [0, 1]$ such that

$$\sup_{x \in [0, 1]} \int_0^1 |k(x, y)|dy = \int_0^1 |k(x^*, y)|dy$$

Thus,

$$\|T\| \leq \int_0^1 |k(x^*, y)| dy$$

Next, we show $\|T\| \geq \int_0^1 |k(x^*, y)| dy$. Define a sequence of functions $(f_n)_n$

$$f_n(y) = \frac{k(x^*, y)}{\frac{1}{n} + |k(x^*, y)|}$$

Note $f_n(y) \rightarrow \text{sign}(k(x^*, y))$ for each $y \in [0, 1]$, i.e. $f_n \rightarrow \text{sign}(k(x^*, \cdot))$ pointwise.

$$\begin{aligned} \sup_{x \in [0, 1]} \int_0^1 |k(x, y)| dy &= \int_0^1 |k(x^*, y)| dy \\ &= \int_0^1 k(x^*, y) \text{sign}(k(x^*, y)) dy \\ &= \int_0^1 k(x^*, y) \lim_{n \rightarrow \infty} f_n(y) dy \end{aligned}$$

Since $\|f_n\|_\infty \leq 1$ for all n , we can employ Lebesgue's Dominated Convergence Theorem to pull the limit out of the integral:

$$\int_0^1 k(x^*, y) \lim_{n \rightarrow \infty} f_n(y) dy = \lim_{n \rightarrow \infty} \int_0^1 k(x^*, y) f_n(y) dy$$

Thus,

$$\begin{aligned} \sup_{x \in [0, 1]} \int_0^1 |k(x, y)| dy &= \lim_{n \rightarrow \infty} \int_0^1 k(x^*, y) f_n(y) dy \\ &= \lim_{n \rightarrow \infty} (Tf_n)(x^*) \\ &\leq \lim_{n \rightarrow \infty} \|Tf_n\|_\infty \end{aligned}$$

But since $\|f_n\| \leq 1$ for all n ,

$$\sup_{x \in [0, 1]} \int_0^1 |k(x, y)| dy \leq \lim_{n \rightarrow \infty} \|Tf_n\|_\infty \leq \lim_{n \rightarrow \infty} \sup_{\|f\| \leq 1} \|Tf\| = \lim_{n \rightarrow \infty} \|T\| = \|T\|$$

Thus,

$$\|T\| \leq \sup_{x \in [0, 1]} \int_0^1 |k(x, y)| dy \leq \|T\| \implies \|T\| = \sup_{x \in [0, 1]} \int_0^1 |k(x, y)| dy$$

□

b)

Argue that the sup in (1) is attained in some $x \in [0, 1]$.

As argued in part **a)**, since k is a continuous function of x and y , then $\int_0^1 |k(x, y)| dy$ is a continuous function of x . Thus $\int_0^1 |k(x, y)| dy$ must reach its maximum on a compact set. Since $[0, 1]$ is compact, $\exists x^* \in [0, 1]$ such that $\sup_{x \in [0, 1]} \int_0^1 |k(x, y)| dy = \int_0^1 |k(x^*, y)| dy$.

c)

Is it possible that $\|T\| = 1$ but $\|T^2\| = 0$? Prove your answer.

Define $k(x, y) : [0, 1] \times [0, 1] \rightarrow \mathbb{R}$ as

$$k(x, y) = \begin{cases} 0, & \text{if } y \leq x + \frac{1}{2} \\ 4(2y - 2x - 1), & \text{else} \end{cases}$$

Then if T is defined using k , then

$$\begin{aligned} \|T\| &= \sup_{x \in [0, 1]} \int_0^1 |k(x, y)| dy \\ &= 4 \sup_{x \in [0, 1]} \int_{x + \frac{1}{2}}^1 |2y - 2x - 1| dy \\ &= 4 \sup_{x \in [0, 1]} \left(y^2 - (2x + 1)y \right) \Big|_{x + \frac{1}{2}}^1 \\ &= 4 \sup_{x \in [0, 1]} \left(x - \frac{1}{2} \right)^2 = 4 \cdot \frac{1}{4} = 1 \end{aligned}$$

However,

$$\begin{aligned} \|T^2\| &= \sup_{\|f\|=1} \|T^2 f\| \\ &= \sup_{\|f\|=1} \int_0^1 k(x, y) \left(\int_0^1 k(y, s) f(s) ds \right) dy \\ &= \sup_{\|f\|=1} \int_0^1 \int_0^1 k(x, y) k(y, s) f(s) ds dy \end{aligned}$$

However, if $y \leq \frac{1}{2} \implies k(x, y) = 0$ and $y \geq \frac{1}{2} \implies k(y, s) = 0$, and thus $\forall y \in [0, 1]$, $k(x, y)k(y, s) = 0$. Thus,

$$\begin{aligned} \|T^2\| &= \sup_{\|f\|=1} \int_0^1 \int_0^1 k(x, y) k(y, s) f(s) ds dy \\ &= \sup_{\|f\|=1} \int_0^1 \int_0^1 0 ds dy = 0 \end{aligned}$$

So it is possible for $\|T\| = 1$ and $\|T^2\| = 0$. □

Problem 2

Study Section 5.4 of the textbook.

Problem 3

Let X be the Banach space $\ell^2(\mathbb{N})$, defined by

$$\ell^2(\mathbb{N}) = \left\{ z = (z_n)_{n=1}^{\infty} \mid z_n \in \mathbb{C}, \sum_{n=1}^{\infty} |z_n|^2 < \infty \right\}$$

For $m = 1, 2, \dots$, define $e_m \in X$ to be the sequence with elements $(e_m)_n = \delta_{n,m}$, and define $P_m : X \rightarrow X$ by $P_m z = z_m e_m$, for all $z \in X$.

a)

Prove that $P_m \in \mathcal{B}(X)$, for all $m \geq 1$.

$$\|P_m\| = \sup_{\|z\|=1} \|P_m z\| = \sup_{\|z\|=1} \|z_m e_m\| = \sup_{\|z\|=1} \|z_m\| \|e_m\| \leq \|e_m\|$$

Thus P_m is a bounded linear operator on X , i.e $P_m \in \mathcal{B}(X)$. □

b)

Verify $P_m P_n = \delta_{n,m} P_m$, for all $n, m \geq 1$.

$$(P_m P_n)(z) = P_m(P_n z) = P_m(z_n e_n) = z_n P_m(e_n) = z_n (e_{n,m}) e_m = z_n \delta_{n,m} e_m$$

If $n \neq m$, then $\delta_{n,m} = 0$ and

$$(P_m P_n)(z) = z_n \delta_{n,m} e_m = 0 = 0 P_m(z) = \delta_{n,m} P_m(z) = (\delta_{n,m} P_m)(z)$$

If $n = m$, then $z_m = z_n$ and

$$(P_m P_n)(z) = z_n \delta_{n,m} e_m = \delta_{n,m} z_m e_m = \delta_{n,m} P_m(z) = (\delta_{n,m} P_m)(z)$$

In either case, $P_m P_n = \delta_{n,m} P_m$. □

c)

Prove that $\|P_m\| = 1$, for all $m \geq 1$.

d)

For $m \geq 1$, define S_m by

$$S_m = \sum_{k=1}^m P_k$$

Calculate $S_m S_n$, for all $n, m \geq 1$.

e)

Show that $\|S_m\| = 1$, for all $n \geq 1$.