

HW #8

Sam Fleischer

November 30, 2015

Problem 1

Consider the Banach space $C([0, 1])$ with the supremum norm. For $x \in [0, 1]$ let δ_x denote the linear functional on $C([0, 1])$ given by

$$\delta_x(f) = f(x), \quad \text{for all } f \in C([0, 1])$$

a)

Show that $\|\delta_x\| = 1$.

b)

Show that there does not exist a Riemann integrable function $k : [0, 1] \rightarrow \mathbb{R}$, such that

$$\delta_x(f) = \int_0^1 k(y)f(y)dy, \quad \text{for all } f \in C([0, 1])$$

Problem 2

Prove that there does not exist an inner product on $C([0, 1])$ such that the supremum norm is derived from this inner product.

Problem 3

Let \mathcal{H} be a Hilbert space and let M be a subset of \mathcal{H} .

a)

Prove that M^\perp is a closed linear subspace of \mathcal{H} .

First we show M^\perp is a linear subspace. Let $x, y \in M^\perp$ and $\lambda, \mu \in \mathbb{C}$. Then for each $m \in M$,

$$\langle m, \lambda x + \mu y \rangle = \lambda \langle m, x \rangle + \mu \langle m, y \rangle = 0$$

Thus $\lambda x + \mu y \in M^\perp$. Thus M^\perp is a linear subspace of \mathcal{H} . Next, let (x_n) be a convergent sequence in M^\perp , and let $x_n \rightarrow x$. Then for each $m \in M$,

$$\langle x, m \rangle = \langle \lim_{n \rightarrow \infty} x_n, m \rangle$$

but since $\langle \cdot, \cdot \rangle$ is continuous,

$$\langle \lim_{n \rightarrow \infty} x_n, m \rangle = \lim_{n \rightarrow \infty} \langle x_n, m \rangle = \lim_{n \rightarrow \infty} 0 = 0$$

Thus $\langle x, m \rangle = 0$, which shows $x \in M^\perp$, proving M^\perp is closed. \square

b)

Prove that $M \cap M^\perp \subset \{0\}$.

Let $x \in M \cap M^\perp$. Then by the definition of M^\perp ,

$$\langle x, x \rangle = 0$$

Then $\|x\| = 0$, which shows $x = 0$. Thus $M \cap M^\perp \subset \{0\}$. \square

c)

If M is a linear subspace of \mathcal{H} , prove that $(M^\perp)^\perp = \overline{M}$.

Assume $x \in \overline{M}$. Then there is a sequence $x_n \in M$ such that $x_n \rightarrow x$. Then $\langle x_n, y \rangle = 0$ for every $y \in M^\perp$. Then by continuity of $\langle \cdot, \cdot \rangle$, $\langle x, y \rangle = 0$ for every $y \in M^\perp$. Then $x \in (M^\perp)^\perp$ by the definition of $(M^\perp)^\perp$. Thus $\overline{M} \subset (M^\perp)^\perp$.

Now assume $x \notin \overline{M}$. Since \overline{M} is closed, then by the Projection Theorem, $\exists y \in \overline{M}$ such that $(x - y) \perp \overline{M}$. Since $y \in \overline{M}$, $\langle x - y, y \rangle = 0$. Since $x \neq y$ ($x \notin \overline{M}$ and $y \in \overline{M}$), then $\langle x - y, x - y \rangle \neq 0$. However, $\langle x - y, x - y \rangle = \langle x - y, x \rangle - \langle x - y, y \rangle = \langle x - y, x \rangle$. Since $x - y \perp \overline{M}$, then $x - y \perp M$. So $x - y \in M^\perp$. Then since $\langle x - y, x \rangle \neq 0$, then $x \notin (M^\perp)^\perp$. Then $(M^\perp)^\perp \subset \overline{M}$.

Thus, $\overline{M} = (M^\perp)^\perp$. \square

Problem 4

Let \mathcal{H} be a Hilbert space and $A \in \mathcal{B}(\mathcal{H})$. If $\langle x, Ay \rangle = 0$ for all $x, y \in \mathcal{H}$, prove $A = 0$.

Since $\langle x, Ay \rangle = 0$ for all $x, y \in \mathcal{H}$, then in particular, take $x = Ay$, and so $\langle Ay, Ay \rangle = 0$ for all $y \in \mathcal{H}$. Then $Ay = 0$ for all $y \in \mathcal{H}$. Thus $A = 0$. \square

Problem 5

Let \mathcal{H} be a Hilbert space and P and Q two orthogonal projections on \mathcal{H} .

a)

Prove that PQ is an orthogonal projection if and only if $PQ - QP = 0$, i.e., if and only if P and Q commute.

First note that

$$\langle PQx, y \rangle = \langle Qx, Py \rangle = \langle x, QPy \rangle \quad (1)$$

Assume PQ is an orthogonal projection. Then by the definition of orthogonal projection, and by (1), $\langle PQx, y \rangle = \langle x, PQy \rangle$. Then for all $x, y \in \mathcal{H}$,

$$\begin{aligned} \langle x, QPy \rangle &= \langle x, PQy \rangle \\ \implies \langle x, (QP - PQ)y \rangle &= 0 \\ \implies QP - PQ &= 0 \\ \implies QP &= PQ \end{aligned}$$

Thus P and Q commute.

Now assume $PQ = QP$. Then $(PQ)^2 = PQPQ = PPQQ = PQ$ since P and Q are orthogonal projections. Also, by (1), $\langle PQx, y \rangle = \langle x, QPy \rangle = \langle x, PQy \rangle$. Thus PQ is an orthogonal projection. \square

b)

Prove that for commuting orthogonal projections P and Q , one has $\text{ran}(PQ) = \text{ran}(P) \cap \text{ran}(Q)$.

Let $x \in \text{ran}(PQ)$. Then $\exists y$ such that $PQy = x$. Then P maps Qy on to x . Then $x \in \text{ran}(P)$. However, since P and Q commute, then $QPy = x$, and thus Q maps Py on to x , and so $x \in \text{ran}(Q)$. Thus $x \in \text{ran}(P) \cap \text{ran}(Q)$. So $\text{ran}(PQ) \subset \text{ran}(P) \cap \text{ran}(Q)$.

Now let $x \in \text{ran}(P) \cap \text{ran}(Q)$. Then $x \in \text{ran}(P)$ and $x \in \text{ran}(Q)$. So $\exists y_1, y_2$ such that $Py_1 = Qy_2 = x$. Thus, $PQy_2 = P^2y_1 = Py_1 = x$, and thus $x \in \text{ran}(PQ)$. So $\text{ran}(P) \cap \text{ran}(Q) \subset \text{ran}(PQ)$.

Thus, $\text{ran}(PQ) = \text{ran}(P) \cap \text{ran}(Q)$. \square

c)

Prove that $P + Q$ is an orthogonal projection if and only if $PQ = 0$.

Assume $PQ = 0$. Then $\langle PQx, y \rangle = 0$ for all $x, y \in \mathcal{H}$. But by (1), $\langle x, QPy \rangle = 0$ for all $x, y \in \mathcal{H}$. Thus $QP = 0$. Then $(P + Q)^2 = P^2 + PQ + QP + Q^2 = P^2 + 0 + 0 + Q^2 = P + Q$ since P and Q are orthogonal projections. Also,

$$\begin{aligned} \langle (P + Q)x, y \rangle &= \langle Px + Qx, y \rangle \\ &= \langle Px, y \rangle + \langle Qx, y \rangle \\ &= \langle x, Py \rangle + \langle x, Qy \rangle \\ &= \langle x, Py + Qy \rangle \\ &= \langle x, (P + Q)y \rangle \end{aligned}$$

Thus $P + Q$ is an orthogonal projection.

Assume $P + Q$ is an orthogonal projection. Then $(P + Q)^2 = P + Q$, but $(P + Q)^2 = P^2 + PQ + QP + Q^2 = P + PQ + QP + Q$. Thus $PQ + QP = 0$, i.e. $PQ = -QP$.

Assume $x \in \text{ran}(P) \cap \text{ran}(Q)$ and note $0 = (PQ + QP)x = PQx + QPx$. Since $x \in \text{ran}(P)$, $Px = x$. Also, since $x \in \text{ran}(Q)$, $Qx = x$. Then $PQx = Px = x$ and $QPx = Qx = x$. So $0 = PQx + QPx = 2x$. Thus $x = 0$.

Now, take any $x \in \mathcal{H}$, then certainly $PQx \in \text{ran}(P)$ and since $PQx = -QPx = Q(-Px)$, then $PQx \in \text{ran}(Q)$. Then $PQx = 0$ by the paragraph above, and thus $PQ = 0$, i.e. $\text{ran}(PQ) = \{0\}$.

Thus, $P + Q$ is an orthogonal projection if and only if $\text{ran}(PQ) = \{0\}$. \square

d)

Prove that if $PQ = 0$, we have $\text{ran}(P + Q) = \text{ran}(P) \oplus \text{ran}(Q)$.

Let $PQ = 0$ and assume $y \in \text{ran}(P + Q)$. Then $\exists x \in \mathcal{H}$ such that $Px + Qx = y$. Then y is the sum of an element in $\text{ran}(P)$ and an element in $\text{ran}(Q)$. Thus $y \in \text{ran}(P) \oplus \text{ran}(Q)$. Assume $y \in \text{ran}(P) \oplus \text{ran}(Q)$. Then $\exists x_1, x_2 \in \mathcal{H}$ such that $y = Px_1 + Qx_2$. Then $Py = P^2x_1 + PQx_2 = Px_1$ and $Qy = QPx_1 + Q^2x_2 = Qx_2$. Thus $y = Px_1 + Qx_2 = Py + Qy = (P + Q)y$. Thus $y \in \text{ran}(P + Q)$, which shows $\text{ran}(P + Q) = \text{ran}(P) \oplus \text{ran}(Q)$. \square

Problem 6

Let \mathcal{H} be a Hilbert space and $P \in \mathcal{B}(\mathcal{H})$ such that $P^2 = P$ and $\dim \text{ran}(P) = 1$.

a)

Show that $\|P\| \geq 1$.

Let $x \in \text{ran}(P)$ such that $\|x\| = 1$. Then $Px = x$, and so $\|Px\| = \|x\| = 1$. Thus $\|P\| \geq 1$. \square

b)

Suppose $\dim \mathcal{H} \geq 2$. Find

$$\sup \{ \|P\| \mid P \in \mathcal{B}(\mathcal{H}), P^2 = P, \dim \text{ran}(P) = 1 \}$$

Problem 7

Let $\{e_1, e_2, \dots\}$ be an orthonormal basis of a separable Hilbert space \mathcal{H} .

a)

Let $(a_n) \in \ell^1(\mathbb{N})$. Show that $\sum_{n=1}^{\infty} a_n e_n$ converges absolutely to a limit in \mathcal{H} .

b)

Let $\alpha \in (0, \infty)$ and define $a_n = n^{-\alpha}$, $n \geq 1$. For which values of α does $\sum_{n=1}^{\infty} a_n e_n$ converge unconditionally but not absolutely?

Problem 8

Define the Legendre polynomials P_n by

$$P_n(x) = \frac{1}{2^n n!} \frac{d^n}{dx^n} (x^2 - 1)^n$$

a)

Show that the Legendre polynomials are orthogonal in $L^2([-1, 1])$, and that they are obtained by Gram-Schmidt orthogonalization of the monomials.

b)

Show that

$$\int_{-1}^1 P_n(x)^2 dx = \frac{2}{2n+1}$$

c)

Prove that the Legendre polynomials form an orthogonal basis of $L^2([-1, 1])$.

d)

Prove that the Legendre polynomial P_n is an eigenfunction of the differential operator

$$L = -\frac{d}{dx}(1-x^2)\frac{d}{dx}$$

with eigenvalue $\lambda_n = n(n+1)$, meaning that

$$LP_n = \lambda_n P_n$$