

HW #2

Sam Fleischer

January 29, 2015

Hunter and Nachtergaele 7.1

Let ϕ_n be the functions defined in (7.7)

$$\phi_n(x) = c_n(1 + \cos x)^n$$

where c_n is chosen such that

$$\int_{\mathbb{T}} \phi_n(x) dx = 1$$

for all n .

(a) Prove (7.5).

$$\lim_{n \rightarrow \infty} \int_{\delta \leq |x| \leq \pi} \phi_n(x) dx = 0$$

for every $\delta > 0$.

Let $\delta > 0$ and for ease, define $\mathbb{D} = [-\pi, -\delta] \cup [\delta, \pi]$.

$$\int_{\mathbb{D}} \phi_n(x) dx = \frac{\int_{\mathbb{D}} (1 + \cos x)^n dx}{\int_{\mathbb{T}} (1 + \cos x)^n dx}$$

since

$$c_n = \frac{1}{\int_{\mathbb{T}} (1 + \cos x)^n dx}$$

Note that

$$\phi'_n(x) = -nc_n(1 + \cos x)^{n-1} \sin x$$

which is positive on $[-\pi, 0)$ and negative on $(0, \pi]$, and thus

$$\max_{x \in \mathbb{D}} \phi_n(x) = \phi_n(\delta)$$

So,

$$\int_{\mathbb{D}} \phi_n(x) dx = \frac{\int_{\mathbb{D}} (1 + \cos x)^n dx}{\int_{\mathbb{T}} (1 + \cos x)^n dx} \leq \frac{2\pi(1 + \cos \delta)^n}{\int_{\mathbb{T}} (1 + \cos x)^n dx}$$

where $\mathbb{E} = [-\frac{\delta}{2}, \frac{\delta}{2}]$. Again, since ϕ_n is decreasing on $(0, \frac{\pi}{2}]$ and ϕ is an even function,

$$\min_{x \in \mathbb{E}} \phi_n(x) = \phi_n\left(\frac{\delta}{2}\right)$$

Thus,

$$\int_{\mathbb{D}} \phi_n(x) dx \leq \frac{2\pi(1 + \cos \delta)^n}{\int_{\mathbb{E}} (1 + \cos x)^n dx} \leq \frac{2\pi}{\delta} \left(\frac{1 + \cos \delta}{1 + \cos \frac{\delta}{2}} \right)^n$$

but

$$\frac{1 + \cos \delta}{1 + \cos \frac{\delta}{2}} < 1$$

since \cos is a decreasing function on $[0, \pi]$. Thus,

$$\lim_{n \rightarrow \infty} \frac{2\pi}{\delta} \left(\frac{1 + \cos \delta}{1 + \cos \frac{\delta}{2}} \right)^n = 0$$

and by the comparison test,

$$\lim_{n \rightarrow \infty} \int_{\mathbb{D}} \phi_n(x) dx = 0$$

- (b) *Prove that if the set \mathcal{P} of trigonometric polynomials is dense in the space of periodic continuous functions on \mathbb{T} with the uniform norm, then \mathcal{P} is dense in the space of all continuous functions on \mathbb{T} with the L^2 -norm.*
- (c) *Is \mathcal{P} dense in the space of all continuous functions on $[0, 2\pi]$ with the uniform norm?*

Hunter and Nachtergaele 7.2

Suppose that $f : \mathbb{T} \rightarrow \mathbb{C}$ is a continuous function, and

$$S_N = \frac{1}{\sqrt{2\pi}} \sum_{n=-N}^N \hat{f}_n e^{inx}$$

is the N^{th} partial sum of its Fourier series.

- (a) *Show that $S_N = D_N * f$, where D_N is the Dirichlet kernel*

$$D_N(x) = \frac{1}{2\pi} \frac{\sin \left[(N + \frac{1}{2})x \right]}{\sin \left(\frac{x}{2} \right)}.$$

- (b) *Let T_N be the mean of the first $N + 1$ partial sums,*

$$T_N = \frac{1}{N + 1}.$$

*Show that $T_N = F_N * f$, where F_N is the Fejér kernel*

$$F_N(x) = \frac{1}{2\pi(N + 1)} \left(\frac{\sin \left[(N + 1)\frac{x}{2} \right]}{\sin \left(\frac{x}{2} \right)} \right)^2.$$

- (c) Which of the families (D_N) and (F_N) are approximate identities as $N \rightarrow \infty$? What can you say about the uniform convergence of the partial sums S_N and the averaged partial sums T_N to f ?

Hunter and NachterGaele 7.3

Prove that the sets $\{e_n \mid n \geq 1\}$ defined by

$$e_n(x) = \sqrt{\frac{2}{\pi}} \sin nx,$$

and $\{f_n : n \geq 1\}$ defined by

$$f_0(x) = \sqrt{\frac{1}{\pi}}, \quad f_n(x) = \sqrt{\frac{2}{\pi}} \cos nx \quad \text{for } n \geq 1,$$

are both orthonormal bases of $L^2([0, \pi])$.

First we show $\{e_n\}_{n=1}^{\infty}$ and $\{f_n\}_{n=0}^{\infty}$ are orthonormal. Suppose $n \neq m$. Then

$$\begin{aligned} \langle e_n, e_m \rangle &= \int_0^{\pi} e_n(x) e_m(x) dx \\ &= \frac{2}{\pi} \int_0^{\pi} \sin(nx) \sin(mx) dx \\ &= \frac{2}{\pi} \int_0^{\pi} \frac{1}{2} [\cos(nx - mx) - \cos(nx + mx)] dx \\ &= \frac{1}{\pi} \int_0^{\pi} \cos(x(n - m)) dx - \frac{1}{\pi} \int_0^{\pi} \cos(x(n + m)) dx \\ &= \frac{1}{\pi} \left[\frac{\sin(x(n - m))}{n - m} - \frac{\sin(x(n + m))}{n + m} \right]_0^{\pi} \\ &= 0 \end{aligned}$$

Also,

$$\begin{aligned} \langle e_n, e_n \rangle &= \frac{2}{\pi} \int_0^{\pi} \sin^2(nx) dx \\ &= \frac{1}{\pi} \int_0^{\pi} 1 - \cos(2nx) dx \\ &= \frac{1}{\pi} \left[\pi - \frac{1}{2n} \sin(2n\pi) \right] \end{aligned}$$

Let $f \in L^2([0, 1])$. Then extend f to its odd expansion $\tilde{f} \in L^2([-1, 1])$ by

$$\tilde{f}(x) = \begin{cases} f(x) & \text{if } x \in (0, 1) \\ 0 & \text{if } x = 0 \\ -f(x) & \text{if } x \in (-1, 0) \end{cases}$$

Then $\{e_n\}_{n=1}^{\infty} \cup \{f_n\}_{n=0}^{\infty}$.

Hunter and Nachtergaele 7.4

Let $T, S \in L^2(\mathbb{T})$ be the triangular and square wave, respectively, defined by

$$T(x) = |x|, \quad \text{if } |x| \leq \pi, \quad S(x) = \begin{cases} 1 & \text{if } 0 < x < \pi \\ -1 & \text{if } -\pi < x < 0 \end{cases}$$

(a) Compute the Fourier series of T and S .

Since T is an even function, we can represent T with a cosine series

$$T(x) = \frac{1}{2}\hat{T}_0 + \sum_{n=1}^{\infty} \hat{T}_n \cos(nx)$$

where

$$\begin{aligned} \hat{T}_0 &= \frac{1}{\pi} \int_{\mathbb{T}} T(x) dx \quad \text{and} \\ \hat{T}_n &= \frac{1}{\pi} \int_{\mathbb{T}} T(x) \cos(nx) dx, \quad n = 1, 2, \dots \end{aligned}$$

Because \cos is even and T is even, $T \sin$ is even, and so

$$\hat{T}_0 = \frac{1}{\pi} \int_{\mathbb{T}} T(x) dx = \frac{2}{\pi} \int_0^{\pi} x dx = \frac{2}{\pi} \frac{\pi^2}{2} = \pi$$

and for $n = 1, 2, \dots$,

$$\hat{T}_n = \frac{2}{\pi} \int_0^{\pi} x \cos(nx) dx$$

Utilizing integration by parts, we find

$$\begin{aligned} \hat{T}_n &= \frac{2}{\pi} \int_0^{\pi} x \cos(nx) dx \\ &= \frac{2}{\pi} \left[\left(\frac{x}{n} \sin(nx) \right) \Big|_0^{\pi} - \frac{1}{n} \int_0^{\pi} \sin(nx) dx \right] \\ &= \frac{2}{\pi} \left[\frac{1}{n^2} \cos(nx) \right]_0^{\pi} \\ &= \frac{2}{\pi} \left[\frac{(-1)^n - 1}{n^2} \right] \\ &= \begin{cases} 0 & \text{if } n \text{ is even} \\ -\frac{4}{\pi n^2} & \text{if } n \text{ is odd} \end{cases} \end{aligned}$$

Thus,

$$T(x) = \frac{\pi}{2} - \frac{4}{\pi} \sum_{n=1}^{\infty} \left[\frac{1}{(2n-1)^2} \cos((2n-1)x) \right]$$

Since S is an odd function, we can represent S with a sin series

$$S(x) = \sum_{n=1}^{\infty} \hat{S}_n \sin(nx)$$

where

$$\hat{S}_n = \frac{1}{\pi} \int_{\mathbb{T}} S(x) \sin(nx) dx, \quad n = 1, 2, \dots$$

Because \sin is odd and S is odd, $\sin S$ is even, and thus

$$\begin{aligned} \hat{S}_n &= \frac{1}{\pi} \int_{\mathbb{T}} S(x) \sin(nx) dx \\ &= \frac{2}{\pi} \int_0^{\pi} \sin(nx) dx \\ &= \frac{2}{\pi} \left[-\frac{1}{n} \cos(nx) \right]_0^{\pi} \\ &= -\frac{2}{\pi n} ((-1)^n - 1) \\ &= \begin{cases} 0 & \text{if } n \text{ is even} \\ \frac{4}{\pi n} & \text{if } n \text{ is odd} \end{cases} \end{aligned}$$

Thus,

$$S(x) = \frac{4}{\pi} \sum_{n=1}^{\infty} \left[\frac{1}{(2n-1)} \sin((2n-1)x) \right]$$

(b) Show that $T \in H^1(\mathbb{T})$ and $T' = S$.

(c) Show that $S \notin H^1(\mathbb{T})$.

Hunter and Nachtergaele 7.5

Consider $f : \mathbb{T}^d \rightarrow \mathbb{C}$ defined by

$$f(x) = \sum_{n \in \mathbb{Z}^d} a_n e^{in \cdot x},$$

where $x = (x_1, x_2, \dots, x_d)$, $n = (n_1, n_2, \dots, n_d)$, and $n \cdot x = n_1 x_1 + n_2 x_2 + \dots + n_d x_d$. Prove that if

$$\sum_{n \in \mathbb{Z}^d} |n|^{2k} |a_n|^2 < \infty$$

for some $k > \frac{d}{2}$, then f is continuous.

Hunter and Nachtergaele 7.6

Suppose that $f \in H^1([a, b])$ and $f(a) = f(b) = 0$. Prove the Poincaré inequality

$$\int_a^b |f(x)|^2 dx \leq \frac{(b-a)^2}{\pi^2} \int_a^b |f'(x)|^2 dx.$$

Hunter and Nachtergaele 7.7

Solve the following initial-boundary value problem for the heat equation,

$$\begin{aligned} u_t &= u_{xx}, \\ u(0, t) &= 0, \quad u(L, t) = 0 \quad \text{for } t > 0 \\ u(x, 0) &= f(x) \quad \text{for } 0 \leq x \leq L \end{aligned}$$

Suppose $u(x, t) = F(x)G(t)$ is a solution. Then

$$\begin{aligned} u_t &= u_{xx} \\ \implies F(x)G'(t) &= F''(x)G(t) \\ \implies \frac{F''(x)}{F(x)} &= \frac{G'(t)}{G(t)} \end{aligned}$$

Since the left hand side is a function of x and the right hand side is a function of t , they can only be equal if they are both constant, i.e.

$$\frac{F''(x)}{F(x)} = C = \frac{G'(t)}{G(t)}$$

for some $C \in \mathbb{R}$. Thus,

$$F''(x) - CF(x) = 0, \quad \text{and} \tag{1}$$

$$G'(t) - CG(t) = 0 \tag{2}$$

Let $\lambda = \sqrt{|C|}$. The solutions of (1) are

$$F(x) = c_1 e^{\lambda i t} + c_2 e^{-\lambda i t}$$