

---

# Homework #2

---

Sam Fleischer

April 12, 2016

<b>Problem 1</b>	<b>2</b>
<b>Problem 2</b>	<b>2</b>
<b>Problem 3</b>	<b>2</b>
<b>Problem 4</b>	<b>2</b>
<b>Problem 5</b>	<b>2</b>
<b>Problem 6</b>	<b>3</b>
<b>Problem 7</b>	<b>3</b>
<b>Problem 8</b>	<b>3</b>
<b>Problem 9</b>	<b>3</b>

**Problem 1**

A function  $f \in L^p(\mathbb{R}^n)$  is said to be  $L^p$ -continuous if  $\tau_h f \rightarrow f$  in  $L^p(\mathbb{R}^n)$  as  $h \rightarrow 0$  in  $\mathbb{R}^n$ , where  $\tau_h f(x) = f(x-h)$  is the translation of  $f$  by  $h$ . Prove that, if  $1 \leq p < \infty$ , every  $f \in L^p(\mathbb{R}^n)$  is  $L^p$ -continuous. Give a counter-example to show that this result is not true when  $p = \infty$ . [Hint: Approximate an  $L^p$  function by a  $C_c$  function.]

*Proof.* Define  $f \in L^\infty(\mathbb{R})$  as  $f(x) = \mathcal{X}_{[0,1]^n}$  where  $\mathcal{X}$  is the characteristic function. Note that  $f(1-\varepsilon) = 1$  for all  $\varepsilon > 0$ . Let  $h$  be a small perturbation, i.e.  $0 < |h| \ll 1$ , and choose  $\varepsilon = \frac{h}{2}$ . Then  $\forall x \in (0, \varepsilon)$ ,  $\tau_h f(x) = 0$  but  $f(x) = 1$ , and thus  $|\tau_h f(x) - f(x)| = 1$ . This shows that  $\forall h > 0$ ,  $\exists$  an interval  $I_h$  (of positive measure,  $\mu(I_h) > 0$ ) such that  $|\tau_h f(x) - f(x)| = 1$  for all  $x \in I_h$ . Thus  $\tau_h f \not\rightarrow f$  in  $L^\infty(\mathbb{R}^n)$ .  $\square$

**Problem 2**

Show that  $L^\infty(\mathbb{R})$  is not separable. [Hint: There is an uncountable set  $\mathcal{F} \subset L^\infty$  such that  $\|f - g\|_\infty \geq 1$  for all  $f, g \in \mathcal{F}$  with  $f \neq g$ .]

*Proof.*

$\square$

**Problem 3**

Prove Chebyshev's Inequality: If  $f \in L^p$  ( $1 \leq p < \infty$ ), then for any  $\alpha > 0$ ,

$$\mu(\{x : |f(x)| > \alpha\}) \leq \left( \frac{\|f\|_p}{\alpha} \right)^p.$$

[Note that you can find the proof of this simple fact in many texts but you should see if you can figure it out yourself. Also, note that this inequality holds for all  $0 < p < \infty$ .]

*Proof.*

$\square$

**Problem 4**

Assume that  $f, g \in L^1(\mathbb{R}^n)$ . Prove that the convolution

$$(f * g)(x) = \int_{\mathbb{R}^n} f(x-y)g(y)dy$$

is measurable and in  $L^1(\mathbb{R}^n)$ .

*Proof.*

$\square$

**Problem 5**

Let  $f_n = \sqrt{n}\mathcal{X}_{(0, \frac{1}{n})}$ . Prove that  $f_n$  converges weakly to 0 in  $L^2(0, 1)$  and  $f_n \rightarrow 0$  in  $L^1(0, 1)$  but  $f_n$  does not converge strongly in  $L^2(0, 1)$ .

*Proof.*

$$\|f_n\|_2^2 = \int_0^1 n \mathcal{X}_{(0, \frac{1}{n})} dx = \int_0^{\frac{1}{n}} n dx = 1$$

Thus  $\|f_n\|_2 = 1$  for all  $n$ , and thus does not converge strongly to 0 in  $L^2(0, 1)$ .

$$\|f_n\|_1 = \int_0^1 \sqrt{n} \mathcal{X}_{(0, \frac{1}{n})} dx = \int_0^{\frac{1}{n}} \sqrt{n} dx = \frac{1}{\sqrt{n}}$$

Thus  $\|f_n\|_1 \rightarrow 0$  as  $n \rightarrow \infty$ , which shows  $f_n \rightarrow 0$  strongly in  $L^1(0, 1)$ . Let  $L \in L^2(0, 1)^* \cong L^2(0, 1)$ . Thus  $\exists \ell \in L^2(0, 1)$  such that

$$L(f) = \int_0^1 \ell(x) f(x) dx$$

for all  $f \in L^2$ . Then

$$L(f_n) = \int_0^1 \ell(x) \sqrt{n} \mathcal{X}_{(0, \frac{1}{n})} dx \leq \left( \int_0^1 |\ell(x)|^2 dx \right)^{\frac{1}{2}} \left( \int_0^1 n \mathcal{X}_{(0, \frac{1}{n})} dx \right)^{\frac{1}{2}} = \|\ell\|_2 \int_0^{\frac{1}{n}} n dx = \|\ell\|_2$$

□

### Problem 6

Find a sequence of functions with the property that  $f_j$  converges to 0 in  $L^2(\Omega)$  weakly, to 0 in  $L^{\frac{3}{2}}(\Omega)$  strongly, but it does not converge to 0 strongly in  $L^2(\Omega)$ .

*Proof.*

□

### Problem 7

Let  $f_n$  and  $g_n$  denote two sequences in  $L^p(\Omega)$  with  $1 \leq p \leq \infty$  such that  $f_n \rightarrow f$  in  $L^p(\Omega)$ , and  $g_n \rightarrow g$  in  $L^p(\Omega)$ . Set  $h_n = \max\{f_n, g_n\}$  and prove that  $h_n \rightarrow h$  in  $L^p(\Omega)$ .

*Proof.*

□

### Problem 8

Let  $f_n$  be a sequence in  $L^p(\Omega)$  with  $1 \leq p < \infty$ , and let  $g_n$  be a bounded sequence in  $L^\infty(\Omega)$ . Suppose that  $f_n \rightarrow f$  in  $L^p(\Omega)$  and that  $g_n \rightarrow g$  pointwise a.e. Prove that  $f_n g_n \rightarrow f g$  in  $L^p(\Omega)$ .

*Proof.*

□

### Problem 9

Prove that the space of continuous functions with compact support  $\mathcal{C}_c^0(\mathbb{R}^n)$  is dense in  $L^p(\mathbb{R}^n)$  for  $1 \leq p < \infty$ .

*Proof.*

□