

---

# Homework #7

---

Sam Fleischer

March 18, 2016

<b>Problem 1</b>	.....	<b>2</b>
<b>Problem 2</b>	.....	<b>3</b>

---

## Problem 1

---

- (a) Use separation of variables to find the eigenvalues  $\lambda$  and eigenfunctions  $u(x, y)$  of the Dirichlet Laplacian on the unit square that satisfy

$$\begin{aligned} -(u_{xx} + u_{yy}) &= \lambda u & 0 < x < 1, 0 < y < 1 \\ u(x, 0) &= 0, & u(x, 1) &= 0 & 0 \leq x \leq 1 \\ u(0, y) &= 0, & u(1, y) &= 0 & 0 \leq y \leq 1. \end{aligned}$$

Suppose  $u(x, y) = F(x)G(y)$ . Then

$$\begin{aligned} -F''G - FG'' &= \lambda FG & \implies & -F''G = F(G'' + \lambda G) \\ & & \implies & -\frac{F''}{F} = \frac{G''}{G} + \lambda \end{aligned}$$

Since the left hand side is a function of  $x$  and the right hand side is a function of  $y$ , then they can only be equal if

$$-\frac{F''}{F} = \frac{G''}{G} + \lambda = \mu$$

where  $\mu$  is a constant. Note the boundary conditions imply

$$F(0) = F(1) = G(0) = G(1) = 0$$

thus  $F'' + \mu F = 0$  and the homogeneous Dirichlet boundary conditions imply  $\mu > 0$  and

$$F(x) = A \sin(\sqrt{\mu}x)$$

The condition  $F(1) = 0$  implies

$$0 = A \sin(\sqrt{\mu}) \implies \mu = \pi^2 n^2$$

for  $n \geq 1$ . Then  $G'' + (\lambda - \mu)G = 0$  and the homogeneous Dirichlet boundary conditions imply  $\lambda - \mu > 0$  and

$$G(y) = B \sin(\sqrt{\lambda - \mu}y)$$

The condition  $G(1) = 0$  implies

$$0 = B \sin(\sqrt{\lambda - \mu}) \implies \lambda - \mu = \pi^2 m^2 \implies \lambda = \pi^2(n^2 + m^2)$$

for  $n, m \geq 1$ . Thus the solution to  $-(u_{xx} + u_{yy}) = \lambda u$  is

$$\sum_{n, m \geq 1} A_{n, m} \sin(n\pi x) \sin(m\pi y)$$

for some constants  $A_{n, m}$ . Then the eigenvalues of

$$-\nabla^2 u = \lambda u; \quad u(x, 0) = u(x, 1) = u(0, y) = u(1, y) = 0$$

are  $\lambda = \pi^2(m^2 + n^2)$  for all  $n, m \geq 1$ .

(b) What is the smallest eigenvalue that is not a simple eigenvalue?

The smallest eigenvalue that is not a simple eigenvalue is  $5\pi^2$  since this can be achieved when  $n = 1$ ,  $m = 2$  or when  $n = 2$ ,  $m = 1$ . The only smaller eigenvalue is  $2\pi^2$ , but that is simple since it can only be achieved when  $n = m = 1$ .

---

## Problem 2

---

(a) Let  $\vec{x} = (x, y)$ ,  $\vec{\xi} = (\xi, \eta)$ , and  $\vec{\xi}^* = (\xi, -\eta)$  where  $\eta > 0$ . Show that

$$G(\vec{x}, \vec{\xi}) = -\frac{1}{2\pi} \log \left( \frac{|\vec{x} - \vec{\xi}|}{|\vec{x} - \vec{\xi}^*|} \right)$$

is the solution of

$$\begin{aligned} -(G_{xx} + G_{yy}) &= \delta(\vec{x} - \vec{\xi}) & \text{in } -\infty < x < \infty, \quad y > 0 \\ G(\vec{x}, \vec{\xi}) &= 0 & \text{on } y = 0. \end{aligned}$$

$$\begin{aligned} -\nabla^2 G &= -\nabla^2 \left[ -\frac{1}{2\pi} \log \left( \frac{|\vec{x} - \vec{\xi}|}{|\vec{x} - \vec{\xi}^*|} \right) \right] \\ &= -\nabla^2 \left[ -\frac{1}{2\pi} \log |\vec{x} - \vec{\xi}| - \left( -\frac{1}{2\pi} \log |\vec{x} - \vec{\xi}^*| \right) \right] \\ &= -\nabla^2 \left[ G_F(|\vec{x} - \vec{\xi}|) - G_F(|\vec{x} - \vec{\xi}^*|) \right] \end{aligned}$$

where

$$G_F(r) = -\frac{1}{2\pi} \log r$$

is the free-space Green's function in two dimensions. By definition,

$$-\nabla^2 G_F(|\vec{x} - \vec{\xi}|) = \delta(\vec{x} - \vec{\xi}) \quad \text{and} \quad -\nabla^2 G_F(|\vec{x} - \vec{\xi}^*|) = \delta(\vec{x} - \vec{\xi}^*).$$

But  $\delta(\vec{x} - \vec{\xi}^*) = 0$  for  $y > 0$  since  $\xi^* = (\xi, -\eta)$  where  $\eta > 0$ . Thus,

$$-\nabla^2 G = -\nabla^2 G_F(|\vec{x} - \vec{\xi}|) + \cancel{\nabla^2 G_F(|\vec{x} - \vec{\xi}^*|)}^0 = \delta(\vec{x} - \vec{\xi}) \quad \text{for } (x, y) \in \{(x, y) : y > 0\}$$

When  $y = 0$ ,  $|\vec{x} - \vec{\xi}| = |\vec{x} - \vec{\xi}^*|$ , thus

$$\frac{|\vec{x} - \vec{\xi}|}{|\vec{x} - \vec{\xi}^*|} = 1 \quad \implies \quad G(\vec{x}, \vec{\xi}) = -\frac{1}{2\pi} \log(1) = 0$$

- (b) Write down the Green's function representation for the solution  $u(x, y)$  of the Dirichlet problem for the Laplacian in the upper half plane

$$u_{xx} + u_{yy} = 0 \quad \text{in } -\infty < x < \infty, \quad y > 0$$

$$u(x, 0) = f(x).$$

You can assume that  $u(x, y) \rightarrow 0$  sufficiently rapidly as  $|(x, y)| \rightarrow \infty$ .

We utilize Green's identity

$$\int_{\Omega} (u \nabla^2 v - v \nabla^2 u) d\vec{\xi} = \int_{\partial\Omega} \left( u \frac{\partial v}{\partial n(\vec{\xi})} - v \frac{\partial u}{\partial n(\vec{\xi})} \right) ds(\vec{\xi})$$

by setting  $\Omega = \{(x, y) : y > 0\}$  and  $v = G$  where  $G$  is the free-space Green's function which solves  $-\nabla^2 G = \delta(x)$ . Thus,

$$\int_{\Omega} (u \nabla^2 G - G \nabla^2 u) d\vec{\xi} = \int_{\partial\Omega} \left( u \frac{\partial G}{\partial n(\vec{\xi})} - G \frac{\partial u}{\partial n(\vec{\xi})} \right) ds(\vec{\xi})$$

However,  $u$  is harmonic in  $\Omega$ , i.e.  $\nabla^2 u \equiv 0$  for  $(x, y) \in \Omega$ , and  $-\nabla^2 G = \delta(\vec{x} - \vec{\xi})$ , i.e.

$$\int_{\Omega} (u \nabla^2 G - G \nabla^2 u) d\vec{\xi} = - \int_{\Omega} u(\vec{\xi}) \delta(\vec{x} - \vec{\xi}) d\vec{\xi} = -u(\vec{x})$$

Also, the Dirichlet condition  $u(\vec{x}) = f(x)$  on  $\partial\Omega$  give us  $G \equiv 0$  for  $y = 0$  (i.e. for  $(x, y) \in \partial\Omega$ ). Thus,

$$-u(\vec{x}) = \int_{\partial\Omega} \left( u \frac{\partial G}{\partial n(\vec{\xi})} - G \frac{\partial u}{\partial n(\vec{\xi})} \right) ds(\vec{\xi}) = \int_{\partial\Omega} u(\vec{\xi}) \frac{\partial G(x, y; \xi, \eta)}{\partial n(\vec{\xi})} ds(\vec{\xi})$$

Finally, the unit normal vector on the boundary of  $\Omega$  is  $-\eta$ , and thus

$$u(\vec{x}) = \int_{\mathbb{R}} u(\xi) \frac{\partial G(x, y; \xi, \eta)}{\partial \eta} d\xi$$

- (c) Use the Green's function representation to show that

$$u(x, y) = \frac{y}{\pi} \int_{-\infty}^{\infty} \frac{f(t)}{(x-t)^2 + y^2} dt.$$

Rewrite  $G(\vec{x}; \vec{\xi}) = G(x, y; \xi, \eta)$ . Then

$$G(\vec{x}, \vec{\xi}) = \frac{1}{2\pi} \log \left( \frac{|\vec{x} - \vec{\xi}|}{|\vec{x} - \vec{\xi}^*|} \right) = -\frac{1}{4\pi} \log \left( \frac{(x-\xi)^2 + (y-\eta)^2}{(x-\xi)^2 + (y+\eta)^2} \right)$$

$$\Rightarrow \frac{\partial G}{\partial \eta} \Big|_{\eta=0} = \left[ -\frac{1}{4\pi} \cdot \frac{[(x-\xi)^2 + (y+\eta)^2](-2)(y-\eta) - [(x-\xi)^2 + (y-\eta)^2](2)(y+\eta)}{((x-\xi)^2 + (y+\eta)^2)^2} \right]_{\eta=0}$$

$$= -\frac{1}{4\pi} \cdot \frac{[(x-\xi)^2 + y^2](-2y) - [(x-\xi)^2 + y^2](2y)}{((x-\xi)^2 + y^2)^2}$$

$$\begin{aligned}
 &= -\frac{1}{4\pi} \cdot \frac{1}{(x-\xi)^2 + y^2} \cdot (-4y) \\
 &= \frac{y}{\pi} \frac{1}{(x-\xi)^2 + y^2}
 \end{aligned}$$

Thus,

$$u(x, y) = u(\vec{x}) = \int_{\mathbb{R}} u(\vec{\xi}) \frac{\partial G(x, y; \xi, \eta)}{\partial \eta} d\xi = \frac{y}{\pi} \int_{\mathbb{R}} \frac{u(\vec{\xi})}{(x-\xi)^2 + y^2} d\xi$$

Define  $t := \xi$  and note  $u(\vec{\xi}) = f(\xi)$  when  $\eta = 0$ . Thus,

$$u(x, y) = \frac{y}{\pi} \int_{\mathbb{R}} \frac{f(t)}{(x-t)^2 + y^2} dt$$