
Homework #1

Sam Fleischer

April 8, 2016

Problem 1	2
Problem 2	2

Problem 1

Find the two-term asymptotic expansion for small ε for all real roots x of the below equations.

(a) $\varepsilon^2 x^3 - x + \varepsilon = 0$

(b) $\varepsilon \exp(x^2) = 1 + \frac{\varepsilon}{1+x^2}$

(a) First note that if $\varepsilon = 0$, then $x = 0$. Let $x = x_0 + x_1\varepsilon + x_2\varepsilon^2 + \dots$. Then

$$\begin{aligned} & \varepsilon^2 (x_0 + x_1\varepsilon + x_2\varepsilon^2 + \dots)^3 - (x_0 + x_1\varepsilon + x_2\varepsilon^2 + \dots) + \varepsilon = 0 \\ \Rightarrow & \varepsilon^2 (x_0^3 + 3x_0^2x_1\varepsilon + (3x_0^2x_2 + 3x_0x_1^2)\varepsilon^2 + \dots) - (x_0 + x_1\varepsilon + x_2\varepsilon^2 + \dots) + \varepsilon = 0 \\ \Rightarrow & x_0 = 0 \quad \text{and} \quad x_1 = 1 \quad \text{and} \quad x_2 = 0 \\ \Rightarrow & x = \varepsilon + O(\varepsilon^3) \end{aligned}$$

Next we rescale to find the other two roots. Let $x = \varepsilon^{-1}y$. Then

$$\begin{aligned} & \varepsilon^{-1}y^3 - \varepsilon^{-1}y + \varepsilon = 0 \\ \Rightarrow & y^3 - y + \varepsilon^2 = 0 \end{aligned}$$

Let $y = y_0 + y_1\varepsilon + y_2\varepsilon^2 + \dots$. Then

$$\begin{aligned} & (y_0 + y_1\varepsilon + y_2\varepsilon^2 + \dots)^3 - (y_0 + y_1\varepsilon + y_2\varepsilon^2 + \dots) + \varepsilon^2 = 0 \\ \Rightarrow & (y_0^3 + 3y_0^2y_1\varepsilon + (3y_0^2y_2 + 3y_1^2)\varepsilon^2 + \dots) - (y_0 + y_1\varepsilon + y_2\varepsilon^2 + \dots) + \varepsilon^2 = 0 \\ \Rightarrow & y_0^3 - y_0 = 0 \iff y_0 = 0, \pm 1 \quad \text{and} \quad (3y_0^2 - 1)y_1 = 0 \Rightarrow y_0^2 \neq \frac{1}{3} \text{ or } y_1 = 0 \\ \Rightarrow & 3y_0^2y_2 - y_2 + 1 = 0 \Rightarrow y_2 = \frac{1}{1 - 3y_0^2} \end{aligned}$$

If $y_0 = 0$, then $y_2 = 1$. If $y_0 = \pm 1$, then $y_2 = -\frac{1}{2}$. Thus the asymptotic solutions are

- $y = \varepsilon^2 + O(\varepsilon^3)$ (this matches with the asymptotic solution reached above).
- $y = \pm 1 - \frac{1}{2}\varepsilon^2 + O(\varepsilon^3)$. These are impossible to get in the original scaling since as $y \rightarrow \pm 1$, $x \rightarrow \pm\infty$.

Problem 2

The Exponential integral function is defined as

$$\text{Ei}(x) = \int_x^\infty \frac{\exp(-s)}{s} ds.$$

Derive an asymptotic expansion for $\text{Ei}(x)$ for large x . Use a computer (e.g. `expint(x)` in MATLAB) to check the accuracy of your expansion for different values of x and for different numbers of terms. Discuss your results.

By integration by parts,

$$\int_x^\infty \frac{\exp(-s)}{s} ds = \frac{-\exp(-s)}{s} \Big|_x^\infty - \int_x^\infty \frac{\exp(-s)}{s^2} ds$$

$$\begin{aligned}
&= \frac{\exp(-x)}{x} - \left[\frac{-\exp(-s)}{s^2} \right]_x^\infty - 2 \int_x^\infty \frac{\exp(-s)}{s^3} ds \\
&\vdots \\
&= \exp(-x) \left[\frac{(-1)^0 0!}{x} + \frac{(-1)^1 1!}{x^2} + \frac{(-1)^2 2!}{x^3} + \cdots + \frac{(-1)^n n!}{x^{n+1}} \right] + R_n(x)
\end{aligned}$$

where

$$R_n(x) = (-1)^{n+1} (n+1)! \int_x^\infty \frac{\exp(-s)}{s^{n+2}} ds$$

Thus the asymptotic expansion for $\text{Ei}(x)$ for large x is

$$\text{Ei}(x) \sim \exp(-x) \sum_{n=0}^{\infty} \frac{(-1)^n n!}{x^{n+1}}.$$

This is a divergent series in n , but converges to 0 as $x \rightarrow \infty$ when n is fixed. The following graph shows the error between the asymptotic expansion and the exponential integral function, as calculated by MATLAB and graphed in Python.

