

MAT 228A Notes

Sam Fleischer

November 29, 2016

1 Conjugate Gradient Method

$Au = f$ where A is symmetric and positive definite (s.p.d.). It turns out u minimizes ϕ where

$$\phi(u) = u^T Au - u^T f.$$

Remember the definition

$$\kappa = \|A\|_2 \|A^{-1}\|_2.$$

For symmetric matrices,

$$\kappa = \frac{\max_j |\lambda_j|}{\min_j |\lambda_j|}.$$

In conjugate gradient method, we will not always go in the direction of steepest descent (the residual direction). Let p_k be the vector to follow.

$$u_{k+1} = u_k + \alpha p_k.$$

Follow this search direction until ϕ increases, i.e. minimize ϕ on a 1D space. We find

$$\alpha = \frac{p_k^T r_k}{p_k^T A p_k}.$$

1.1 Start in 2D

- Initial guess u_0
- compute residual r_0
- Initialize $p_0 = r_0$
- Get $u_1 = u_0 + \alpha p_0$
- Pick p_1 so that $p_1^T A p_0 = 0$.
 - In otherwords, p_0 and p_1 are A -conjugate, or

$$p_1^T A p_0 = \langle p_0, p_1 \rangle_A$$

because A is s.p.d.

- Why do this? Notice that p_0 is tangent to the level set of ϕ at $\phi(u_1)$, i.e. we know $p_0^T r_1 = 0$. So,

$$\begin{aligned} p_0^T r_1 &= 0 \\ p^T (f - Au_1) &= 0 \\ p^T (Au - Au_1) &= 0 \\ p^T A(u - u_1) &= 0 \end{aligned}$$

So p_1 will be pointing in the direction of $u - u_1$. So u_2 is the exact solution.

1.2 3D

- initial guess u_0
- compute residual r_0
- Initialize $p_0 = r_0$
- Get $u_1 = u_0 + \alpha p_0$
- Pick p_1 to be A -conjugate to p_0 .
 - This is a 2D space to choose from.. just pick a direction in that space.
- p_1 and r_1 define α
- calculate $u_2 = u_1 + \alpha p_1$.
 - p_0 and p_1 span a plane. This plane, which is $c_0 p_0 + c_1 p_1 + u_2$, is tangent to the level surface at $\phi(u_2)$.
 - At this point, we've minimized on this plane.
- Pick p_2 to be A -conjugate to $\{p_0, p_1\}$.
- Minimizing ϕ along this direction gives us the exact solution.

$$\begin{aligned}
 (c_0 p_0 + c_1 p_1)^T r_2 &= 0 \\
 (c_0 p_0 + c_1 p_1)^T (f - Au_2) &= 0 \\
 (c_0 p_0 + c_1 p_1)^T (Au - Au_2) &= 0 \\
 (c_0 p_0 + c_1 p_1)^T A(u - u_2) &= 0
 \end{aligned}$$

So p_2 is pointing right at the solution.

1.3 In general

So CG will (no round off) give the exact solution in N steps. Usually we get close in far fewer steps.

1.3.1 Pseudocode

- Initialize $u_0, r_0 = f - Au_0, p_0 = r_0$
- loop in k
 - $w = Ap_k$
 - $\alpha = \frac{r_k^T r_k}{w^T p_k}$
 - $u_{k+1} = u_k + \alpha p_k$
 - $r_{k+1} = r_k - \alpha w$
 - check $\|r_{k+1}\|$ for stopping
 - compute $\beta = \frac{r_{k+1}^T r_{k+1}}{r_k^T r_k}$
 - $p_{k+1} = r_{k+1} + \beta p_k$
 - * This is r_{k+1} without components in $\{p_0, p_1, \dots, p_{k-1}\}$.

2 Analysis of Convergence

Because A is s.p.d. it defines a norm $\|u\|_A^2 = u^T A u$. We can show $\|e_j\|_A \leq 2 \left(\frac{\sqrt{\kappa} - 1}{\sqrt{\kappa} + 1} \right)^j \|e_0\|_A$, which shows convergence since $\|e_j\|_A \rightarrow 0$ as $j \rightarrow \infty$. So, $\kappa \approx 1$ gives fast convergence and $\kappa \gg 1$ gives slow convergence.

2.1 Theorem from Book

The vectors in the CG algorithm have the following properties provided $r_k \neq 0$:

1. p_k is A -conjugate to all previous search directions.
2. r_k is orthogonal to all previous residuals.
3. The following subspaces of \mathbb{R}^n are identical

- (a) $\text{span}\{p_0, p_1, \dots, p_{k-1}\}$

- (b) $\text{span}\{r_0, Ar_0, A^2r_0, \dots, A_{k-1}r_0\}$

- (c) $\text{span}\{e_0, Ae_0, A^2e_0, \dots, A_{k-1}e_0\}$

Define $K_n = \text{span}\{r_0, Ar_0, A^2r_0, \dots, A_{n-1}r_0\}$ as the n -dimensional Krylov space associated with r_0 .

$$u_n \in u_0 + K_n.$$

u_n minimizes ϕ in this space. Minimizing $\phi(u)$ is equivalent to minimizing $\|e_j\|_A^2$.

$$\begin{aligned} \|e_j\|_A^2 &= e_j^T A e_j \\ &= (u_j - u)^T A (u_j - u) \\ &= u_j^T A u_j - 2u_j^T A u + u^T A u \\ &= u_j^T A u - 2u_j^T f + u^T A u \\ &= 2\phi(u_j) + C \end{aligned}$$