Section 5.2 - Binomial Probability Distributions

Chris Godbout

Outline

Introduction

Examples

Introduction

Definition (Binomial Probability Distribution)

Definition (Binomial Probability Distribution)

A binomial probability distribution results from a procedure that meets the following requirements:

1. The procedure has a fixed number of trials.

Definition (Binomial Probability Distribution)

- 1. The procedure has a fixed number of trials.
- 2. The trials must be independent.

Definition (Binomial Probability Distribution)

- 1. The procedure has a fixed number of trials.
- 2. The trials must be independent.
- 3. Each trial must have all outcomes classified into *two categories* (usually called success and failure).

Definition (Binomial Probability Distribution)

- 1. The procedure has a fixed number of trials.
- 2. The trials must be independent.
- 3. Each trial must have all outcomes classified into *two categories* (usually called success and failure).
- 4. The probability must remain the same for all trials.

Definition (Binomial Probability Distribution)

- 1. The procedure has a fixed number of trials.
- 2. The trials must be independent.
- 3. Each trial must have all outcomes classified into *two categories* (usually called success and failure).
- 4. The probability must remain the same for all trials.
- 5. The random variable *X* counts the number of successes in *n* trials.

 \cdot The probability of success is p.

- The probability of success is *p*.
- The probability of failure is 1 p.

- The probability of success is *p*.
- The probability of failure is 1 p.
- *n* is the fixed number of trials.

- The probability of success is *p*.
- The probability of failure is 1 p.
- *n* is the fixed number of trials.
- P(X = x) is the probability of getting exactly x successes.

- The probability of success is *p*.
- The probability of failure is 1 p.
- *n* is the fixed number of trials.
- P(X = x) is the probability of getting exactly x successes.
- P(X < x) is the probability of getting fewer than x successes.

Formula

Theorem (Binomial Probability Formula)

$$P(X = x) = {}_{n}C_{x} \cdot p^{x} \cdot (1-p)^{n-x}$$

Mean and Standard Deviation

Mean and Standard Deviation

Mean

The mean of a binomial probability distribution is

$$E(X) = \mu = np$$

Mean and Standard Deviation

Mean

The mean of a binomial probability distribution is

$$E(X) = \mu = np$$

Standard Deviation

The standard deviation of a binomial probability distribution is

$$\sigma = \sqrt{np(1-p)}$$

Examples

You are in the middle of a nightmare and have to take an exam for which you haven't studied. Fortunately, it's a multiple choice test with 10 question. Each question as 4 choices.

What is *p*?

- What is *p*?
- What is P(X = 7)?

- What is *p*?
- What is P(X = 7)?
- What is E(X)?

- What is *p*?
- What is P(X = 7)?
- What is E(X)?
- What is the probability that the student gets at least 7 correct?

Assume that 45% of all people have blood that is in Group O.

Assume that 45% of all people have blood that is in Group O.

• What is the probability that exactly 3 out of 8 people have blood in Group O?

7

Assume that 45% of all people have blood that is in Group O.

- What is the probability that exactly 3 out of 8 people have blood in Group O?
- What is the probability that no more than 3 out of 8 people have blood in Group O?

7

Assume that 45% of all people have blood that is in Group O.

- · What is the probability that exactly 3 out of 8 people have blood in Group O?
- What is the probability that no more than 3 out of 8 people have blood in Group O?
- What is the probability that exactly 16 out of 21 people have blood in Group
 O?