

Section 10.3 — Hypothesis Testing for Population Means (σ unknown)

Chris Godbout

Performing a Hypothesis Test

Hypothesis test for Population Means (σ unknown)

Performing a Hypothesis Test

Basic Steps

1. State the null and alternative hypotheses.

Basic Steps

1. State the null and alternative hypotheses.
2. Determine which distribution to use for the test statistic and state the level of significance.

Basic Steps

1. State the null and alternative hypotheses.
2. Determine which distribution to use for the test statistic and state the level of significance.
3. Gather data and calculate the necessary sample statistics.

Basic Steps

1. State the null and alternative hypotheses.
2. Determine which distribution to use for the test statistic and state the level of significance.
3. Gather data and calculate the necessary sample statistics.
4. Draw a conclusion and interpret the decision.

Types of Hypothesis Tests

Symbol in H_a	Tail
\neq	Two-tailed test
$<$	Left-tailed test
$>$	Right-tailed test

Definition (Critical region)

The **critical region** or **rejection region** is the set of values of the test statistic that cause us to reject the null hypothesis.

Critical values

Definition (Critical region)

The **critical region** or **rejection region** is the set of values of the test statistic that cause us to reject the null hypothesis.

Definition (Critical values)

The **critical values** are the values that separate the critical region from the values of the test statistic that do not lead to rejection of the null hypothesis. These are determined by α .

Definition (p -values)

A p -value is the probability of obtaining a sample statistic as extreme or more extreme than the one observed in the data when the null hypothesis is assumed to be true.

p -values

Definition (p -values)

A p -value is the probability of obtaining a sample statistic as extreme or more extreme than the one observed in the data when the null hypothesis is assumed to be true.

Determining p -values

Left-tailed	p -value = area to the <i>left</i> of the test statistic
Right-tailed	p -value = area to the <i>right</i> of the test statistic
Two-tailed	p -value = <i>twice</i> the area in the tail beyond the test sta

Make a decision

Critical value method

If the test statistic is in the critical region, reject H_0 .
Otherwise, fail to reject.

p -value method

If $p\text{-value} \leq \alpha$, reject H_0 . Otherwise, fail to reject H_0 .

Hypothesis test for Population Means (σ unknown)

$$t = \frac{\bar{x} - \mu}{s/\sqrt{n}}$$

with

$$df = n - 1$$

Teachers' salaries

Teacher's salaries in one district are very low, so low that educators in that state regularly complain about their compensation. The state mean is \$33,600, but teachers in one district claim that the mean in their district significantly lower. They survey a simple random sample of 22 teachers in the district and calculate the mean salary of \$32,400 with a standard deviation of \$1520. Test the teacher's claim at the 0.01 level of significance.

Children's Clothing

A children's clothing company sells hand-smocked dresses for girls. The length of one particular size dress is designed to be 26 inches. The company regularly tests the lengths of the garments to ensure quality control. If the mean length is found to be significantly longer or shorter than 26 inches, then the machines must be adjusted. The most recent simple random sample of 28 dresses had a mean length of 26.30 inches with a standard deviation of 0.77 inches. Perform a hypothesis test on the accuracy of the machines at the 0.01 level of significance.