

# VFTS682: a confirmed dynamical ejection?

M. Renzo<sup>1</sup> and ■ [TBD] ■.

Astronomical Institute Anton Pannekoek, University of Amsterdam, 1098 XH Amsterdam, The Netherlands

## ABSTRACT

**Key words.** stars: kinematics, stars: runaways, stars: individual: VFTS682

## 1. Introduction

How do stars form is one longstanding question in astrophysics (Lada & Lada (2003); Zinnecker & Yorke (2007)). It is particularly difficult for massive stars, because these are intrinsically rare (e.g., Salpeter 1955; Kroupa 2001; Schneider et al. 2018), evolve fast, and remain enshrouded in their parent cloud during the formation process. Moreover, observations of young massive stars reveal a complicated multiplicity structure which requires explanation (Kobulnicky & Fryer 2007; Mason et al. 2009; Sana & Evans 2011; Sana et al. 2012; Kiminki & Kobulnicky 2012; Chini et al. 2012; Kobulnicky et al. 2014; Almeida et al. 2017; De Marco & Izzard 2017). Understanding massive star formation, possibly as a function of metallicity, is a key question given the present and upcoming transient survey (e.g., LSST, Black-Gem, LIGO/Virgo O3 ) which will reveal transients associated to massive stars evolution and death.

The second data release (DR2) from the Gaia satellite (Gaia Collaboration et al. 2016, 2018) allows us to test these hypothesis using one particular star, VFTS682. This star is a very massive ( $M_{\text{ZAMS}} \simeq 150 M_{\odot}$ , Bestenlehner et al. 2011; Schneider et al. 2018) WNh5 star in the 30 Doradus region of the Large Magellanic Cloud (LMC), and it is presently observed at a projected distance of  $\sim 29$  pc from the nearest cluster of massive stars R136 (Bestenlehner et al. 2011). Based on the extremely high mass of this star and its present day apparent isolation, Bestenlehner et al. (2011) proposed it might be a candidate for isolated star formation, or a “slow runaway” ejected from R136 in the past. This second option is also supported by the N-body simulations of Fujii & Portegies Zwart (2011); Banerjee et al. (2012). Many other very massive stars are present in the surroundings of R136, and a more detailed analysis on the larger sample is desirable.

Massive stars can in principle be ejected from R136 as a consequence of dynamical interactions (Poveda et al. 1967; Leonard 1991; Evans et al. 2010; Fujii & Portegies Zwart 2011; Allison 2012; Oh & Kroupa 2016), or by the disruption of a binary by the first core-collapse supernova (Zwicky 1957; Blaauw 1961; De Donder et al. 1997; Eldridge et al. 2011; Renzo et al. 2018). However, R136 has an estimated age of  $\lesssim 2$  Myr (Sabbi et al. 2012), which is shorter than the shortest stellar lifetime ( $\sim 3$  Myr, e.g., Zapartas et al. 2017), so one would not expect the binary disruption scenario to be relevant for this cluster.

In this study, we combine the radial velocity measurements from the VFTS survey (Evans et al. 2011) with the proper motion from Gaia DR2 to reconstruct the three-dimensional velocity of VFTS682, and test the null hypothesis that this star was ejected from R136. ■ [check the following] ■ Our results indicate that R136 is the likely origin of this star, and therefore isolated star formation is *not* required to explain it. However, we find a mild discrepancy between the apparent age of VFTS682, its kinematic age, and the age estimates for R136.

In Sec. 2, we describe the data selection and validation process for VFTS 682 and for stars in the surrounding used to define the local reference frame. Sec. 3 presents our main findings. We conclude by discussing the implications for theories of star formation, N-body interactions, and binary evolution in Sec. 4.

## 2. Gaia DR2 data selection

VFTS682 is labeled in the Gaia DR2 catalog<sup>1</sup> with the source id 4657685637907503744. The star has a `visibility_period` = 17, which counts how many observations have been used to reconstruct its astrometric solution (Lindgren et al. 2018). Its reported G-band magnitude is 15.65, cf. the V-band magnitude of 16.08 (Evans et al. 2011; Bestenlehner et al. 2011), and the reported `astrometric_excess_noise` = 0. These values suggest that the Gaia data for VFTS682 are trustworthy. However, the effective temperature reported in Gaia DR2 is one order of magnitude lower than what found by Bestenlehner et al. (2011), and the best fit parallax of this star is negative. We do not use the effective temperature of the star anywhere in this study, and we attribute the unphysical value of the parallax to the large distance to the LMC. Our main findings do not rely on the parallax nor the effective temperature values reported in the Gaia DR2 catalog.

We retrieve for VFTS682 the position in right ascension (RA) and declination (DE) in the ICRS frame (Gaia Collaboration et al. 2018), its proper motion components ( $\mu_{\text{RA}}$ , and  $\mu_{\text{DE}}$ , respectively). For the radial velocity of VFTS682 and of the 30 Doradus region as a whole, we instead use the VFTS data as quoted in Bestenlehner et al. (2011). Table 1 lists the values adopted throughout this work for each of these quantities.

To compare the astrometry of VFTS682 and derive its peculiar motion, we then select data from the Gaia DR2 catalog for two regions: the “surroundings” of VFTS 682, and

<sup>1</sup> <https://vizier.u-strasbg.fr/viz-bin/VizieR-3?-source=I/345/gaia2>

**Table 1.** Astrometric parameters for VFTS682. The peculiar radial velocity  $\delta v_{\text{rad}}$  is obtained as the difference between the average radial velocity of the 30 Doradus region ( $270 \pm 10 \text{ km s}^{-1}$ ) minus the radial velocity measured from the HeII  $\lambda 4686$  line for VFTS682 ( $315 \pm 15 \text{ km s}^{-1}$ ).

Parameter	Value	Source
RA [degree]	$84.73 \pm 0.03$	Gaia DR2
DE [degree]	$-69.07 \pm 0.05$	
$\mu_{\text{RA}}$ [mas yr $^{-1}$ ]	$1.84 \pm 0.07$	
$\mu_{\text{DE}}$ [mas yr $^{-1}$ ]	$0.78 \pm 0.08$	
$\delta v_{\text{rad}}$ [km s $^{-1}$ ]	$-45 \pm 25$	Bestenlehner et al. (2011)

the “R136 cluster”. The surrounding region is defined by all the stars in a target of 10 arcminutes around VFTS682 fulfilling the following criteria: we require `visibility_period`  $\geq 5$ , `astrometric_excess_noise`  $\leq 1$ , the error on the proper motion components to be smaller than  $0.1 \text{ mas yr}^{-1}$ , and the proper motion components themselves to be smaller than  $2 \text{ mas yr}^{-1}$  in absolute value. At the distance to the LMC,  $1 \text{ mas yr}^{-1} \approx 250 \text{ km s}^{-1}$  (e.g., ?), so the cut on the values of the proper motions removes stars that would have projected tangential velocities in excess of  $\sim 500 \text{ km s}^{-1}$ , which are most likely to be foreground stars. This selection yields 437 stars.

The “R136 cluster” is effectively defined by taking all the stars within 25 arcseconds from R136a, one of the most massive members of the cluster itself (Crowther et al. 2010), requiring the same “quality” criteria applied above. This selection yields 62 stars. ■ **[improve definition of stars from R136, describe accordingly – check with Danny]** ■

Throughout this study, we assume the same distance of 50 kpc to the star, and to the 30 Doradus region as a whole.

For each of the two local rest frames (“surroundings” and “R136 cluster”), we compute the average transverse velocity as:

$$\langle \mu_i \rangle = \frac{\sum_{\text{stars}} \frac{1}{\Delta \mu_i} \mu_i}{\sum_{\text{stars}} \frac{1}{\Delta \mu_i}}, \quad \Delta \langle \mu_i \rangle = \frac{\sqrt{N}}{\sum_{\text{stars}} \frac{1}{\Delta \mu_i}}, \quad (1)$$

where  $i = \text{RA, DEC}$ , and  $\Delta \mu_i$  is the error on the proper motion component reported by Gaia, and the sums run over all the  $N$  stars of the frame considered. We evaluate each proper motion component separately.

The data retrieved, and the ipython notebook used for the analysis presented here will be made available at ■ **[probably git repo on bitbucket?]** ■.

### 3. The kinematics of VFTS682

#### 3.1. Is it a runaway star?

We first address the question of whether VFTS682 is a typical star from the kinematic point of view, or whether it is a runaway star with a significantly large peculiar velocity compared to its surrounding population. The former is what should be expected if it formed where we observe it today, in relative isolation from other massive stars.

Using the 437 stars selected as the “surrounding population” in Sec. 2, we find averaged proper motion components of  $\mu_{\text{RA}} = 1.695 \pm 0.003 \text{ mas yr}^{-1}$  and  $\mu_{\text{DE}} = 0.691 \pm 0.003 \text{ mas yr}^{-1}$ . Subtracting these values from the proper motions of VFTS682 (see Table 1), we obtain the components of proper motion of the star relative to the surrounding region  $\mu_{\text{RA}}^{\text{sur}} = 0.14 \pm 0.07 \text{ mas yr}^{-1}$  and  $\mu_{\text{DE}}^{\text{sur}} = 0.09 \pm 0.08 \text{ mas yr}^{-1}$ . These can be converted in the

the components of the transverse velocity  $v_{\text{RA}}^{\text{sur}} = 35 \pm 17 \text{ km s}^{-1}$ ,  $v_{\text{DE}}^{\text{sur}} = 22 \pm 19 \text{ km s}^{-1}$ , assuming a distance of 50 kpc (we do not account for the uncertainty in the distance estimate when propagating errors). The radial velocity from Bestenlehner et al. (2011) then gives the third component along the line of sight, allowing us to calculate the speed of the star:

$$v = \sqrt{(v_{\text{RA}}^{\text{sur}})^2 + (v_{\text{DE}}^{\text{sur}})^2 + (\delta v_{\text{rad}})^2} = 61 \pm 1 \text{ km s}^{-1}. \quad (2)$$

This value for the peculiar 3D velocity of VFTS682 make it a “bona fide” runaway star.

■ **[errors are wrong]** ■

#### 3.2. Does it come from the R136 cluster?

The inset plot in Fig. 1 shows a zoom in of the region around R136 which includes VFTS682. To test the hypothesis that the star was indeed ejected from this region, we check the relative orientation of its proper motion. The error cone, obtained by propagating the errors on the proper motion components indicates that R136 is indeed the most likely origin, as expected by Fujii & Portegies Zwart (2011); Banerjee et al. (2012).

■ **[kinematic age]** ■


### 4. Summary and Discussion

- VFTS682 is a bona fide runaway with  $v \sim 60 \text{ km s}^{-1}$  thrown out from R136. Both its speed and the age of the cluster are consistent with a dynamical ejection.
- VFTS682 comes from R136 as was expected by Bestenlehner et al. (2011); Fujii & Portegies Zwart (2011); Banerjee et al. (2012), so it does not require isolated SFH to be explained
- apparent age tension (connect to VFTS16 as well).
- is R136 a single young cluster or a merger
- estimate the influence of the gravitational potential of R136, what is its total mass and relaxation time?

Random notes:  $v \sin(i) < 200 \text{ km s}^{-1}$  from Schneider et al. (2018), age  $1.0 \pm 0.2 \text{ Myr}$  from Schneider et al. (2018)

### References

- Allison, R. J. 2012, MNRAS, 421, 3338  
 Almeida, L. A., Sana, H., Taylor, W., et al. 2017, A&A, 598, A84  
 Banerjee, S., Kroupa, P., & Oh, S. 2012, ApJ, 746, 15  
 Bestenlehner, J. M., Vink, J. S., Gräfenr, G., et al. 2011, A&A, 530, L14  
 Blaauw, A. 1961, Bull. Astron. Inst. Netherlands, 15, 265  
 Chini, R., Hoffmeister, V. H., Nasser, A., Stahl, O., & Zinnecker, H. 2012, MNRAS, 424, 1925  
 Crowther, P. A., Schnurr, O., Hirschi, R., et al. 2010, MNRAS, 408, 731  
 De Donder, E., Vanbeveren, D., & van Bever, J. 1997, A&A, 318, 812  
 De Marco, O. & Izzard, R. G. 2017, PASA, 34, e001  
 Eldridge, J. J., Langer, N., & Tout, C. A. 2011, MNRAS, 414, 3501  
 Evans, C. J., Taylor, W. D., Hénault-Brunet, V., et al. 2011, A&A, 530, A108  
 Evans, C. J., Walborn, N. R., Crowther, P. A., et al. 2010, ApJ, 715, L74  
 Fujii, M. S. & Portegies Zwart, S. 2011, Science, 334, 1380  
 Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, ArXiv e-prints  
 Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., et al. 2016, A&A, 595, A1  
 Kiminki, D. C. & Kobulnicky, H. A. 2012, ApJ, 751, 4  
 Kobulnicky, H. A. & Fryer, C. L. 2007, ApJ, 670, 747  
 Kobulnicky, H. A., Kiminki, D. C., Lundquist, M. J., et al. 2014, ApJS, 213, 34  
 Kroupa, P. 2001, MNRAS, 322, 231  
 Lada, C. J. & Lada, E. A. 2003, ARA&A, 41, 57  
 Leonard, P. J. T. 1991, AJ, 101, 562  
 Lindegren, L., Hernandez, J., Bombrun, A., et al. 2018, ArXiv e-prints



**Fig. 1.** position and projected relative velocity to R136. ■ [orient properly, load picture on background, add cone of uncertainty] ■

- Mason, B. D., Hartkopf, W. I., Gies, D. R., Henry, T. J., & Helsel, J. W. 2009, AJ, 137, 3358
- Oh, S. & Kroupa, P. 2016, A&A, 590, A107
- Poveda, A., Ruiz, J., & Allen, C. 1967, Boletín de los Observatorios Tonantzintla y Tacubaya, 4, 86
- Renzo, M., Zapartas, E., de Mink, S. E., et al. 2018, ArXiv:1804.09164
- Sabbi, E., Lennon, D. J., Gieles, M., et al. 2012, ApJ, 754, L37
- Salpeter, E. E. 1955, ApJ, 121, 161
- Sana, H., de Mink, S. E., de Koter, A., et al. 2012, Science, 337, 444
- Sana, H. & Evans, C. J. 2011, in IAU Symposium, Vol. 272, Active OB Stars: Structure, Evolution, Mass Loss, and Critical Limits, ed. C. Neiner, G. Wade, G. Meynet, & G. Peters, 474–485
- Schneider, F. R. N., Sana, H., Evans, C. J., et al. 2018, Science, 359, 69
- Zapartas, E., de Mink, S. E., Izzard, R. G., et al. 2017, A&A, 601, A29
- Zinnecker, H. & Yorke, H. W. 2007, ARA&A, 45, 481
- Zwicky, F. 1957, ZAp, 44, 64