

## SPACE ASTROMETRY OF THE VERY MASSIVE $\sim 150 M_{\odot}$ CANDIDATE RUNAWAY STAR VFTS682

M. RENZO<sup>\*1</sup>, S. E. DE MINK<sup>1</sup>, D. J. LENNON<sup>2</sup>, I. PLATAIS<sup>3</sup>, R. P. VAN DER MAREL<sup>3,4</sup>, E. LAPLACE<sup>1</sup>, J. M. BESTENLEHNER<sup>5</sup>, C. J. EVANS<sup>6</sup>, V. HÉNAULT-BRUNET<sup>7</sup>, S. JUSTHAM<sup>1,8,9</sup>, A. DE KOTER<sup>1</sup>, N. LANGER<sup>10</sup>, F. NAJARRO<sup>11</sup>, H. SANA<sup>12</sup>, F. R. SCHNEIDER<sup>13</sup>, J. S. VINK<sup>14</sup>

\* Corresponding author: [m.renzo@uva.nl](mailto:m.renzo@uva.nl). A list of the affiliations can be found at the end of this paper.

*Draft version August 23, 2018*

### ABSTRACT

How very massive stars form is still an intriguing open question in stellar astrophysics. VFTS682 is among the most massive stars known, with an inferred initial mass of  $\sim 150 M_{\odot}$ . It is located in 30 Doradus at a projected distance of 29 pc from the central cluster R136. The absence of other massive stars in its immediate vicinity led to two intriguing hypotheses: either it formed in relative isolation or it was ejected dynamically from the central cluster. In this context, we investigate the kinematics of VFTS682 as obtained by *Gaia* and multi-epoch *Hubble Space Telescope* astrometry. We derive a projected velocity relative to the cluster of  $38 \pm 17 \text{ km s}^{-1}$  ( $1\sigma$  confidence interval). Although the error bars are substantial, two completely independent measures suggest that VFTS682 might be a slow runaway ejected from the central cluster, given its distance and the inferred age. This hypothesis is further supported by a variety of circumstantial clues. The central cluster is known to harbor several other stars more massive than  $150 M_{\odot}$  of similar in spectral type. The very massive O stars VFTS16 (and possibly VFTS72), which has recently been proposed as a dynamically ejected runaway, provides direct evidence that R136 is indeed capable of ejecting some of its most massive members, consistent with the predictions of numerical simulations. If future data confirm the runaway nature, this would make VFTS682 the most massive runaway star known to date.

*Subject headings:* stars: kinematics, stars: runaways, stars: individual: VFTS682

### 1. INTRODUCTION

How massive stars form is one of the major longstanding questions in astrophysics (e.g., [Zinnecker & Yorke 2007](#)). Improving our understanding of massive star formation, and its possible dependence on environment and metallicity, is crucial for understanding the role massive stars play within their host galaxies, but also for understanding the transients that mark their death and the compact remnants they leave behind. Obtaining clues from observations is challenging, because massive stars are intrinsically rare, evolve fast, typically reside in dense groups, and remain enshrouded in their parent cloud during the entirety of the formation process. Important progress has been made on the theoretical side, (e.g. [Bate 2009](#); [Kuiper et al. 2015](#); [Rosen et al. 2016](#)), but the simulations of this multi-scale and multi-physics problem are computationally expensive and remain challenging.

It has been proposed that most, if not all, stars form in clusters ([Lada & Lada 2003](#)). In this picture, field stars are primarily the result of the dissolution of dense groups. However, a significant population of massive stars exists in relative isolation, far from dense clusters or OB associations and their origin remains a matter of debate ([Gvaramadze et al. 2012](#); [Lamb et al. 2016](#); [Ward & Kruijssen 2018](#)). One hypothesis to explain the population of relatively isolated massive stars is that they formed in the field (e.g., [Parker & Goodwin 2007](#)). Another, not necessarily mutually exclusive, hypothesis is that these massive stars were ejected from the clusters in which they formed. Such ejections may result from dynamical interactions (e.g., [Poveda et al. 1967](#)) or from the disruption of binary systems at the death of the companion star (e.g., [Blaauw 1961](#); [Renzo et al. 2018](#)).

One of the most extreme examples that has been considered in this debate is the very massive star VFTS682 ([Bestenlehner et al. 2011](#); [Bressert et al. 2012](#)). This star is located in the field of the 30 Doradus region in the Large

Magellanic Cloud (LMC) and was studied as part of the multi-epoch spectroscopic VLT-FLAMES Tarantula Survey (VFTS, [Evans et al. 2011](#)). It is a hydrogen-rich Wolf-Rayet star of spectral type WNh5. Spectral analysis and comparison with evolutionary models lead to an inferred present-day mass of  $\sim 137.8^{+27.5}_{-15.9} M_{\odot}$  corresponding to an initial mass of  $\sim 150.0^{+28.7}_{-17.4} M_{\odot}$  ([Schneider et al. 2018](#)). This makes VFTS682 one of the most massive stars known and one of the most extreme objects in the region. From the spectral point of view, it is reminiscent of the very massive stars in the core of the R136 cluster ([de Koter et al. 1997](#); [Crowther et al. 2010, 2016](#)). In particular, a remarkable similarity exist between the spectrum of VFTS682 and R136a3 ([Rubio-Díez et al. 2017](#)).

VFTS682 stands out by its relative isolation at a projected distance of 119.4 arcseconds, corresponding to  $\sim 29$  pc, from the star cluster R136. [Bestenlehner et al. \(2011\)](#) considered two possible explanations for the offset: either the star formed in situ as an isolated massive star, or it was ejected from R136. N-body simulations indicate that the ejection of very massive stars like VFTS682 is expected (e.g. [Fujii & Portegies Zwart 2011](#); [Banerjee et al. 2012](#)). The capability of young cluster of ejecting a large number of (very) massive stars is supported by the recent findings of proper motion studies ([Lennon et al. 2018](#); [Drew et al. 2018](#)).

[Platais et al. \(2015, 2018\)](#) analyzed multi-epoch *Hubble Space Telescope* (HST) photometry and identified 10 stars likely ejected from R136. [Lennon et al. \(2018\)](#) investigate the kinematics of isolated O-type stars in the region using the second *Gaia* data release (DR2, [Gaia Collaboration et al. 2016, 2018](#)) and show that the proper motion, position and direction of the  $\sim 100 M_{\odot}$  star VFTS16 is consistent with a runaway origin from R136. They found a less clear case for VFTS72, and in both cases some tension between the kinematic age of these stars and their apparent age remains.

In this paper we present an analysis of the new kinemati-

TABLE 1  
STELLAR PARAMETERS OF VFTS682.

Parameter	Units	Value	Ref.
present day mass	$[M_\odot]$	$137.8^{+27.5}_{-15.9}$	(1)
initial mass	$[M_\odot]$	$150.0^{+28.7}_{-17.4}$	(1)
age	[Myr]	$1.0 \pm 0.2$	(1)
mass loss rate	$\log_{10}(\dot{M}/[M_\odot \text{ yr}^{-1}])$	$-4.1 \pm 0.2$	(2)

NOTE. — The quoted uncertainties are statistical, and do not include systematic effects in the modeling.

(1) Schneider et al. (2018) (2) Bestenlehner et al. (2011)

cal constraints for VFTS682 provided by *Gaia* DR2 and constraints from HST proper motions by Platais et al. (2018). We discuss the implications for the hypothesis that VFTS682 is a slow runaway star ejected from R136.

## 2. OBSERVATIONS

The WNh5 star VFTS682, located at right ascension (RA)  $05^{\text{h}}38^{\text{m}}55.510^{\text{s}}$  and declination (DEC)  $-69^{\circ}04'26.72''$  J2000, was observed as part of the multi-epoch, spectroscopic VFTS campaign covering  $\lambda 4000\text{--}7000$  (Evans et al. 2011). Bestenlehner et al. (2011) analyzed the spectra to infer the stellar parameters and measure an extinction of  $A_V = 4.45 \pm 0.12$ , implying a luminosity of  $\log_{10}(L/L_\odot) = 6.5 \pm 0.2$ , making this one of the most luminous stars in the region. The absence of radial velocity variations suggests that the star is unlikely to have close binary companions (Bestenlehner et al. 2011), unless the orbital inclination is very high. Bayesian fits of the stellar parameters against evolutionary tracks (Brott et al. 2011; Köhler et al. 2015) using the BONNSAI code (Schneider et al. 2014, 2017) provide estimates for the age, present mass and initial mass, Table 1.

Townsley et al. (2006) did not find a bright X-ray point source at the position of VFTS682 in their *Chandra* survey of the region. The X-ray luminosity of VFTS682 is significantly lower than known massive binaries in the region, which suggests the absence of colliding winds. These would be expected in the presence of companions even for extreme mass ratios, given the extremely large mass of VFTS682. This star is also relatively isolated in the (near) infrared. The nearest bright (near-)infrared sources detected by *Spitzer* (Meixner et al. 2006) and resolved in the *VISTA* Magellanic Clouds Survey (Cioni et al. 2011) are located at a distance of about 10 arcsecond, i.e. about 2.4 pc. Walborn et al. (2013) speculate that these nearby young stars may represent a case of star formation triggered by the the wind of VFTS682.

The V-band light curve of VFTS682 shows variations at a  $\sim 10\%$  level on a timescale of years, which is unusual for Wolf-Rayet stars and more typical for Luminous Blue Variable (LBV) stars (Parker 1993; Udalski et al. 2008; Bestenlehner et al. 2011). It also shows a mid-infrared excess (Gruendl & Chu 2009).

Estimates of the radial velocity are complicated by the variable, possibly inhomogeneous, optically thick wind typical of emission line stars. Bestenlehner et al. (2011) estimate a mass loss rate of  $\log_{10}(\dot{M}/[M_\odot \text{ yr}^{-1}]) = -4.1 \pm 0.2$ , not accounting for the possible effect of clumping. We therefore caution for over-interpreting the radial velocities estimates. Bestenlehner et al. (2011) estimate a radial velocity (RV) of  $300 \pm 10 \text{ km s}^{-1}$  using the Nv  $\lambda 4944$  line, which is offset from the average radial velocity of the region of  $270 \pm 10 \text{ km s}^{-1}$ . This was suggested as indicating a runaway nature, but it is no proof of

TABLE 2  
KINEMATICS OF VFTS682.

Parameter	Units	Value	Ref.
<i>Absolute and relative position</i>			
RA <sub>VFTS682</sub>	[degrees]	84.7314	(1)
DEC <sub>VFTS682</sub>	[degrees]	-69.0741	(1)
RA <sub>R136</sub>	[degrees]	84.6767	SIMBAD
DEC <sub>R136</sub>	[degrees]	-69.1009	SIMBAD
$\delta\text{RA}$	[mas]	0.0547	(2, 5)
$\delta\text{DEC}$	[mas]	0.0268	(2, 5)
$d_{\parallel}$	[arcsec]	119.4	SIMBAD
$L_{\parallel}$	[pc]	29	(2)
<i>Gaia absolute proper motion for VFTS682 and the region</i>			
$\mu_{\text{RA}}$	[mas yr $^{-1}$ ]	$1.843 \pm 0.070$	(1)
$\mu_{\text{DEC}}$	[mas yr $^{-1}$ ]	$0.786 \pm 0.080$	(1)
$\rho(\mu_{\text{RA}}, \mu_{\text{DEC}})$		0.0226	(1)
$\langle \mu_{\text{RA}} \rangle_{\text{R136}}$	[mas yr $^{-1}$ ]	$1.74 \pm 0.01$	(3)
$\langle \mu_{\text{DEC}} \rangle_{\text{R136}}$	[mas yr $^{-1}$ ]	$0.70 \pm 0.02$	(3)
<i>Gaia relative proper motion for VFTS682</i>			
$\delta\mu_{\text{RA}}$	[mas yr $^{-1}$ ]	$0.10 \pm 0.08$	(1, 5)
$\delta\mu_{\text{DEC}}$	[mas yr $^{-1}$ ]	$0.086 \pm 0.100$	(1, 5)
$\delta\mu_{\text{Gaia}}$	[mas yr $^{-1}$ ]	$0.13 \pm 0.09$	(1, 5)
$v_{2D}$	[km s $^{-1}$ ]	$32 \pm 21$	(1, 5)
$\theta_{\text{Gaia}}$	[degrees]	$14^{+36}_{-31}$	(1, 5)
<i>HST relative proper motion for VFTS682</i>			
$\delta\mu_{\text{RA,HST}}$	[mas yr $^{-1}$ ]	$0.01 \pm 0.13$	(4)
$\delta\mu_{\text{DEC,HST}}$	[mas yr $^{-1}$ ]	$0.20 \pm 0.10$	(4)
$\delta\mu_{\text{HST}}$	[mas yr $^{-1}$ ]	$0.20 \pm 0.10$	(4)
$v_{2D,\text{HST}}$	[km s $^{-1}$ ]	$47 \pm 24$	(4)
$\theta_{\text{HST}}$	[degrees]	$-35^{+24}_{-51}$	(1, 5)
<i>Weighted average relative proper motion for VFTS682</i>			
$\delta\mu_{\text{RA,avg}}$	[mas yr $^{-1}$ ]	$0.08 \pm 0.07$	(5)
$\delta\mu_{\text{DEC,avg}}$	[mas yr $^{-1}$ ]	$0.14 \pm 0.07$	(5)
$\delta\mu_{\text{avg}}$	[mas yr $^{-1}$ ]	$0.16 \pm 0.07$	(5)
$v_{2D,\text{avg}}$	[km s $^{-1}$ ]	$38 \pm 17$	(5)
<i>Expected proper motion if ejected from R136 at age zero</i>			
$v_{2D}$	[km s $^{-1}$ ]	$29 \pm 6$	(2)
$\theta$	[degrees]	$\sim 0$	

NOTE. — Proper motions are based on Gaia data unless indicated otherwise in subscript. The conversion factor assuming a distance of 50 kpc is such that 1 mas yr $^{-1}$  corresponds to 237 km s $^{-1}$ . The correlation coefficient is indicated as  $\rho$ . The angle  $\theta$  is defined such that  $\theta = 0$  for motions pointing away radially from R136, we neglect the errors on  $\langle \mu_{\text{RA}} \rangle_{\text{R136}}$  and  $\langle \mu_{\text{DEC}} \rangle_{\text{R136}}$  to determine the uncertainty on  $\theta_{\text{Gaia}}$ . (1) Gaia Collaboration et al. (2018), (2) Bestenlehner et al. (2011), (3) Lennon et al. (2018), (4) Platais et al. (2018) and (5) this study.

it. Bressert et al. (2012) note an offset compared between the RV of the star and the nebular lines from the gas filaments in its vicinity. This is consistent with the expectation if the star was not formed in situ. We will refrain from using the RV measurements in this work, and focus on the velocities on the plane of the sky.

We adopt a distance to the LMC of 50 kpc. The error on the distance determination is small ( $\lesssim 2\%$ , Pietrzyński et al. 2013) and any possible offset in the radial direction between R136 or VFTS682 and the distance we adopted for the LMC is probably much smaller ( $\sim 0.5\%$ , e.g., Luks & Rohlfs 1992). These uncertainties are negligible compared to the errors in the proper motion discussed below.

### 2.1. Gaia astrometry for VFTS682

VFTS682 is identified with the source id 4657685637907503744 in the *Gaia* DR2 catalog as a




FIG. 1.— *Left panel:* distribution of OB-type and Wolf-Rayet stars in proper motion relative to R136 from *Gaia* DR2. VFTS682 is not an outlier, but its relative proper motion matches the expected value (green vertical band) if it were indeed ejected from R136 assuming an age of  $1.0 \pm 0.2$  Myr. The top axis shows the conversion to  $\text{km s}^{-1}$  assuming a distance of 50 kpc. *Right panel:* distribution of angles between the relative proper motion direction and the radial direction from the center of the cluster to the star. The error bars from *Gaia* for VFTS682 are large, but the best value is in agreement with the hypothesis of dynamical ejection (green vertical band). The peak at  $\theta \approx \pi/2$  is due to stars belonging to NGC2060. In both panels, the dark blue histograms contain 317 stars with error smaller than  $0.1 \text{ mas yr}^{-1} \approx 25 \text{ km s}^{-1}$  at 50 kpc, the lighter blue histograms contain 36 stars with errors smaller than  $0.05 \text{ mas yr}^{-1}$ .




FIG. 2.— The thick yellow arrows indicate proper motions relative to R136 (blue circle) from *Gaia* DR2, multiplied by 0.4 Myr. The prolongations in the opposite direction are proportional to the apparent ages of the stars, and the thin lines illustrate the error cone on the potential origin. The green thick arrow indicates the HST proper motion of VFTS682, with the relative error cone and prolongation proportional to its apparent age. The two blue circles indicate the regions of radii 0.01 and 0.05 degrees around the core of R136.

15.65 mag star in the G band ([Gaia Collaboration et al. 2016, 2018](#)). The number of visibility periods, i.e., groups of observations separated from other groups by a gap of at least four days, used in the astrometric solution is seventeen. The reported astrometric excess noise is zero. These values suggest that the *Gaia* DR2 data for VFTS682 are reliable.

*Gaia* provides absolute proper motions. To determine the relative proper motion with respect to R136, we follow [Lennon et al. \(2018\)](#) to define the motion of the local frame of reference using the average proper motion of nearby stars with reliable astrometric data. They selected bright ( $G < 17$ ) stars within 0.05 degrees of R136 and exclude sources with proper motion errors greater than  $0.01 \text{ mas yr}^{-1}$  in both coordinates (see their Sect. 2.1). Using this definition of the local frame, we compute the relative proper motion  $\delta\mu_{RA}$  and  $\delta\mu_{DEC}$ . We also compute the total projected 2D velocity  $v_{2D}$  and the angle  $\theta$  between the direction of motion and the vector connecting the center of R136 with the current position of VFTS682, i.e. such that  $\theta = 0$  is the angle corresponding to perfectly radial motion away from R136. All kinematic quantities are provided in Table 2.

## 2.2. HST astrometry for VFTS682

The 30 Doradus region was targeted by a two-epoch photometric campaign with HST providing observations in the F775W filter in October 2011 and October 2014 (GO-12499; P.I.: D. J. Lennon, [Sabbi et al. 2013](#)). [Platais et al. \(2015, 2018\)](#) analyzed the HST data to determine the relative proper motions and identify candidate runaway stars. The brightest stars ( $V < 14$ ) are saturated in the data set and have been excluded from the analysis. The high extinction around VFTS682 prevents this issue. VFTS682 is relatively slow and is therefore not discussed in [Platais et al. 2018](#). However, their measurements provide useful complementary estimate of the proper motion of VFTS682 independent from the *Gaia* data.

The HST study provides proper motions that are relative to the bulk motion of the majority of the stars in the field of view. The full 30 Dor field is covered by different pointings and there is some systematic distortion. However, even for stars far from 30 Dor the effect is small, no more than  $0.05 \text{ mas yr}^{-1}$  across the whole 30 Dor field. The effect is much smaller for stars close to the center of field, such as VFTS 682 ([Platais et al. 2018](#)). We can therefore use the relative proper motion (Table 2) as a good estimate for the proper motion relative to R136.

## 3. THE KINEMATICS OF VFTS682

The black points in Fig. 1 show the proper motion (left panel) and the projected flight direction (right panel) relative to R136 of VFTS682 from *Gaia* DR2.  $\theta$  is the angle between the projected radial direction from the center of R136 to each star and its projected relative proper motion (see also Fig. 2). Dynamical ejections from the cluster should produce close to radial ejections, i.e.  $\theta \approx 0$ . The green vertical bands highlight the expectations for these two quantities if VFTS682 were ejected from R136 early in its lifetime. For comparison, we also show the relative proper motion of VFTS16 and VFTS72 (gray points), and the distribution in relative proper motion and flight direction for all the VFTS OB-type and Wolf-Rayet stars with *Gaia* DR2 errors on the proper motion components of less than  $0.1 \text{ mas yr}^{-1}$  (dark blue lines, including VFTS682), and less than  $0.05 \text{ mas yr}^{-1}$  (light hatched blue). Although the error bars are substantial and VFTS682 is not an outlier compared to other OB-type and Wolf-Rayet

stars, the agreement suggests that the star is indeed a “slow-runaway” as suggested by [Bestenlehner et al. \(2011\)](#).

Subtracting the mean motion of R136, we obtain relative proper motions (projected velocities) of  $\delta\mu_{Gaia} = 0.13 \pm 0.09 \text{ mas yr}^{-1}$  ( $32 \pm 21 \text{ km s}^{-1}$ ) and  $\delta\mu_{HST} = 0.20 \pm 0.10 \text{ mas yr}^{-1}$  ( $47 \pm 24 \text{ km s}^{-1}$ ). Both values are consistent with each other, but also with no relative motion within  $2\sigma$ . The weighted average of these two independent measurements is  $\delta\mu_{avg} = 0.16 \pm 0.07 \text{ mas yr}^{-1}$  ( $38 \pm 17 \text{ km s}^{-1}$ ).

Figure 2 shows the projected motion of VFTS682 relative to R136 on the sky. We also show VFTS16 and VFTS72 for completeness (see [Lennon et al. 2018](#)). The thick yellow arrows are proportional to the relative proper motion from *Gaia* DR2, and the thick green line illustrates the relative proper motion from HST. The error cone on the direction of motion is illustrated by the corresponding prolongation in the direction opposite to the motion, and we also show the most likely origin of the stars accounting for their apparent age. This figure illustrates that R136 is the most likely origin of these stars, although the large error bars prevent a robust identification for VFTS682, and there is some tension between the apparent age and the present day distance from the cluster core for VFTS16 and VFTS72 ([Lennon et al. 2018](#)). We note that accurate distances along the line of sight are needed to constrain the flight direction in three dimensions.

Assuming VFTS682 indeed originates from R136, we can calculate its kinematic age as:

$$\tau_{kin} = \frac{d_{||}}{\delta\mu_{avg}} \simeq \frac{119.4 \text{ arcsec}}{0.16 \text{ mas yr}^{-1}} \simeq 0.7 \pm 0.3 \text{ Myr} , \quad (1)$$

where  $d_{||} = 119.4 \text{ arcsec}$  is the angular distance from VFTS682 to the core of the cluster ([Bestenlehner et al. 2011](#)). The kinematic age  $\tau_{kin}$  is consistent with a very early ejection from the cluster (see Table 1).

In summary, both *Gaia* and HST relative proper motions are consistent with the dynamical ejection of VFTS682 from the cluster, although we cannot confidently rule out the counter hypothesis.

## 4. DISCUSSION

Based on our results, we consider it plausible that VFTS682 is the most massive runaway known to date, with a two-dimensional projected velocity with respect to R136 of  $38 \pm 17 \text{ km s}^{-1}$  (averaging the *Gaia* DR2 and HST results). Due to the large error bars, this result will need to be revisited with future astrometric data.

If confirmed, it means that isolated star formation is *not* required to explain the isolation of VFTS682. Its proper motion suggests that it was ejected from the cluster R136  $0.7 \pm 0.3 \text{ Myr}$  ago. If the cluster age ( $\lesssim 2 \text{ Myr}$ , [Crowther et al. 2010; Sabbi et al. 2012](#)) is indeed smaller than the shortest stellar lifetime ( $\sim 3 \text{ Myr}$ , [Brott et al. 2011; Köhler et al. 2015; Zapartas et al. 2017](#)), the ejection of VFTS682 from the disruption of a binary is excluded. The kinematic age we infer is smaller than the kinematic age of  $\sim 1.5 \text{ Myr}$  for VFTS16 found in [Lennon et al. \(2018\)](#), which indicates that VFTS682 was ejected later than VFTS16, and potentially later than VFTS72 too.

If the star were ejected dynamically, its isolation makes it an ideal target to constrain the stellar physics of stars with masses well above  $\sim 100 M_\odot$  while avoiding crowding issues. Moreover, its exceptionally large mass raises the question of which stars must populate the core of the cluster. N-body

dynamics typically ejects the least massive star among those interacting (e.g., [Banerjee et al. 2012](#)). Just based on the kinematic properties of VFTS682, we would expect several stars with initial masses larger than  $\sim 150 M_{\odot}$  in the cluster R136, which is consistent with the detection of extremely massive stars in the core of the cluster.

The simulations of [Fujii & Portegies Zwart \(2011\)](#) suggest that early in the evolution of a cluster, dynamical interactions form an extremely massive binary, which then tightens its orbit by ejecting other stars. The spectral similarities between VFTS682 and stars in the core of R136 are in agreement with this “bully binary” model. Interpreting the kinematics of VFTS682 through the lens of their simulations suggests the presence of a close binary with total mass  $M_1 + M_2 \gtrsim 300 M_{\odot}$  in the core of the cluster. The kinematic age of VFTS682 puts an upper limit to the timescale to form the “bully binary” in R136. If identified, such a binary might be an ideal observational candidate for a dynamically formed progenitor system of a binary black-hole, provided that stars this massive can avoid a pair-instability supernova (e.g., [Rakavy & Shaviv 1967](#)) at LMC metallicity (see also [Langer et al. 2007](#)).

Similarly, the final fate of VFTS682 could be either a pair-instability supernova without compact remnant formation, or possibly direct collapse to a black hole above the pair-instability mass gap. The amount of mass loss of these stars will determine their final core mass and thus their final fate.

Also [Banerjee et al. \(2012\)](#) suggested, based on their N-body simulations, that VFTS682 was ejected from R136. In their model, they relied on (dynamically driven) stellar mergers to explain the high masses of VFTS682 and the massive members of R136. They demonstrated that the cluster potential does not significantly change the velocity of the star after the ejection. To eject such a massive object, the cluster is expected to have produced a large number of massive runaways. Indeed, several isolated massive stars are observed in the region ([Evans et al. 2010; Lennon et al. 2018](#)), some with known large radial velocities and/or proper motion. A comprehensive study of the kinematic properties of all the massive stars surrounding R136 could shed light on whether some can be unequivocally identified as merger products, but also on the initial conditions for the cluster dynamics (e.g., [Oh & Kroupa 2016](#)), and whether it formed via a monolithic collapse, or as a (potentially ongoing) merger of several sub-structures (e.g., [Sabbí et al. 2012](#)).

VFTS682 is potentially the most massive runaway known to date, and its possible ejection from the cluster R136 likely implies that it is only the “tip of the iceberg” of massive runaways in the region. Studies of this population, enabled by recent HST and *Gaia* observations will put constraints on the evolution of these extreme stars, together with the formation and evolution of the central cluster itself.

## REFERENCES

- Banerjee, S., Kroupa, P., & Oh, S. 2012, *ApJ*, 746, 15  
 Bate, M. R. 2009, *MNRAS*, 392, 590  
 Bestenlehner, J. M., Vink, J. S., Gräfener, G., et al. 2011, *A&A*, 530, L14  
 Blaauw, A. 1961, *Bull. Astron. Inst. Netherlands*, 15, 265  
 Bressert, E., Bastian, N., Evans, C. J., et al. 2012, *A&A*, 542, A49  
 Brott, I., de Mink, S. E., Cantiello, M., et al. 2011, *A&A*, 530, A115  
 Cioni, M.-R. L., Clementini, G., Girardi, L., et al. 2011, *A&A*, 527, A116  
 Crowther, P. A., Schnurr, O., Hirschi, R., et al. 2010, *MNRAS*, 408, 731  
 Crowther, P. A., Caballero-Nieves, S. M., Bostroem, K. A., et al. 2016, *MNRAS*, 458, 624  
 de Koter, A., Heap, S. R., & Hubeny, I. 1997, *ApJ*, 477, 792  
 Drew, J. E., Herrero, A., Mohr-Smith, M., et al. 2018, *MNRAS*, arXiv:1807.06486; 1807.06486  
 Evans, C. J., Walborn, N. R., Crowther, P. A., et al. 2010, *ApJ*, 715, L74  
 Evans, C. J., Taylor, W. D., Hénault-Brunet, V., et al. 2011, *A&A*, 530, A108  
 Fujii, M. S., & Portegies Zwart, S. 2011, *Science*, 334, 1380  
 Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, *ArXiv:1804.09365*, arXiv:1804.09365  
 Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., et al. 2016, *A&A*, 595, A1  
 Gruendl, R. A., & Chu, Y.-H. 2009, *ApJS*, 184, 172  
 Gvaramadze, V. V., Weidner, C., Kroupa, P., & Pflamm-Altenburg, J. 2012, *MNRAS*, 424, 3037  
 Köhler, K., Langer, N., de Koter, A., et al. 2015, *A&A*, 573, A71  
 Kuiper, R., Yorke, H. W., & Turner, N. J. 2015, *ApJ*, 800, 86  
 Lada, C. J., & Lada, E. A. 2003, *ARA&A*, 41, 57  
 Lamb, J. B., Oey, M. S., Segura-Cox, D. M., et al. 2016, *ApJ*, 817, 113  
 Langer, N., Norman, C. A., de Koter, A., et al. 2007, *A&A*, 475, L19  
 Lennon, D. J., Evans, C. J., van der Marel, R. P., et al. 2018, *ArXiv:1805.08277*, arXiv:1805.08277  
 Luks, T., & Rohlfs, K. 1992, *A&A*, 263, 41  
 Meixner, M., Gordon, K. D., Indebetouw, R., et al. 2006, *AJ*, 132, 2268

## AFFILIATIONS

<sup>1</sup>Astronomical Institute Anton Pannekoek, University of Amsterdam, 1098 XH Amsterdam, The Netherlands

<sup>2</sup>ESA, European Space Astronomy Centre, Apdo. de Correos 78, E-28691 Villanueva de la Cañada, Madrid, Spain

<sup>3</sup> Department of Physics & Astronomy, Johns Hopkins University, Baltimore, MD 21218, USA

<sup>4</sup> Space Telescope Science Institute, 3700 San Martin Drive,

Baltimore, MD 21218, USA

<sup>5</sup>Department of Physics and Astronomy, Hicks Building, Hounsfield Road, University of Sheffield, Sheffield S3 7RH, UK

<sup>6</sup>UK Astronomy Technology Centre, Royal Observatory Edinburgh, Blackford Hill, Edinburgh, EH9 3HJ, UK

<sup>7</sup>National Research Council, Herzberg Astronomy & Astrophysics, 5071 West Saanich Road, Victoria, BC, V9E 2E7,

Canada

<sup>8</sup> School of Astronomy & Space Science, University of the Chinese Academy of Sciences, Beijing 100012, China

<sup>9</sup> National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100012, China

<sup>10</sup> Argelander-Institut für Astronomie, Universität Bonn, Auf dem Hügel 71, 53121, Bonn, Germany

<sup>11</sup> Centro de Astrobiología, CSIC-INTA, Carretera de Torrejón a Ajalvir km-4, E-28850 Torrejón de Ardoz, Madrid, Spain

<sup>12</sup> Institute of Astronomy, KU Leuven, Celestijnenlaan 200 D, B-3001 Leuven, Belgium

<sup>13</sup> Department of Physics, University of Oxford, Keble Road, Oxford OX1 3RH, UK

<sup>14</sup> Armagh Observatory, College Hill, Armagh BT61 9DG, UK

We are grateful to J. Heyl, M. C. Ramirez-Tannus, S. N. Shore, and S. Torres for help and discussions. We also thank the HST-PROMO (High-resolution Space Telescope PROPer MOTion) Collaboration for sharing and discussing ideas relevant to this work. SdM acknowledges the European Unions Horizon 2020 research and innovation programme from the European Research Council (ERC), Grant agreement No. 715063 and VHB the NRC-Canada Plaskett Fellowship. This work has made use of data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement.