

2.24 Use  $S$  to be the event "is smoker"  
 $W$  to be the event "is a woman"  
 $M$  to be the event "is a man"  
 $C$  to be the event "has cancer"

We have been told

$$\Pr[C | M \cap S] = 23 \cdot \Pr[C | M \cap S^c],$$

$$\Pr[C | W \cap S] = 13 \cdot \Pr[C | W \cap S^c],$$

$$\Pr[S | M] = 0.231, \Pr[S | W] = 0.183 \Rightarrow \Pr[S^c | M] = 0.769, \Pr[S^c | W] = 0.817.$$

$$(a) \Pr[W | S] = \frac{\# \text{ of women who smoke}}{\# \text{ of smokers}} = \frac{21.1}{21.1 + 24.8} = 0.46$$

$$\begin{aligned} (b) \Pr[S | W \cap C] &= \frac{\Pr[S \cap W \cap C]}{\Pr[W \cap C]} = \frac{\Pr[C | S \cap W] \cdot \Pr[S \cap W]}{\Pr[C \cap W \cap S] + \Pr[C \cap W \cap S^c]} \\ &= \frac{\Pr[C | S \cap W] \Pr[S \cap W]}{\Pr[C | S \cap W] \cdot \Pr[S \cap W] + \Pr[C | S^c \cap W] \cdot \Pr[S^c \cap W]} \\ &= \frac{13 \cdot \Pr[C | S^c \cap W] \Pr[S \cap W]}{13 \cdot \Pr[C | S^c \cap W] \cdot \Pr[S \cap W] + \Pr[C | S^c \cap W] (1 - \Pr[S \cap W])} \\ &= \frac{13 \Pr[S \cap W]}{13 \Pr[S \cap W] + \Pr[S^c \cap W]} = \frac{13 \Pr[S | W] \Pr[W]}{13 \Pr[S | W] \Pr[W] + \Pr[S^c | W] \Pr[W]} \\ &= \frac{13 \Pr[S | W]}{13 \Pr[S | W] + \Pr[S^c | W]} = \frac{13(0.183)}{13(0.183) + 0.817} \approx 0.744 \end{aligned}$$

(c) Similarly to part (b),

$$\Pr[S | M \cap C] = \frac{23(0.231)}{23(0.231) + 0.769} = \boxed{0.874}$$

$$2.51 \quad X \sim \text{Geom}(\pi) \Rightarrow \Pr(X = x) = (1 - \pi)^{x-1} \pi$$

$$\begin{aligned} (a) \Pr(X \geq k) &= [(1 - \pi)^k + (1 - \pi)^{k+1} + \dots] \pi \\ &= (1 - \pi)^k \pi [1 + (1 - \pi) + (1 - \pi)^2 + \dots] = \frac{(1 - \pi)^k \pi}{1 - (1 - \pi)} = (1 - \pi)^k \end{aligned}$$

$$\begin{aligned}
 (b) \quad \Pr(X=x \mid X \geq k) &= \frac{\Pr(X \geq k \text{ and } X=x)}{\Pr(X \geq k)} \\
 &= \begin{cases} 0, & x < k \\ \frac{\pi(1-\pi)^x}{(1-\pi)^k}, & x \geq k \end{cases} = \begin{cases} 0, & x < k \\ \pi(1-\pi)^{x-k}, & x \geq k \end{cases} = \Pr(X=x-k).
 \end{aligned}$$

(c) Saying  $X \geq k$  is like starting over.

$$2.62 \quad (a) \quad X=2: \quad \binom{4}{2} \left[ \binom{26}{5} - 2 \binom{13}{5} \right] / \binom{52}{5} = 0.1459$$

$$X=4: \quad \binom{4}{1} \binom{13}{2} \binom{13}{1}^3 / \binom{52}{5} = 0.2637$$

$$X=3: \quad 1 - (\Pr(X=1) + \Pr(X=2) + \Pr(X=4)) = 0.5884$$

$x$	$\Pr(X=x)$
1	0.00198
2	0.1459
3	0.5884
4	0.2637

$$(b) \quad E(X) = (0.00198) + (2)(0.1459) + (3)(0.5884) + (4)(0.2637) = 3.114$$

2.97  $\text{hyper}(2, 12, 18, 17)$  yields the same answer. So do

$1 - \text{hyper}(14, 17, 13, 18)$  and  $\text{hyper}(3, 18, 12, 13)$ .

$$3.5 \quad \text{For } X \sim \exp(\lambda), \text{ we have cdf } F_X(x) = \begin{cases} 0, & x < 0 \\ 1 - e^{-\lambda x}, & x \geq 0 \end{cases}$$

For the median, we solve  $0.5 = 1 - e^{-\lambda x} \Rightarrow x = \frac{1}{\lambda} \ln 2$ .

The first quartile  $x$  satisfies  $0.25 = 1 - e^{-\lambda x} \Rightarrow x = -\frac{1}{\lambda} \ln(3/4)$ .

The third quartile  $x$  satisfies  $0.75 = 1 - e^{-\lambda x} \Rightarrow x = \frac{2}{\lambda} \ln 2$ .

$$3.22 \quad \text{For } X \sim \text{Geom}(\pi), \quad f_X(x) = (1-\pi)^x \pi$$

$$\begin{aligned}
 \Rightarrow M_X(t) &= E(e^{tX}) = \sum_{x=0}^{\infty} e^{tx} (1-\pi)^x \pi = \pi \sum_{x=0}^{\infty} [e^t(1-\pi)]^x \\
 &= \pi \cdot \frac{1}{1 - e^t(1-\pi)}
 \end{aligned}$$

$$\begin{aligned}
3.23 \quad M_Y(t) &= \int_{-\infty}^{\infty} e^{ty} f(y) dy = \int_0^1 y e^{ty} dy + \int_1^2 (2-y) e^{ty} dy \\
&= \left. \frac{1}{t} y e^{ty} \right|_0^1 - \frac{1}{t} \int_0^1 e^{ty} dy + \left. \frac{1}{t} (2-y) e^{ty} \right|_1^2 + \frac{1}{t} \int_1^2 e^{ty} dy \\
&= \frac{1}{t} e^t - \frac{1}{t^2} [e^{ty}]_0^1 - \frac{1}{t} e^t + \frac{1}{t^2} [e^{ty}]_1^2 \\
&= \frac{1}{t^2} (1 - e^t) + \frac{1}{t^2} (e^{2t} - e^t) = \frac{1}{t^2} (1 - 2e^t + e^{2t}) = \frac{1}{t^2} (1 - e^t)^2.
\end{aligned}$$

$$\begin{aligned}
3.31 \quad M'_X(t) &= 2e^{2t}(1-t^2)^{-1} + 2te^{2t}(1-t^2)^{-2} \rightarrow E(X) = M'_X(0) = 2 \\
M''_X(t) &= 4e^{2t}(1-t^2)^{-1} + 8te^{2t}(1-t^2)^{-2} + 2e^{2t}(1-t^2)^{-2} + 8t^2e^{2t}(1-t^2)^{-3} \\
&\rightarrow E(X^2) = M''_X(0) = 6
\end{aligned}$$

Thus,  $\text{Var}(X) = 6 - 2^2 = 2.$

$$\begin{aligned}
3.33 \quad M'_X(t) &= \frac{18}{(3-t)^3} \rightarrow E(X) = M'_X(0) = 2/3 \\
M''_X(t) &= \frac{54}{(3-t)^4} \rightarrow E(X^2) = M''_X(0) = 2/3 \\
\text{So, } \text{Var}(X) &= \frac{2}{3} - \left(\frac{2}{3}\right)^2 = \frac{2}{9}.
\end{aligned}$$

3.37 (a) Since  $X \sim \text{Binom}(n, \pi)$  has MGF  $M_X(t) = (1 - \pi + \pi e^t)^n$ ,  
when  $M_X(t) = \left(\frac{1}{2}(e^t + 1)\right)^{10}$ ,  $X \sim \text{Binom}(10, 1/2).$

(b) Since  $X \sim \text{Norm}(\mu, \sigma)$  has MGF  $M_X(t) = e^{\mu t + \sigma^2 t^2/2}$ ,  
when  $M_X(t) = e^{t + t^2/2}$ ,  $X \sim \text{Norm}(1, 1).$

(c) Since  $X \sim \text{Exp}(\lambda)$  has MGF  $M_X(t) = \frac{1}{1 - t/\lambda}$ ,  
when  $M_X(t) = \frac{1}{1 - 2t}$ ,  $X \sim \text{Exp}(1/2).$

(d) Since  $X \sim \text{Gamma}(\alpha, \lambda)$  has MGF  $M_X(t) = \frac{1}{(1 - t/\lambda)^\alpha}$   
when  $M_X(t) = (1 - 2t)^{-3}$ ,  $X \sim \text{Gamma}(\alpha=3, \lambda=1/2)$ , or  $\text{Gamma}(\alpha=3, \beta=2).$

3.38  $X \sim \text{Gamma}(\alpha, \lambda)$ , so  $M_X(t) = \frac{1}{(1 - t/\lambda)^\alpha}$ . Setting  $Y = 3X$ , we have

$$M_Y(t) = E(e^{tY}) = E(e^{t(3X)}) = E(e^{(3t)X}) = M_X(3t) = \frac{1}{(1 - 3t/\lambda)^\alpha}$$

$$\Rightarrow Y \sim \text{Gamma}(\alpha, \lambda/3).$$

3.39 (a)  $p_{\text{exp}}(2) - p_{\text{exp}}(0) = 0.865$

(d)  $E(X) = 1/3$ ,  $\text{Var}(X) = 2/63$

(b)  $p_{\text{exp}}(1, 2) - p_{\text{exp}}(0, 2) = 0.865$

$\text{diff}(\text{pbeta}(1/3 + c(-1, 1) * \text{sqrt}(2/63)), 2, 4))$   
 $= 0.6522.$

(c)  $\frac{2}{2\sqrt{3}} \cdot (b-a) \cdot \frac{1}{b-a} = \frac{1}{\sqrt{3}} \doteq 0.5774.$

3.62 (a) Because  $R \sim \text{Norm}(100, 20)$ , his obtaining 150 would correspond to a Z-score

$$Z_R = \frac{150 - 100}{20} = 2.5$$

For  $C \sim \text{Norm}(110, 15)$ ,  $Z_C = \frac{150 - 110}{15} = 2.667$


A higher Z-score corresponds to a rarer event. Thus, Ralph should reach scores of 150 (or higher) more often than Claudia.

(b) By normality and independence,

$$R - C \sim \text{Norm}(-10, \sqrt{15^2 + 20^2}) = \text{Norm}(-10, 25)$$

$$\Pr(R > C) = 1 - \text{pnorm}(0, -10, 25)$$

$$\doteq 0.345$$



(c) Let  $\bar{R}$ ,  $\bar{C}$  be their averages over three games.

$$\bar{R} \sim \text{Norm}(100, 20/\sqrt{3}), \bar{C} \sim \text{Norm}(110, 15/\sqrt{3}) \text{ and } \bar{R} - \bar{C} \sim \text{Norm}(-10, 25/\sqrt{3}).$$

$$\Pr(\bar{R} > \bar{C}) = 1 - \text{pnorm}(0, -10, 25/\sqrt{3}) \doteq 0.244.$$

(d) Let  $X = \#$  of games won by Ralph. Assuming independence,  $X \sim \text{Binom}(3, 0.345)$ .

$$\Pr(X \geq 2) = 1 - \text{pbinom}(1, 3, 0.345) = 0.275.$$

C.4 (a)  $\text{proj}(\vec{u} \rightarrow \vec{v}) = \frac{\langle 1, 0 \rangle \cdot \langle 1, 1 \rangle}{\langle 1, 1 \rangle \cdot \langle 1, 1 \rangle} \langle 1, 1 \rangle = \frac{1}{2} \langle 1, 1 \rangle = \langle \frac{1}{2}, \frac{1}{2} \rangle.$

(b)  $\text{proj}(\vec{u} \rightarrow \vec{v}) = \frac{\langle 1, 0 \rangle \cdot \langle 1, -1 \rangle}{\langle 1, -1 \rangle \cdot \langle 1, -1 \rangle} \langle 1, -1 \rangle = \frac{1}{2} \langle 1, -1 \rangle = \langle \frac{1}{2}, -\frac{1}{2} \rangle.$

(c)  $\text{proj}(\vec{u} \rightarrow \vec{v}) = \frac{\langle 1, 0 \rangle \cdot \langle 1, 2 \rangle}{\langle 1, 2 \rangle \cdot \langle 1, 2 \rangle} \langle 1, 2 \rangle = \frac{1}{5} \langle 1, 2 \rangle = \langle \frac{1}{5}, \frac{2}{5} \rangle$

$$(d) \text{proj}(\vec{u} \rightarrow \vec{v}) = \frac{\langle 1, 2, 3 \rangle \cdot \langle 1, 1, 1 \rangle}{\langle 1, 1, 1 \rangle \cdot \langle 1, 1, 1 \rangle} \langle 1, 1, 1 \rangle = \frac{6}{3} \langle 1, 1, 1 \rangle = \langle 2, 2, 2 \rangle.$$

$$(e) \text{proj}(\vec{u} \rightarrow \vec{v}) = \frac{\langle 1, 1, 1 \rangle \cdot \langle 1, 2, 3 \rangle}{\langle 1, 2, 3 \rangle \cdot \langle 1, 2, 3 \rangle} \langle 1, 2, 3 \rangle = \frac{6}{14} \langle 1, 2, 3 \rangle = \left\langle \frac{3}{7}, \frac{6}{7}, \frac{9}{7} \right\rangle$$

$$(f) \text{proj}(\vec{u} \rightarrow \vec{v}) = \frac{\langle 1, 2, 3 \rangle \cdot \langle 1, -1, 0 \rangle}{\langle 1, -1, 0 \rangle \cdot \langle 1, -1, 0 \rangle} \langle 1, -1, 0 \rangle = -\frac{1}{2} \langle 1, -1, 0 \rangle = \left\langle -\frac{1}{2}, \frac{1}{2}, 0 \right\rangle$$

C.17 If  $A$  is  $m \times n$ , then  $A^T$  is  $n \times m$ , and the product  $AA^T$  is  $m \times m$  (square).

C.21 This statement is true. To demonstrate it, let  $B = (A^T)^{-1}$ . Then  $I = BA^T$ .

Taking transposes of both sides and noting  $I^T = I$ , we have  $I = (BA^T)^T = AB^T$ , showing that  $B^T = A^{-1}$ . Transposing again gives  $B = (A^{-1})^T$ .

$$C.24 (a) \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \begin{bmatrix} ad-bc & 0 \\ 0 & ad-bc \end{bmatrix} = (ad-bc) I.$$

(b) It is evident that  $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ , when multiplied by the now-rescaled  $\frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$ , is  $I$ .

$$(c) \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}^{-1} = -\frac{1}{2} \begin{bmatrix} 4 & -2 \\ -3 & 1 \end{bmatrix} = \begin{bmatrix} -2 & 1 \\ 3/2 & -1/2 \end{bmatrix}.$$