


3.55 (a)  $f_{X,Y}(x,y) = 1$  for  $(x,y) \in [5,6] \times [5,6]$ .

(b)  $\Pr((X,Y) \in [5,5.5] \times [5,5.5]) = 1/4$ .

(c) For them to arrive within 10 minutes of each other,  $(X,Y)$  must be a point in the orange shaded region, which has area (= probability)

$$1 - (2)(\frac{1}{2})(5/6)^2 = \frac{11}{36}.$$



3.57 (a) Let  $A = \{(x,y) \mid x^2 + y^2 \leq R^2\}$ . For the pdf to be a constant  $k$ , we require

$$1 = k \cdot \text{Area}(A) = \pi R^2 k \Rightarrow k = \frac{1}{\pi R^2}.$$

Thus


$$f(x,y) = \begin{cases} \frac{1}{\pi R^2}, & (x,y) \in A, \\ 0, & \text{otherwise} \end{cases}$$

(b) Let  $H = \{(x,y) \mid x^2 + y^2 \leq \frac{R}{2}\}$ .  $\Pr((x,y) \in H) = \frac{\text{Area}(H)}{\text{Area}(A)} = \frac{\pi(R/2)^2}{\pi R^2} = \frac{1}{4}$ .

(c)  $|X - Y| \leq R \Leftrightarrow X - R \leq Y \leq X + R$

The desired probability corresponds to the area of the orange shaded region:

$$\left(\frac{1}{2} \pi R^2 + R^2\right) \cdot \frac{1}{\pi R^2} = \frac{1}{2} + \frac{1}{\pi}.$$



(d) Let  $x \in [-R, R]$ . Then

$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x,y) dy = \int_{-\sqrt{R^2-x^2}}^{\sqrt{R^2-x^2}} \frac{1}{\pi R^2} dy = \frac{2}{\pi R^2} \sqrt{R^2-x^2}$$


(e)  $X$  and  $Y$  are not independent. By symmetry,  $f_Y(y) = \frac{2}{\pi R^2} \sqrt{R^2-y^2}$  for  $y \in [-R, R]$ .

But  $f_X(x)f_Y(y) = \frac{4}{\pi^2 R^4} \sqrt{(R^2-x^2)(R^2-y^2)} \neq \frac{1}{\pi R^2}$ .

3.58 This gives a different distribution. Under this approach,

$$\Pr((x,y) \in H) = \frac{1}{2\pi R} \cdot \text{Area}(\text{shaded region of } r\theta\text{-plane})$$

$$= \frac{1}{2\pi R} \cdot (2\pi)(\frac{1}{2}R) = \frac{1}{2}$$



a different result than in part (b) of Exercise 3.57.

3.63  $X \sim \text{Pois}(\lambda_1)$  and  $Y \sim \text{Pois}(\lambda_2)$ , so  $M_X(t) = e^{e^t \lambda_1 - \lambda_1}$ , and  $M_Y(t) = e^{e^t \lambda_2 - \lambda_2}$ .

By independence of  $X, Y$ ,

$$\begin{aligned} M_{X+Y}(t) &= M_X(t) M_Y(t) = e^{e^t \lambda_1 - \lambda_1} \cdot e^{e^t \lambda_2 - \lambda_2} \\ &= e^{e^t \lambda_1 + e^t \lambda_2 - (\lambda_1 + \lambda_2)} = e^{e^t (\lambda_1 + \lambda_2) - (\lambda_1 + \lambda_2)}. \end{aligned}$$

This is the mgf for another Poisson r.v. with parameter  $\lambda_1 + \lambda_2$ . Thus

$$X + Y \sim \text{Pois}(\lambda_1 + \lambda_2).$$

3.67 Each  $X_i \sim \text{Gamma}(\alpha, \lambda)$ . By independence,

$$M_S(t) = \prod_{i=1}^n \left( \frac{\lambda}{\lambda - t} \right)^\alpha = \left( \frac{\lambda}{\lambda - t} \right)^{n\alpha} \Rightarrow S \sim \text{Gamma}(n\alpha, \lambda).$$

$$M_{\bar{X}}(t) = M_{\frac{1}{n}S}(t) = M_S\left(\frac{1}{n}t\right) = \left( \frac{\lambda}{\lambda - t/n} \right)^{n\alpha} = \left( \frac{n\lambda}{n\lambda - t} \right)^{n\alpha} \Rightarrow \bar{X} \sim \text{Gamma}(n\alpha, n\lambda).$$

4.1 Let  $\bar{X} = \frac{1}{n} \sum X_i$  be the first sample moment about the origin (a.k.a. the sample mean). The population mean for  $\text{Binom}(1, \pi)$  is  $1 \cdot \pi = \pi$ . Our estimate is  $\hat{\pi} = \bar{X}$ .

4.4 For  $X \sim \text{NBinom}$ ,  $E(X) = \frac{\Delta}{\pi} - \Delta$ . So, we set

$$\frac{\Delta}{\hat{\pi}} - \Delta = \bar{X} \Rightarrow \hat{\pi} = \frac{\Delta}{\Delta + \bar{X}}.$$

4.7 favstats reveals  $\bar{X} \doteq 0.6091$ ,  $s \doteq 0.248$ ,  $n = 134 \Rightarrow v = 0.06105$ .

From the formulas:

$$\hat{\alpha} = \bar{X} \left( \frac{\bar{X}(1-\bar{X})}{v} - 1 \right) \doteq 1.7665$$

$$\hat{\beta} = (1-\bar{X}) \left( \frac{\bar{X}(1-\bar{X})}{v} - 1 \right) \doteq 1.1337$$

The beta distribution using these shape parameters gives a very poor fit to the data. By filtering out the players with  $\text{FTPct} = 0$ , the new parameter estimates from remaining players are

$$\hat{\alpha} = 4.824, \quad \hat{\beta} = 2.387,$$

and the fit is vastly improved.

4.9 Take  $\bar{x} = \frac{1}{n} \sum x_i$  and  $v = \frac{1}{n} \sum (x_i - \bar{x})^2$ .

For an exponential distribution,  $\hat{\lambda} = \frac{1}{\bar{x}} \doteq 0.00763$

For a gamma distribution,

$$\frac{\hat{\alpha}}{\hat{\lambda}} = \bar{x} \Rightarrow \hat{\alpha} = \hat{\lambda} \bar{x}$$

$$\frac{\hat{\alpha}}{\hat{\lambda}^2} = v \Rightarrow \frac{\hat{\lambda} \bar{x}}{\hat{\lambda}^2} = \frac{\bar{x}}{\hat{\lambda}} = v$$

$$\Rightarrow \hat{\lambda} = \frac{\bar{x}}{v} \doteq 1.858 \quad \text{and} \quad \hat{\alpha} = \hat{\lambda} \bar{x} = \frac{\bar{x}^2}{v} \doteq 0.0141.$$

4.14 We know  $SE_{\bar{x}} = \frac{\sigma}{\sqrt{n}} = \frac{10}{\sqrt{25}} = 2$ .

Thus,  $Pr(|\bar{X} - \mu| < 3) = \text{pnorm}(3, 0, 2) - \text{pnorm}(-3, 0, 2) \doteq 0.866$ .

4.16 (a) The 10 different SRS, along w/ resulting sample means:

| sample | $\bar{x}$ | sample | $\bar{x}$ |
|--------|-----------|--------|-----------|
| 1, 6   | 3.5       | 6, 8   | 7         |
| 1, 6   | 3.5       | 6, 9   | 7.5       |
| 1, 8   | 4.5       | 6, 8   | 7         |
| 1, 9   | 5         | 6, 9   | 7.5       |
| 6, 6   | 6         | 8, 9   | 8.5       |

$$\Rightarrow \mu_{\bar{x}} = (3.5 + 7 + 7.5)(2/10) + (4.5 + 5 + 6 + 8.5)(1/10) = 6.$$

$$\begin{aligned} \text{Var}(\bar{x}) &= (3.5^2 + 7^2 + 7.5^2)(2/10) + (4.5^2 + 5^2 + 6^2 + 8.5^2)(1/10) - \mu_{\bar{x}}^2 \\ &= 2.85. \end{aligned}$$

(b) From Coro. 4.3.3, we have

$$E(\bar{X}) = \mu = (1+6+6+8+9)(1/5) = 6,$$

a match with part (a). Furthermore,

$$\text{Var}(\bar{X}) = \frac{\sigma^2}{n} \cdot \frac{N-n}{N-1} = \frac{\sigma^2}{2} \cdot \frac{5-2}{5-1} = \frac{3}{8} \sigma^2,$$

where  $\sigma^2 = (1^2 + 6^2 + 6^2 + 8^2 + 9^2)(1/5) - 6^2 = 7.6$ . So,

$$\text{Var}(\bar{X}) = 3/8 \sigma^2 = 2.85, \quad \text{also matching part (a).}$$

(c) We may treat an iid sample as if we were rolling 5-sided dice, yielding pairings:

(1,1) (1,6) (1,6) (1,8) (1,9)  
 (6,1) (6,6) (6,6) (6,8) (6,9)  
 (6,1) (6,6) (6,6) (6,8) (6,9)  
 (8,1) (8,6) (8,6) (8,8) (8,9)  
 (9,1) (9,6) (9,6) (9,8) (9,9)

The 5-by-5 table of means corresponds directly to these pairings

|     |     |     |     |     |
|-----|-----|-----|-----|-----|
| 1   | 3.5 | 3.5 | 4.5 | 5   |
| 3.5 | 6   | 6   | 7   | 7.5 |
| 3.5 | 6   | 6   | 7   | 7.5 |
| 4.5 | 7   | 7   | 8   | 8.5 |
| 5   | 7.5 | 7.5 | 8.5 | 9   |

So,

$$\mu = (1+8+9)(1/25) + (4.5+5+8.5)(2/25) + (3.5+6+7+7.5)(4/25) = 6,$$

and

$$\sigma^2 = (1^2 + 8^2 + 9^2)(1/25) + (4.5^2 + 5^2 + 8.5^2)(2/25) + (3.5^2 + 6^2 + 7^2 + 7.5^2)(4/25) - 6^2 = 3.8.$$

$$4.39 \text{ (a)} \quad |\vec{u}_i|^2 = \left(\frac{1}{\sqrt{n}}\right)^2 + \dots + \left(\frac{1}{\sqrt{n}}\right)^2 = n \cdot \left(\frac{1}{n}\right) = 1 \Rightarrow |\vec{u}_i| = 1.$$

For  $i = 2, 3, \dots, n$ ,

$$\begin{aligned} |\vec{u}_i|^2 &= \frac{1}{i(i-1)} \left[ (i-1)^2 + \sum_{j=1}^{i-1} 1^2 \right] = \frac{1}{i(i-1)} [(i-1)^2 + (i-1)] \\ &= \frac{i-1}{i(i-1)} [(i-1) + 1] = \frac{1}{i} (i) = 1 \Rightarrow |\vec{u}_i| = 1. \end{aligned}$$

Moreover, for  $1 < i < j$ ,

$$\vec{u}_i \cdot \vec{u}_j = 1 - i + \sum_{m=1}^{i-1} 1 = (1-i) + (i-1) = 0.$$

It is more transparent that each  $\vec{u}_i \cdot \vec{u}_i = 0$ ,  $i > 1$ .

(b) Let  $\vec{x} = \langle x_1, x_2, \dots, x_n \rangle$ . Then,

$$\vec{x} \cdot \vec{u}_1 = \frac{1}{\sqrt{n}} (x_1 + x_2 + \dots + x_n) = \sqrt{n} \cdot \frac{1}{n} \sum x_i = \bar{x} \sqrt{n}.$$

(c) With  $\vec{x}$  as in part (b),

$$(\vec{x} \cdot \vec{u}_1) \vec{u}_1 = (\bar{x} \sqrt{n}) \cdot \frac{1}{\sqrt{n}} \langle 1, 1, \dots, 1 \rangle = \langle \bar{x}, \bar{x}, \dots, \bar{x} \rangle.$$

(d) For  $\vec{x} = \langle 3, 4, 4, 7, 7 \rangle$ ,  $\bar{x} = \frac{1}{5}(25) = 5$ , and

$$\vec{v} = \langle 3, 4, 4, 7, 7 \rangle - \langle 5, 5, 5, 5, 5 \rangle = \langle -2, -1, -1, 2, 2 \rangle.$$

Thus,

| $i$ | $\vec{u}_i$  | $\vec{x} \cdot \vec{u}_i$ | $\vec{v} \cdot \vec{u}_i$ | $ \text{proj}(\vec{x} \rightarrow \vec{u}_i) $ | $ \text{proj}(\vec{v} \rightarrow \vec{u}_i) $ |
|-----|--|---------------------------|---------------------------|--|--|
| 1   | $\frac{1}{\sqrt{5}} \langle 1, 1, 1, 1, 1 \rangle$   | $5^{3/2}$                 | 0                         | $5^{3/2}$                                      | 0  |
| 2   | $\frac{1}{\sqrt{2}} \langle 1, -1, 0, 0, 0 \rangle$  | $-1/\sqrt{2}$             | $-1/\sqrt{2}$             | $1/\sqrt{2}$                                   | $1/\sqrt{2}$                                   |
| 3   | $\frac{1}{\sqrt{6}} \langle 1, 1, -2, 0, 0 \rangle$  | $-1/\sqrt{6}$             | $-1/\sqrt{6}$             | $1/\sqrt{6}$                                   | $1/\sqrt{6}$                                   |
| 4   | $\frac{1}{\sqrt{12}} \langle 1, 1, 1, -3, 0 \rangle$ | $-10/\sqrt{12}$           | $-10/\sqrt{12}$           | $10/\sqrt{12}$                                 | $10/\sqrt{12}$                                 |
| 5   | $\frac{1}{\sqrt{20}} \langle 1, 1, 1, 1, -4 \rangle$ | $-10/\sqrt{20}$           | $-10/\sqrt{20}$           | $10/\sqrt{20}$                                 | $10/\sqrt{20}$                                 |

For  $i \geq 2$ ,  $\vec{x} \cdot \vec{u}_i = \vec{v} \cdot \vec{u}_i$ . This is true for other  $\vec{x} \in \mathbb{R}^5$ , too, since

$$\vec{v} \cdot \vec{u}_i = (\vec{x} - \bar{x} \vec{u}_1) \cdot \vec{u}_i = \vec{x} \cdot \vec{u}_i - \bar{x} \cdot \vec{u}_i = \vec{x} \cdot \vec{u}_i - 0$$

since  $\bar{x} \vec{u}_1$ , being parallel to  $\vec{u}_1$ , is orthogonal to each  $\vec{u}_i$  with  $i \geq 2$ .

4.40 (a)  $\bar{x} = (3+4+5+8)/4 = 5$

$$s^2 = \frac{1}{3} \left[ (3-5)^2 + (4-5)^2 + (5-5)^2 + (8-5)^2 \right] = \frac{14}{3}.$$

(b)  $\vec{p}_1 = \langle 5, 5, 5, 5 \rangle$ , as determined in Exercise 4.39.

$$\vec{p}_2 = \frac{1}{\sqrt{2}} (3-4) \vec{u}_2 = \langle -\frac{1}{2}, \frac{1}{2}, 0, 0 \rangle$$

$$\vec{p}_3 = \frac{1}{\sqrt{6}} (3+4-10) \vec{u}_3 = \langle -\frac{1}{2}, -\frac{1}{2}, 1, 0 \rangle$$

$$\vec{p}_4 = \frac{1}{\sqrt{12}} (3+4+5-24) \vec{u}_4 = \langle -1, -1, -1, 3 \rangle$$

and  $\sum \vec{p}_i = \langle 3, 4, 5, 8 \rangle$  as predicted.

$$(c) \quad l_1 = |\vec{p}_1| = \sqrt{4(5^2)} = 10$$

$$l_2 = |\vec{p}_2| = \sqrt{2\left(\frac{1}{2}\right)^2} = \frac{1}{\sqrt{2}}$$

$$l_3 = |\vec{p}_3| = \sqrt{2\left(\frac{1}{2}\right)^2 + 1^2} = \sqrt{3/2}$$

$$l_4 = |\vec{p}_4| = \sqrt{3(-1)^2 + 3^2} = 2\sqrt{3}$$

$$(d) \quad \sum_{i=2}^4 l_i^2 = \frac{1}{2} + \frac{3}{2} + 12 = 14 = 3\left(\frac{14}{3}\right) = 3s^2.$$

$$C.10 \quad \text{proj}(\langle 1, 2, 3 \rangle \rightarrow \langle 1, 1, 1 \rangle) = \frac{\langle 1, 2, 3 \rangle \cdot \langle 1, 1, 1 \rangle}{|\langle 1, 1, 1 \rangle|^2} \langle 1, 1, 1 \rangle = \langle 2, 2, 2 \rangle$$

This is the closest vector to  $\langle 1, 2, 3 \rangle$  in  $W = \text{span}(\langle 1, 1, 1 \rangle)$ , so

$$\langle 1, 2, 3 \rangle - \text{proj}(\langle 1, 2, 3 \rangle \rightarrow \langle 1, 1, 1 \rangle) = \langle -1, 0, 1 \rangle$$

is perpendicular/orthogonal to  $\langle 1, 1, 1 \rangle$ , and

$$\text{span}\{\langle 1, 1, 1 \rangle, \langle 1, 2, 3 \rangle\} = \text{span}\{\langle 1, 1, 1 \rangle, \langle -1, 0, 1 \rangle\}.$$

To get these orthogonal vectors to be of length 1, we divide by their lengths:

$$\vec{u}_1 = \frac{1}{\sqrt{3}} \langle 1, 1, 1 \rangle = \left\langle \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right\rangle, \quad \vec{u}_2 = \frac{1}{\sqrt{2}} \langle -1, 0, 1 \rangle = \left\langle -\frac{1}{\sqrt{2}}, 0, \frac{1}{\sqrt{2}} \right\rangle.$$