

Contents

| | |
|---|-----------|
| Introduction | 1 |
| 1 The Straight Line | 2 |
| 1.1 Straight Line Equations | 2 |
| 1.2 Parallel and Perpendicular Lines | 4 |
| 1.3 The Angle a Line Makes with the x -Axis | 6 |
| 1.4 Collinearity | 8 |
| 1.5 The Midpoint | 10 |
| 1.6 Altitudes of a Triangle | 12 |
| 1.7 Medians of a Triangle | 14 |
| 1.8 Perpendicular Bisectors | 16 |
| Review Exercise | 18 |
| 2 Recurrence Relations | 19 |
| 2.1 Finding Terms in a Sequence | 19 |
| 2.2 Determining a Recurrence Relation | 21 |
| 2.3 The Limit of a Recurrence Relation | 23 |
| Review Exercise | 25 |
| 3 Differentiation I | 26 |
| 3.1 The Derivative of ax^n | 26 |
| 3.2 Differentiable Form | 28 |
| 3.3 Leibniz Notation | 30 |
| 3.4 The Gradient of a Tangent to a Curve | 32 |

| | | |
|----------|---|-----------|
| 3.5 | The Equation of a Tangent to a Curve | 34 |
| 3.6 | The Rate of Change of a Function | 36 |
| | Review Exercise | 38 |
| 4 | Quadratic Theory | 39 |
| 4.1 | Solving Quadratic Inequations | 39 |
| 4.2 | Completing the Square | 41 |
| 4.3 | The Discriminant | 43 |
| 4.4 | The Intersection of a Line and a Curve | 45 |
| 4.5 | Tangency | 47 |
| | Review Exercise | 49 |
| 5 | Sets and Functions | 50 |
| 5.1 | The Domain of a Function | 50 |
| 5.2 | The Range of a Function | 52 |
| 5.3 | Composite Functions | 54 |
| 5.4 | Inverse Functions | 56 |
| | Review Exercise | 58 |
| 6 | Trigonometry | 59 |
| 6.1 | Exact Values | 59 |
| 6.2 | Solving Trig Equations using Exact Values | 61 |
| 6.3 | Radians | 63 |
| 6.4 | Solving Trig Equations and Radians | 65 |
| 6.5 | Solving Phase Angle Equations | 67 |
| 6.6 | Solving Multiple Angle Equations | 69 |
| 6.7 | Review Exercise | 71 |
| 7 | Graph Transformations | 72 |
| 7.1 | Translations | 72 |
| 7.2 | Stretches and Compressions | 74 |

| | | |
|-----------|---|------------|
| 7.3 | Reflections | 76 |
| 7.4 | Composite Transformations | 78 |
| | Review Exercise | 80 |
| 8 | Vectors | 81 |
| 8.1 | Parallel Vectors | 81 |
| 8.2 | Collinearity using Vectors | 83 |
| 8.3 | The Division of a Line in a Ratio | 85 |
| 8.4 | Unit Vectors i , j and k | 87 |
| 8.5 | The Scalar Product | 89 |
| 8.6 | The Angle Between Two Vectors | 91 |
| 8.7 | Properties of the Scalar Product | 93 |
| 8.8 | Vector Pathways | 95 |
| | Review Exercise | 97 |
| 9 | Differentiation II | 98 |
| 9.1 | Increasing and Decreasing Functions | 98 |
| 9.2 | Stationary Points | 102 |
| 9.3 | Closed Intervals | 106 |
| 9.4 | Optimisation | 108 |
| 9.5 | The Graph of the Derivative | 111 |
| | Review Exercise | 114 |
| 10 | Polynomials | 115 |
| 10.1 | Polynomials and Synthetic Division | 115 |
| 10.2 | Remainder Theorem | 117 |
| 10.3 | Finding Unknown Coefficients | 123 |
| 10.4 | Solving Polynomial Equations | 125 |
| 10.5 | Identifying Polynomial Functions | 126 |
| | Review Exercise | 128 |

| | |
|---|------------|
| 11 Integration | 129 |
| 11.1 Indefinite Integrals | 130 |
| 11.2 Differential Equations | 132 |
| 11.3 Definite Integrals | 133 |
| 11.4 The Area Under a Curve | 134 |
| 11.5 The Area Between Two Curves | 138 |
| Review Exercise | 140 |
| 12 Compound Angles | 141 |
| 12.1 The Addition Formulae | 141 |
| 12.2 Applications to Right-Angled Triangles | 143 |
| 12.3 The Double Angle Formulae | 145 |
| Review Exercise | 145 |
| 13 The Circle | 146 |
| 13.1 The Equation of the Circle | 147 |
| 13.2 The General Equation of the Circle | 149 |
| 13.3 The Intersection of a Line and a Circle | 151 |
| 13.4 Finding the Equation of a Tangent to a Circle | 153 |
| 13.5 Points Inside and Outside Circles | 155 |
| 13.6 The Intersections of Circles | 157 |
| Review Exercise | 159 |
| 14 The Wave Function | 160 |
| 14.1 The Wave Function using $k \cos(x - a)$ | 161 |
| 14.2 Other Forms of the The Wave Function | 163 |
| 14.3 Maximum and Minimum Values using the Wave Function | 164 |
| 14.4 Solving Equations using the Wave Function | 165 |
| 14.5 Multiple Angles and Different Variables | 166 |
| Wave Function Review Exercise | 167 |

| | |
|--|------------|
| 15 Logs and Exponentials | 168 |
| 15.1 Graphs of Exponential Functions | 169 |
| 15.2 Evaluating Logarithms | 171 |
| 15.3 Laws of Logs | 172 |
| 15.4 Solving Log Equations | 173 |
| 15.5 Graphs of Logarithmic Functions | 175 |
| 15.6 Solving Exponential Equations | 177 |
| 15.7 Natural Logarithms | 178 |
| 15.8 Exponential Growth and Decay | 179 |
| 15.9 Experimental Data of the form $y = ax^b$ | 181 |
| 15.10 Experimental Data of the form $y = ab^x$ | 183 |
| Logs and Exponentials Review Exercise | 185 |
| 16 Further Calculus | 186 |
| 16.1 Differentiating Trig Functions | 186 |
| 16.2 Integrating Trig Functions | 188 |
| 16.3 The Chain Rule for Differentiation | 190 |
| 16.4 Integrating $(ax + b)^n$ | 192 |
| Review Exercise | 194 |
| Challenge Problems | 195 |
| Answers | 213 |

Introduction

Intro

1.1 Straight Line Equations

Exercise

1.2 Parallel and Perpendicular Lines

Exercise

1.3 The Angle a Line Makes with the x -Axis

Exercise

1.4 Collinearity

Exercise

1.5 The Midpoint

Exercise

1.6 Altitudes of a Triangle

Exercise

1.7 Medians of a Triangle

Exercise

1.8 Perpendicular Bisectors

Exercise

Review Exercise

2.1 Finding Terms in a Sequence

Exercise

2.2 Determining a Recurrence Relation

Exercise

2.3 The Limit of a Recurrence Relation

Exercise

Review Exercise

3.1 The Derivative of ax^n

Exercise

3.2 Differentiable Form

Exercise

3.3 Leibniz Notation

Exercise

3.4 The Gradient of a Tangent to a Curve

Exercise

3.5 The Equation of a Tangent to a Curve

Exercise

3.6 The Rate of Change of a Function

Exercise

Review Exercise

4.1 Solving Quadratic Inequations

Exercise

4.2 Completing the Square

Exercise

4.3 The Discriminant

Exercise

4.4 The Intersection of a Line and a Curve

Exercise

4.5 Tangency

Exercise

Review Exercise

5.1 The Domain of a Function

Exercise

5.2 The Range of a Function

Exercise

5.3 Composite Functions

Exercise

5.4 Inverse Functions

Exercise

Review Exercise

6.1 Exact Values

Exercise

6.2 Solving Trig Equations using Exact Values

Exercise

6.3 Radians

Exercise

6.4 Solving Trig Equations and Radians

Exercise

6.5 Solving Phase Angle Equations

Exercise

6.6 Solving Multiple Angle Equations

Exercise

6.7 Review Exercise

7.1 Translations

Exercise

7.2 Stretches and Compressions

Exercise

7.3 Reflections

Exercise

7.4 Composite Transformations

Exercise

Review Exercise

8.1 Parallel Vectors

Exercise

8.2 Collinearity using Vectors

Exercise

8.3 The Division of a Line in a Ratio

Exercise

8.4 Unit Vectors i , j and k

Exercise

8.5 The Scalar Product

Exercise

8.6 The Angle Between Two Vectors

Exercise

8.7 Properties of the Scalar Product

Exercise


8.8 Vector Pathways

Exercise

Review Exercise

9.1 Increasing and Decreasing Functions

A function can be said to be *increasing* for values of x for which a tangent to the curve has a *positive gradient*, and *decreasing* where a tangent to the curve has a *negative gradient*.



A function $f(x)$ is:


- Increasing when $f'(x) > 0$.
- Decreasing when $f'(x) < 0$.

Function f above is *increasing* for $x < a$, $x > b$ and *decreasing* for $a < x < b$.

Example 9.1.1

Part of the graph of a function $f(x)$ is below. State the values of x for which $f(x)$ is:

- (a) Increasing.
- (b) Decreasing.



- (a) The function $f(x)$ is increasing for $x < -2$ and $x > 6$.
- (b) The function $f(x)$ is decreasing for $-2 < x < 6$.

Evaluating $f'(x)$ can show whether a function is increasing or decreasing at that point:

Example 9.1.2

Determine whether the function $f(x) = 5x^2 - 6\sqrt{x}^3$ is increasing or decreasing when $x = 4$.

$$f(x) = 5x^2 - 6x^{\frac{3}{2}} \quad \leftarrow \text{prepare to differentiate}$$

$$f'(x) = 10x - 9x^{\frac{1}{2}} \quad \leftarrow \text{differentiate}$$

$$= 10x - 9\sqrt{x} \quad \leftarrow \text{write in radical form}$$

$$f'(4) = 10(4) - 9\sqrt{4} \quad \leftarrow \text{substitute}$$

$$= 22 \quad \leftarrow \text{evaluate}$$

Since $f'(4) > 0$, the function f is increasing when $x = 4$.

The range of values of x for which a function is increasing or decreasing can also be determined:

Example 9.1.3

Determine the range of values of x for which the function $f(x) = 2x^2 - 8x + 11$ is increasing.

$$\text{Increasing} \implies f'(x) > 0$$

$$f'(x) = 4x - 8$$

← differentiate

$$4x - 8 > 0$$

← apply condition

$$4x > 8$$

$$x > 2$$

← solve

So $f'(x)$ is increasing when $x > 2$

Quadratic inequations may arise when finding ranges, which require a sketch when solving.

Example 9.1.4

Determine the range of values of x for which $y = \frac{2}{3}x^3 + 3x^2 - 8x + 7$ is decreasing.

$$\text{Decreasing} \implies \frac{dy}{dx} < 0$$

$$\frac{dy}{dx} = 2x^2 + 6x - 8$$

$$2x^2 + 6x - 8 < 0$$

$$x^2 + 3x - 4 < 0$$

quadratic inequation

Consider roots:

$$x^2 + 3x - 4 = 0$$

$$(x + 4)(x - 1) = 0$$

$$x = -4, x = 1$$



So $f'(x)$ is decreasing when $-4 < x < 1$

A function can be said to be **strictly increasing** if $f'(x) > 0$ across its domain.

A function can be said to be **strictly decreasing** if $f'(x) < 0$ across its domain.

Example 9.1.5

Show that $f(x) = \frac{1}{3}x^3 - 4x^2 + 21x - 10$ is strictly increasing for all values of x .

Note that $x^2 - 8x + 1$ can be expressed as $(x - 4)^2 + 5$.

$$\begin{aligned}f'(x) &= x^2 - 8x + 21 \\ &= (x - 4)^2 + 5\end{aligned}$$

← differentiate

← complete the square

∴ minimum value of $f'(x)$ is 5

← state minimum value of f'

Since $5 > 0$, $f'(x) > 0$ for all x ,


← compare to zero

so the function f is strictly increasing.

← conclusion about f

9.2 Stationary Points

A *stationary point* can be found where a tangent to a curve has a *gradient of zero*, or:



Stationary points occur where $\frac{dy}{dx} = 0$

The graph of $y = f(x)$ has stationary points at $(-2, 3)$ and $(6, -2)$.

Example 9.2.1

Find the coordinates of the stationary points on the curve $y = x^3 - \frac{9}{2}x^2 + 6x$.

Find x -coordinate(s)

Stationary points occur where $\frac{dy}{dx} = 0$

$$\frac{dy}{dx} = 3x^2 - 9x + 6$$

$$3x^2 - 9x + 6 = 0$$

$$x^2 - 3x + 2 = 0$$

$$(x - 1)(x - 2) = 0$$

$$x = 1, x = 2$$

Find y -coordinates


$$\begin{aligned} \text{When } x = 1, \quad y &= (1)^3 - \frac{9}{2}(1)^2 + 6(1) \\ &= \frac{5}{2} \end{aligned}$$

$$\begin{aligned} \text{When } x = 2, \quad y &= (2)^3 - \frac{9}{2}(2)^2 + 6(2) \\ &= 2 \end{aligned}$$


\therefore stationary points occur at $(1, \frac{5}{2})$ and $(1, 2)$

Either Leibniz notation $\left(\frac{dy}{dx}\right)$ or function notation $(f'(x))$ may be used depending on the context.


Determining the nature of a stationary point means to find out *what type* it is. There are four:




minimum
turning point



maximum
turning point



rising point
of inflection



falling point
of inflection

The **second derivative**, $f''(x)$ or $\frac{d^2y}{dx^2}$, is obtained by **differentiating the first derivative**.

For example:

$$f(x) = 4x^3 - 3x^2 + 7x - 1$$

$$f'(x) = 12x^2 - 6x + 7$$

$$f''(x) = 24x - 6$$

- Stationary point (a, b) is a:
- **maximum turning point** if $f''(a) < 0$
 - **minimum turning point** if $f''(a) > 0$

Example 9.2.2

Determine the coordinates and nature of the stationary points of $f(x) = x^3 - 12x^2 + 36x$.

Find x -coordinate(s)

SPs occur where $f'(x) = 0$

$$f'(x) = 3x^2 - 24x + 36$$

$$3x^2 - 24x + 36 = 0$$

$$x^2 - 8x + 12 = 0$$

$$(x - 2)(x - 6) = 0$$

$$x = 2, x = 6$$

Find y -coordinates

$$f(2) = (2)(2 - 6)^2 = 32$$

$$f(6) = (6)(6 - 6)^2 = 0$$

Determine Nature

$$f''(x) = 6x - 24$$

$$f''(2) = 6(2) - 24 = -12$$




$$f''(2) < 0 \therefore \text{max TP}$$

$$f''(6) = 6(6) - 24 = 12$$




$$f''(6) > 0 \therefore \text{min TP}$$

\therefore maximum turning point at $(2, 32)$ and minimum turning point at $(6, 0)$




If the second derivative is 0, a **nature table** can be used instead, where $\frac{dy}{dx}$ is evaluated (> 0 or < 0) for values of x either side of a stationary point, to visualise its nature. For stationary point (a, b) :

| | | | |
|-----------------|---|---|---|
| x | \longrightarrow | a | \longrightarrow |
| $\frac{dy}{dx}$ | $-ve$ | 0 | $+ve$ |
| slope |  |  |  |




minimum
turning point

| | | | |
|-----------------|---|---|---|
| x | \longrightarrow | a | \longrightarrow |
| $\frac{dy}{dx}$ | $+ve$ | 0 | $-ve$ |
| slope |  |  |  |

maximum
turning point

| | | | |
|-----------------|---|--|---|
| x | \longrightarrow | a | \longrightarrow |
| $\frac{dy}{dx}$ | $+ve$ | 0 | $+ve$ |
| slope |  |  |  |

rising point
of inflection

| | | | |
|-----------------|---|---|---|
| x | \longrightarrow | a | \longrightarrow |
| $\frac{dy}{dx}$ | $-ve$ | 0 | $-ve$ |
| slope |  |  |  |

falling point
of inflection

Example 9.2.3

Find the coordinates and nature of the stationary point on the curve $y = 13 + 6x^2 - 12x - x^3$.

Find x -coordinate(s)

SPs occur where $\frac{dy}{dx} = 0$

$$\frac{dy}{dx} = 12x - 12 - 3x^2$$

$$12x - 12x - 3x^2 = 0$$

$$4x - 4 - x^2 = 0$$

$$x^2 - 4x + 4 = 0$$




$$(x - 2)(x - 2) = 0$$

$$x = 2$$

Find y -coordinate

When $x = 2$,

$$\begin{aligned} y &= 13 + 6(2)^2 - 12(2) - (2)^3 \\ &= 13 + 24 - 24 - 8 \\ &= 5 \end{aligned}$$

| | | | |
|-----------------|---|---|---|
| x | $\xrightarrow{1}$ | 2 | $\xrightarrow{3}$ |
| $\frac{dy}{dx}$ | $-ve$ | 0 | $-ve$ |
| slope |  |  |  |

Determine Nature

$$\frac{d^2y}{dx^2} = -12 - 6x$$

When $x = 2$,


$$\begin{aligned} \frac{d^2y}{dx^2} &= -12 - 6(2) \\ &= 0 \end{aligned}$$

\therefore nature table

\therefore falling point of inflection at $(2, 5)$

9.3 Closed Intervals

Given a smooth, continuous function across a *closed interval*, finding the coordinates and nature of stationary points can allow the maximum and minimum values to be determined.




Maximum and minimum values can occur at:

- Turning points
- *Bounds* of the interval

For $f(x)$, the maximum value occurs when $x = b$ and the minimum at the minimum turning point.

Example 9.3.1

Function $f(x) = \frac{2}{3}x^3 - x^2 - 4x - \frac{1}{3}$ has two stationary points, and part of the graph of $y = f(x)$ is shown below. Find the maximum and minimum values of f on the interval $-3 \leq x \leq 3$.




Check y -coordinates at the lower and upper bounds:

$$\text{When } x = -3, \quad y = \frac{2}{3}(-3)^3 - (-3)^2 - 4(-3) - \frac{1}{3} = -\frac{46}{3}$$

$$\text{When } x = 3, \quad y = \frac{2}{3}(3)^3 - (3)^2 - 4(3) - \frac{1}{3} = -\frac{10}{3}$$

\therefore maximum value of f is 2, which occurs when $x = -1$

minimum value of f is $-\frac{46}{3}$, which occurs when $x = -3$



Where a function is *strictly increasing* or *strictly decreasing* in a closed interval, its maximum and minimum values can *only* be located at the bounds of the interval.

Example 9.3.2


Given $y = x^3 + \frac{1}{2}x^2 - \frac{7}{2}$ is strictly increasing, find the greatest and least values of y in the interval $-1 \leq x \leq 3$.

$$\text{When } x = -1, \quad y = (-2)^3 + \frac{1}{2}(-1)^2 - \frac{7}{2} = -4$$

$$\text{When } x = 3, \quad y = (3)^3 + \frac{1}{2}(3)^2 - \frac{7}{2} = 28$$

\therefore the maximum value of y is 28

and the minimum value of y is -4



9.4 Maximum and Minimum Values

If a smooth, continuous function has only one stationary point within its domain, and it is a *maximum turning point*, the *greatest value* of the function will be located at the turning point. The corresponding will be true for the *least* value of a function with only a *minimum* turning point. \[0.1em]

Example 9.4.1

Determine the value of x for which the function f , defined by $f(x) = x + \frac{1}{x}$, $x > 0$, has its minimum value.

Find x -coordinate(s)

Stationary points occur where $f'(x) = 0$

$$\begin{aligned} f(x) &= x + x^{-1} \\ f'(x) &= 1 - x^{-2} \\ 1 - x^{-2} &= 0 \\ 1 - \frac{1}{x^2} &= 0 \\ x^2 - 1 &= 0 \\ x^2 &= 1 \\ x &= 1 \text{ (Since } x > 0) \end{aligned}$$

Determine Nature

$$\begin{aligned} f''(x) &= 2x^{-3} \\ &= \frac{2}{x^3} \\ f''(1) &= \frac{2}{1^3} \\ &= 2 \end{aligned}$$

Since $f''(1) > 0$, min TP when $x = 1$

$\therefore f$ is minimised when $x = 1$

When using variables other than x and y (and function f), care should be taken with notation.

Example 9.4.2

Determine the maximum value of P given $P = 6t - t^3$, where $t > 0$, and the value of t for which it occurs.

Find x -coordinate(s)

Stationary points occur where $\frac{dP}{dt} = 0$

$$\begin{aligned} \frac{dP}{dt} &= 6 - 3t^2 \\ 6 - 3t^2 &= 0 \\ 6 &= 3t^2 \\ 2 &= t^2 \\ \sqrt{2} &= t \end{aligned}$$

Determine Nature

$$\begin{aligned} \frac{d^2P}{dt^2} &= -6t \\ \text{When } t &= \sqrt{2}, \quad \frac{d^2P}{dt^2} = -6\sqrt{2} \\ \frac{d^2P}{dt^2} &< 0 \therefore \text{max TP when } t = \sqrt{2} \end{aligned}$$

Maximum Value

$$\begin{aligned} \text{When } t &= \sqrt{2}, P = 6(\sqrt{2}) - \sqrt{2}^3 = 4\sqrt{2} \\ \therefore \text{max value of } P &\text{ is } 4\sqrt{2}, \text{ when } t = \sqrt{2} \end{aligned}$$


If a function has a single stationary point within an interval, which is a maximum turning point, and the *minimum* value of the function is required, then it must be located at one of the bounds, and the value of the function at each must be checked.

9.4 Optimisation

When a mathematical equation is used to model some real-world variables, finding maximum or minimum values on an interval is often desired. This is referred to as *optimisation*.

Example 9.5.1

A company has determined that a mathematical equation can model the amount of revenue in thousands of pounds, R , which may be earned by setting the price for a new product as x pounds.



The diagram illustrates the model used by the company, which is:

$$R(x) = 8x^3 - x^4 \text{ for } 0 < x \leq 8$$

Find the value of x which gives the maximum revenue for the product.

Stationary Points occur where $R'(x) = 0$

$$R'(x) = 24x^2 - 4x^3$$

$$24x^2 - 4x^3 = 0$$

$$6x^2 - x^3 = 0$$

$$x^2(6 - x) = 0$$

$$x = 6 \text{ since } x > 0$$

Determine Nature

$$R''(x) = 48x - 12x^2$$


$$R''(6) = 48(6) - 12(6)^2$$

$$= -144$$

$$R''(6) < 0 \therefore \text{maximum revenue when } x = 6$$

Example 9.5.2

A chicken is aiming to cross a road. It can fly faster than it can walk, but doing so tires it.



If it flies for x metres and walks the rest of the way then the amount of time T it will take to reach its destination is given, in seconds, by:

$$T(x) = x^2 + \frac{16}{x}$$

Determine the value of x which minimises the chicken's time to get to the other side.

Stationary Points occur where $T'(x) = 0$

$$T(x) = x^2 + 16x^{-1}$$

$$T'(x) = 2x - 16x^{-2}$$

$$2x - 16x^{-2} = 0$$

$$2x - \frac{16}{x^2} = 0$$

$$2x^3 - 16 = 0$$

$$x^3 - 8 = 0$$

$$x^3 = 8$$

$$x = 2$$

Determine Nature

$$T''(x) = 2 + 32x^{-3}$$

$$= 2 + \frac{32}{x^3}$$

$$T''(2) = 2 + \frac{32}{2^3}$$

$$= 2 + 4$$

$$= 6$$


$$T''(2) > 0$$

$\therefore T$ is minimised when $x = 2$

It can be asked for the equation used in the mathematical model to be *shown* to be correct. Establishing the equation will require an amount of problem-solving. A common context involves the volume and surface area of a 3D object.

Example 9.5.3

A solid cuboid measures x cm by x cm by h cm. The volume of this cuboid is 125 cm^3 .



- (a) Show that the surface area of the cuboid can be given by:

$$A(x) = 2x^2 + \frac{500}{x}$$

- (b) Find the value of x such that the surface area is minimised, and find the minimum surface area.

It is recommended to first form an equation for the surface area, A , in terms of x and h . By then finding the relationship between x and h , substitution allows h to be eliminated.

(a)

Surface Area

$$\begin{aligned} A &= 2 \times x^2 + 4 \times xh \\ &= 2x^2 + 4xh \\ &= 2x^2 + 4x \left(\frac{125}{x^2} \right) \\ &= 2x^2 + \frac{500}{x} \text{ as required} \end{aligned}$$

Volume

$$\begin{aligned} V &= L \times B \times H \\ 125 &= x^2 h \\ \frac{125}{x^2} &= h \end{aligned}$$

It should be noted that part (b) can be answered in full even if part (a) is not completed.

(b) **Stationary Points** occur where $A'(x) = 0$

$$\begin{aligned} A(x) &= 2x^2 + 500x^{-1} \\ A'(x) &= 4x - 500x^{-2} \\ 4x - 500x^{-2} &= 0 \\ 4x - \frac{500}{x^2} &= 0 \\ 4x^3 - 500 &= 0 \\ x^3 - 125 &= 0 \\ x^3 &= 125 \\ x &= 5 \end{aligned}$$

Determine Nature


$$\begin{aligned} A''(x) &= 4 + 1000x^{-3} \\ &= 4 + \frac{1000}{x^3} \\ A''(5) &= 4 + \frac{1000}{5^3} \\ &= 4 + 8 \\ &= 12 \\ A''(5) &> 0 \\ \therefore A &\text{ is minimised when } x = 5 \end{aligned}$$

Minimum Area


$$\begin{aligned} A(5) &= 2(5)^2 + \frac{500}{5^2} \\ &= 70 \text{ cm}^2 \end{aligned}$$

9.5 The Graph of the Derivative


Part of the graph of $f(x)$ for function f is shown below.




The graph of the *derived function*, $f'(x)$, plots the *rate of change* of f against values of x . Where $f(x)$ is *increasing*, $f'(x)$ has a positive value, $f(x)$ is *decreasing*, $f'(x)$ has a negative value



Stationary points on $f(x)$, where $f'(x) = 0$, give corresponding *roots* on the graph of $f'(x)$.




Above the x -axis when $f(x)$ is increasing, and below when $f(x)$ is decreasing, $f'(x)$ can be sketched.




Example 9.6.1

Part of the graph of $y = f(x)$ is shown below. Sketch $y = f'(x)$.




Stationary points on $y = f(x)$ are roots of $y = f'(x)$:




Example 9.6.2

For cubic function $g(x)$, part of the graph of $y = g(x)$ is shown below. Sketch $y = g'(x)$.



Since $g(x)$ is cubic, $g'(x)$ must be quadratic, and so the graph is parabolic:



Review Exercise

10.1 Polynomials and Synthetic Division

Expressions comprised of sums of *positive, integer* powers of x are referred to as *polynomials in x* .


Constant
Degree 0
e.g. 7

Linear
Degree 1
e.g. $2x + 4$

Quadratic
Degree 2
e.g. $x^2 - 3x + 1$

Cubic
Degree 3
e.g. $4x^3 - 2x$

Quartic
Degree 4
e.g. $x^4 - 3x^3 + 1$



A polynomial function such as $f(x) = 2x^3 - 7x^2 + 4x - 1$ can be evaluated for some value of x :

$$\begin{aligned} f(3) &= 2(3)^3 - 7(3)^2 + 4(3) - 1 \\ &= 54 - 63 + 12 - 1 \\ &= 4 \end{aligned}$$

Another way to evaluate $f(3)$ is to use synthetic division, as follows:

- 1) Draw a table as below and fill in the first row, then bring the first coefficient down:

| | | | | | | |
|---------|---|---|----|---|---|--|
| $x = 3$ | 3 | 2 | -7 | 4 | 1 | ← Coefficients of $2x^3 - 7x^2 + 4x - 1$ |
| | | 2 | | | | |

- 2) The 6 is obtained by *multiplying* 2 by 3, and the -1 by *adding* the two values above:

| | | | | | |
|---|---|----|---|---|---|
| ③ | 2 | -7 | 4 | 1 | Multiply diagonally by 3 Add columns |
| | 2 | 6 | | | |

- 3) By continuing this *multiplying then adding* pattern, the value of $f(3) = 4$ can be obtained:

| | | | | | |
|---|---|----|----|---|-------------------------|
| 3 | 2 | -7 | 4 | 1 | |
| | | 6 | -3 | 3 | |
| | 2 | -1 | 1 | 4 | ← $\therefore f(3) = 4$ |

Example 10.1.1

Given $f(x) = x^4 - 4x^3 + 3x^2 + 9$, use synthetic division to evaluate $f(2)$.

| | | | | | | |
|---|----|----|----|----|----|-----------------------|
| 2 | 1 | -4 | 3 | 0 | 9 | |
| | | 2 | -4 | -2 | -4 | |
| 1 | -2 | -1 | -2 | 5 | | $\therefore f(2) = 5$ |

The top line within a synthetic division box should be *ordered in decreasing powers of x* . Any missing powers are included with a coefficient of zero.

10.2 Remainder Theorem

Addition, subtraction and multiplication of polynomials were all required for the National 5 course.

Since $3 \times 4 + 1 = 13$, this implies that:

$$\frac{13}{3} = 4 \text{ Remainder } 1$$

Since $(x - 2)(x^2 - 3x + 4) + 7 = x^3 - 5x^2 + x - 1$, this implies that:

$$\frac{x^3 - 5x^2 + x - 1}{x - 2} = x^2 - 3x + 4 \text{ Remainder } 7$$

Note that for $f(x) = x^3 - 5x^2 + x - 1$, $f(2) = (2)^3 - 5(2)^2 + (2) - 1 = 7$.

Remainder Theorem:

If polynomial $f(x)$ is divided by a linear divisor $(x - a)$ then the remainder is $f(a)$, and $f(x) = (x - a)Q(x) + f(a)$ where quotient $Q(x)$ is a polynomial of degree one less than $f(x)$.

Example 10.2.1

Find the remainder when $3x^3 - x^2 + 5$ is divided by $(x - 1)$.

| | | | | | | |
|--------------------|---|---|----|---|---|-------------------------------|
| $(x - 1)$ | 1 | 3 | -1 | 0 | 5 | |
| $\rightarrow f(1)$ | | | 3 | 2 | 2 | |
| | | 3 | 2 | 2 | 7 | $\leftarrow \text{Remainder}$ |

So the remainder is 7.

The *quotient*, $Q(x)$, can be determined from the values to the left of the remainder. \ Since $3x^3 - x^2 + 5$ is a *cubic*, the quotient $Q(x)$ is *quadratic*: $3x^2 + 2x + 2$.

$$3x^3 - x^2 + 5 = (x - 1)(3x^2 + 2x + 2) + 7$$

Example 10.2.2

Function f is defined by $f(x) = x^3 + 4x^2 - 3x - 7$. Find the remainder r when $f(x)$ is divided by $(x + 1)$, and hence express $f(x)$ in the form $(x + 1)Q(x) + r$ where $Q(x)$ is a quadratic.

$$\begin{array}{r|rrrr}
 -1 & 1 & 4 & -3 & -7 \\
 & & -1 & -3 & 6 \\
 \hline
 \text{Quotient} \longrightarrow & 1 & 3 & -6 & -1 \longleftarrow \text{Remainder}
 \end{array}$$

So the remainder is -1 and $x^3 + 4x^2 - 3x - 7 = (x + 1)(x^2 + 3x - 6) - 1$.

10.3 Factor Theorem

Since 20 can be divided by 5 to give 4 **with no remainder**, this means 5 is a **factor** of 20.

Factor Theorem:

If polynomial $f(x)$ is divided by a linear **factor** $(x - a)$ then the **remainder** $f(a) = 0$, and $f(x) = (x - a)Q(x)$ where quotient $Q(x)$ is a polynomial of degree one less than $f(x)$. Or:

$$f(a) = 0 \iff (x - a) \text{ is a factor of } f(x)$$

Given a linear factor, synthetic division allows a cubic expression $f(x)$ to be expressed as the product of the *linear* factor and a *quadratic* quotient, which may potentially be further factorised:

$$\begin{aligned} f(x) &= (\text{cubic}) \\ &= (\text{linear})(\text{quadratic}) && \leftarrow \text{Using synthetic division} \\ &= (\text{linear})(\text{linear})(\text{linear}) \end{aligned}$$

Example 10.3.1

Show that $(x + 4)$ is a factor of $x^3 - x^2 - 14x + 24$, and hence factorise it fully.

| | | | | | |
|------------|---|----|-----|-----|-------------|
| -4 | 1 | -1 | -14 | 24 | |
| | | -4 | 20 | -24 | |
| Quotient → | 1 | -5 | 6 | 0 | ← Remainder |

Remainder = 0: $(x + 4)$ is a factor. \leftarrow "Show that.."

$$\begin{aligned} &x^3 - x^2 - 14x + 24 \\ &= (x + 4)(x^2 - 5x + 6) \\ &= (x + 4)(x - 2)(x - 3) \end{aligned}$$

For some cubic expressions, a resulting quadratic quotient may not be able to be factorised.

Example 10.3.2

Show that $(x - 2)$ is a factor of $2x^3 - x^2 + x - 14$, and hence factorise it fully.

$$\begin{array}{r|rrrr}
 2 & 2 & -1 & 1 & -14 \\
 & & 4 & 6 & 14 \\
 \hline
 & 2 & 3 & 7 & 0
 \end{array}$$

"Show that $(x - 2)$ is a factor..." \rightarrow Remainder = 0 $\therefore (x - 2)$ is a factor.

$$\begin{aligned}
 & 2x^3 - x^2 + x - 14 \\
 &= (x - 2)(2x^2 + 3x + 7)
 \end{aligned}$$

Since $3^2 - 4 \times 2 \times 7 = -65$, and $-65 < 0$, the quadratic quotient cannot be factorised
Therefore the cubic cannot be further factorised.

Where a linear factor of a polynomial function is unknown, a suggested approach is to consider the factors of the constant term with the aim of obtaining a value such that the remainder is zero.

Example 10.3.3

Factorise $x^3 - x^2 - 17x = 15$.

Consider $\pm 1, \pm 3 \pm 5 \pm 15$

$f(1) = (1)^3 - (1)^2 - 17(1) - 15 = -32 \therefore (x - 1)$ is **not** a factor.

$f(-1) = (-1)^3 - (-1)^2 - 17(-1) - 15 = 0 \therefore (x + 1)$ **is** a factor.

$$\begin{array}{r|rrrr}
 -1 & 1 & -1 & -17 & -15 \\
 & & -1 & 2 & 15 \\
 \hline
 & 1 & -2 & -15 & 0
 \end{array}$$

$$\begin{aligned}
 & x^3 - x^2 - 17x - 15 \\
 &= (x + 1)(x^2 - 2x - 15) \\
 &= (x + 1)(x + 3)(x - 5)
 \end{aligned}$$

Repeated use of synthetic division may allow a *quartic* (x^4) to be expressed as a product of up to four linear factors.

$$\begin{aligned}
 f(x) &= (\text{quartic}) \\
 &= (\text{linear})(\text{cubic}) && \longleftarrow \text{Using synthetic division} \\
 &= (\text{linear})(\text{linear})(\text{quadratic}) && \longleftarrow \text{Using synthetic division} \\
 &= (\text{linear})(\text{linear})(\text{linear})(\text{linear})
 \end{aligned}$$


Example 10.3.4

Show that $(x - 2)$ is a factor of $x^4 - 2x^3 - 7x^2 + 8x + 12$ and hence factorise it fully.

| | | | | | |
|---|---|----|----|-----|-----|
| 2 | 1 | -2 | -7 | 8 | 12 |
| | | 2 | 0 | -14 | -12 |
| | 1 | 0 | -7 | -6 | 0 |

Remainder = 0: $(x - 2)$ is a factor.

$$x^4 - 2x^3 - 7x^2 + 8x + 12 = (x - 2)(x^3 - 7x - 6)$$

Try factors of 6 

| | | | | |
|---|---|---|----|----|
| 3 | 1 | 0 | -7 | -6 |
| | | 3 | 9 | 6 |
| | 1 | 3 | 2 | 0 |

$$\begin{aligned}
 x^4 - 2x^3 - 7x^2 + 8x + 12 &= (x - 2)(x^3 - 7x - 6) \\
 &= (x - 2)(x - 3)(x^2 + 3x + 2) \\
 &= (x - 2)(x - 3)(x + 1)(x + 2)
 \end{aligned}$$

10.3 Finding Unknown Coefficients

An unknown coefficient of a polynomial expression may be calculated if a factor of the expression is known, or the remainder given a linear divisor is known. It may be preferred to use synthetic division, or to evaluate the expression directly.

Example 10.4.1

Given that $(x + 2)$ is a factor of $2x^3 + kx^2 - 14x + 8$, determine the value of k .

$$\begin{array}{r|rrrr}
 -2 & 2 & k & -14 & 8 \\
 & & -4 & -2k + 8 & 4k + 12 \\
 \hline
 & 2 & k - 4 & -2k - 6 & 4k + 20 \quad \leftarrow \text{Remainder}
 \end{array}$$

Since $(x + 2)$ is a factor, remainder = 0.

$$\begin{aligned}
 4k + 20 &= 0 \\
 4k &= -20 \\
 k &= -5
 \end{aligned}$$

Where a polynomial has *two* unknown coefficients, *simultaneous equations* may be formed and solved.

Example 10.4.2

Find the values of a and b given that:

- $(x - 2)$ is a factor of $x^3 + ax^2 + bx + 12$.
- $x^3 + ax^2 + bx + 12$ divided by $(x - 1)$ has a remainder of 10.

$$\begin{array}{r|rrrr}
 2 & 1 & a & b & 12 \\
 & & 2 & 2a+4 & 4a+2b+8 \\
 \hline
 & 1 & a+2 & 2a+b+4 & 4a+2b+20
 \end{array}$$

Since $(x - 2)$ is a factor, remainder = 0 $\implies 4a + 2b + 20 = 0$.

$$\begin{array}{r|rrrr}
 1 & 1 & a & b & 12 \\
 & & 1 & a+1 & a+b+1 \\
 \hline
 & 1 & a+1 & a+b+1 & a+b+13
 \end{array}$$

Remainder = 10 $\implies a + b + 13 = 10$.

Solve simultaneously:

$$\begin{array}{lcl}
 4a + 2b = -20 & \left. \begin{array}{l} \\ \\ \end{array} \right\} & 4a + 2b = -20 \\
 a + b = -3 & \times 2 & 2a + 2b = -6 \\
 & & \hline
 & & 2a = -14 \quad (-) \\
 & & a = -7
 \end{array}$$

Sub $a = -7$:

$$\begin{array}{l}
 a + b = -3 \\
 (-7) + b = -3 \\
 b = 4
 \end{array}$$


10.4 Solving Polynomial Equations

Quadratic equations such as $x^2 - 4x - 21 = 0$ are typically solved through factorising, or using the quadratic formula. There is no equivalent *general formula* for solving polynomial equations of degree 3 (*cubic*) or higher. Instead, such polynomial equations are solved through *factorising*.

10.5 Identifying Polynomial Functions


Given the form a polynomial function takes and its graph, showing any x -intercepts and at least one other coordinate, the factorised form of the polynomial may be deduced.

- An x -intercept (or *root*) when $x = a$ implies that a polynomial has a *factor* of $(x - a)$.
- A *stationary point* at an x -intercept is a *repeated root*, and its factor will be (at least) squared.



Root when $x = a$


Factor of $(x - a)$



Repeated root when $x = a$

Factor of $(x - a)^n$

The diagram below shows the graph of a quartic function with equation $y = k(x - a)(x - b)(x - c)^2$.



With roots at $x = 1$ and $x = 3$, and a *repeated root* at $x = -2$, the quartic can be expressed as:

$$y = k(x - 1)(x - 3)(x + 2)^2$$

The value of k can only be found if a *non-root* coordinate is also known. By substitution:

$$y = k(x - 1)(x - 3)(x + 2)^2$$

$$6 = k(0 - 1)(0 - 3)(0 + 2)^2$$

$$6 = k(-1)(-3)(2)^2$$

$$6 = 12k$$

$$\frac{6}{12} = k$$

$$\frac{1}{2} = k$$

$$y = \frac{1}{2}(x - 1)(x - 3)(x + 2)^2$$

Substitute in $(0, 6)$


State equation

By comparing the form given and the final equation, it can be seen that: $k = \frac{1}{2}, a = 1, b = 3, c = -2$

Review Exercise

Introduction and Overview

In mathematics, *integration* can be thought of as the process of calculating some kind of “*sum*”, such as the total area under a curve, by breaking it down into sums of increasingly small parts. This technique was known about and used thousands of years ago in both Ancient Greece and other parts of the world.



In the 17th century both Gottfried Wilhelm Leibniz and Isaac Newton independently discovered the *fundamental theorem of calculus*, showing that *integration* can be performed using *antidifferentiation*, which is the ‘reverse’ of differentiation. Now, *integration* is used to refer to both the idea of calculating such a ‘sum’, and to the process of antidifferentiation.



This chapter will first introduce the use of integration to calculate an *antiderivative*, before moving on to using this concept to calculating area enclosed using curves.

Chapter Contents

- 11.1 Indefinite Integrals
- 11.2 Differential Equations
- 11.3 Definite Integrals
- 11.4 The Area Under a Curve
- 11.5 The Area Between Two Curves

Integration skills covered in this chapter will be extended upon in the later chapter of **Further Calculus**.

11.1 Indefinite Integrals

The *derivative* of the function $f(x) = x^3 - 4x^2 - 7x + 3$ is obtained by *multiplying by the power* and then *reducing the power by one* for each term.

$$\begin{aligned}\text{If } f(x) &= x^3 - 4x^2 + 3 \\ \text{Then } f'(x) &= 3x^2 - 8x\end{aligned}$$

Integration can be considered the *inverse* of differentiation, with the aim of taking the *derivative* $f'(x)$ and determining the *original function* $f(x)$. However, given only the derivative $f'(x) = 3x^2 - 8x$ it would not be possible to know any constant term the original function $f(x)$ contained. Finding the *indefinite integral* of a function requires the inclusion of a **Constant of Integration, C** :

$$\begin{aligned}\text{If } f'(x) &= 3x^2 - 8x \\ \text{Then } f(x) &= x^3 - 4x^2 + C\end{aligned}$$

Indefinite Integrals: "The (indefinite) integral of x^n with respect to x ..."

$$\int (x^n) dx = \frac{x^{n+1}}{n+1} + C$$

In other words, **increase the power by 1** then **divide by the new power**.

The *integral sign*, \int , should always appear accompanied by dx , for a function in x .

Note: $\int (f(x) + g(x)) dx = \int f(x) dx + \int g(x) dx$

Example 11.1.1

Find $\int (6x^2 + 8x) dx$.

$$\begin{aligned}\int (6x^2 + 8x) dx \\ &= \frac{6x^3}{3} + \frac{8x^2}{2} + C && \leftarrow \text{Constant of integration} \\ &= 2x^3 + 4x^2 + C && \leftarrow \text{Simplify}\end{aligned}$$

As with differentiation, *preparation for integration* may be needed.

Example 11.1.2

Find $\int \left(4\sqrt{x} - \frac{3}{x^2}\right) dx$, $x > 0$.

$$\begin{aligned}
 & \int \left(4\sqrt{x} - \frac{3}{x^2}\right) dx \\
 &= \int \left(4x^{\frac{1}{2}} - 3x^{-2}\right) dx && \leftarrow \text{Preparing to integrate} \\
 &= \frac{4x^{\frac{3}{2}}}{\frac{3}{2}} - \frac{3x^{-1}}{-1} + C && \leftarrow \text{Integrate: } \frac{1}{2} + 1 = \frac{1}{2} + \frac{2}{2} = \frac{3}{2} \\
 &= \frac{8}{3}x^{\frac{3}{2}} + 3x^{-1} + C && \leftarrow \text{Simplify: } 4 \div \frac{3}{2} = \frac{4}{1} \times \frac{2}{3} = \frac{8}{3}
 \end{aligned}$$

The *derivative* of a *linear* term is a *constant*, so the integral of a *constant* is *linear*.

Example 11.1.3

Find $\int ((2x - 1)(x + 3)) dx$.

$$\begin{aligned}
 & \int ((2x - 1)(x + 3)) dx \\
 &= \int (2x^2 - 5x - 3) dx && \leftarrow \text{Preparing to integrate} \\
 &= \frac{2x^3}{3} - \frac{5x^2}{2} - 3x + C && \leftarrow \text{Integral of } -3 \text{ is } -3x
 \end{aligned}$$

Integration *with respect to variables other than x* should not use dx , and the notation used should match the variable of the function instead.

Example 11.1.4

Find $\int (3t^2 - 5) dt$.

$$\begin{aligned}
 & \int (3t^2 - 5) dt \\
 &= \frac{3t^3}{3} - 5t + C \\
 &= t^3 - 5t + C
 \end{aligned}$$

11.2 Differential Equations

11.3 Definite Integrals

The indefinite integral of the function $f(x)$ can be notated as $F(x)$:

$$\int f(x) \, dx = F(x)$$

The *definite integral* of $f(x)$ from a to b is the difference between $F(b)$ and $F(a)$:

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

Together, they form a core part of the **Fundamental Theorem of Calculus**:

Fundamental Theorem of Calculus:

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

where

$$F(x) = \int f(x) \, dx$$

Since any *constant of integration* within $F(x)$ will cancel through subtraction ($C - C$), it is *not included* when calculating a *definite integral*.


Example 11.3.1

Find $\int_1^2 (3x^2 + 6x - 2) \, dx$.

$$\begin{aligned} & \int_1^2 (3x^2 + 6x - 2) \, dx \\ &= \left[\frac{3x^3}{3} + \frac{6x^2}{2} - 2x \right]_1^2 && \leftarrow \text{Integrate} \\ &= [x^3 + 3x^2 - 2x]_1^2 && \leftarrow \text{Simplify} \\ &= ((2)^3 + 3(2)^2 - 2(2)) - ((1)^3 + 3(1)^2 - 2(1)) && \leftarrow \text{Substitute} \\ &= 16 - 4 \\ &= 12 && \leftarrow \text{Evaluate} \end{aligned}$$


11.4 The Area Under a Curve

The area shaded below, with each "bar" having width 1 unit, is given by the *sum* $1 + 2 + 3 + 1 + 2$.



An integral, \int , can be seen as "sum", but for a continuous function. Given a function $f(x)$, the area enclosed between a section of the curve $y = f(x)$ and the x -axis can be calculated using a definite integral.

Area Under a Curve:




$$\text{Area} = \int_a^b f(x) \, dx = F(b) - F(a)$$

The values of a and b can be referred to as the *bounds* of the integration.


Example 11.4.1

Part of the graph of $y = 5 + 2x - x^2$ is shown. Calculate the shaded area.



$$\begin{aligned} & \int_{-1}^2 (5 + 2x - x^2) dx && \leftarrow \text{Area} \\ &= \left[5x + x^2 - \frac{x^3}{3} \right]_{-1}^2 && \leftarrow \text{Integrate} \\ &= \left(5(2) + (2)^2 - \frac{(2)^3}{3} \right) - \left(5(-1) + (-1)^2 - \frac{(-1)^3}{3} \right) && \leftarrow \text{Substitute} \\ &= \frac{34}{3} - \left(-\frac{11}{3} \right) && \leftarrow \text{Evaluate} \\ &= 15 \text{ square units} \end{aligned}$$

Where the area between a curve and the x -axis lies *under* the x -axis, the definite integral will be negative.




To avoid "*positive*" and "*negative*" areas cancelling each other out, such sections must be calculated as separate integrals, and their *absolute values* added.

$$\text{Area} = A_1 + A_2$$

Example 11.4.2

Part of the graph of $y = x^3 - 10x^2 + 27x - 18$ is shown. Calculate the shaded area.



$$\int_1^3 (x^3 - 10x^2 + 27x - 18) dx$$

$$= \left[\frac{x^4}{4} - \frac{10x^3}{3} + \frac{27x^2}{2} - 18x \right]_1^3$$

$$= \left(\frac{(3)^4}{4} - \frac{10(3)^3}{3} + \frac{27(3)^2}{2} - 18(3) \right)$$

$$- \left(\frac{(1)^4}{4} - \frac{10(1)^3}{3} + \frac{27(1)^2}{2} - 18(1) \right)$$

$$= -\frac{9}{4} - \left(-\frac{91}{12} \right)$$

$$= \frac{16}{3}$$

$$\int_3^6 (x^3 - 10x^2 + 27x - 18) dx$$

$$= \left[\frac{x^4}{4} - \frac{10x^3}{3} + \frac{27x^2}{2} - 18x \right]_3^6$$

$$= \left(\frac{(6)^4}{4} - \frac{10(6)^3}{3} + \frac{27(6)^2}{2} - 18(6) \right)$$

$$- \left(\frac{(3)^4}{4} - \frac{10(3)^3}{3} + \frac{27(3)^2}{2} - 18(3) \right)$$


$$= -18 - \left(-\frac{9}{4} \right)$$

$$= -\frac{63}{4}$$

$$\therefore \text{Area} = \frac{16}{3} + \frac{63}{4} = \frac{253}{12} \text{ square units}$$


11.5 The Area Between Two Curves

The area **between** two curves can be calculated using the subtraction of one definite integral from another:



Note that $\int_a^b f(x) \, dx - \int_a^b g(x) \, dx = \int_a^b (f(x) - g(x)) \, dx$, leading to the following formula:


Area Between Two Curves:



$$\text{Area} = \int_a^b (f(x) - g(x)) \, dx \quad \text{or} \quad \text{Area} = \int_a^b (\text{Upper} - \text{Lower}) \, dx$$

Example 11.5.1

Part of the graphs of $y = -3x$ and $y = 4 - x^2$ are shown. Calculate the shaded area.



$$\int_{-1}^4 (4 - x^2 - (-3x)) dx$$

← Upper – Lower

$$\int_{-1}^4 (4 - x^2 + 3x) dx$$

← Simplify

$$= \left[4x - \frac{x^3}{3} + \frac{3x^2}{2} \right]_{-1}^4$$

← Integrate

$$= \left(4(4) - \frac{(4)^3}{3} + \frac{3(4)^2}{2} \right) - \left(4(-1) - \frac{(-1)^3}{3} + \frac{3(-1)^2}{2} \right)$$

← Substitute

$$= \frac{56}{3} - \left(-\frac{13}{6} \right)$$

← Evaluate

$$= \frac{125}{6} \text{ square units}$$

Review Exercise

12.1 The Addition Formulae

It should be noted that $\sin(A + B) \neq \sin A + \sin B$:

$$\sin(60^\circ + 30^\circ) = \sin 90^\circ = 1$$

$$\text{but } \sin 60^\circ + \sin 30^\circ = \frac{\sqrt{3}}{2} + \frac{1}{2} = \frac{\sqrt{3} + 1}{2}$$

Instead, the following expansions are provided in the formula sheet, known as the **addition formulae**:

Addition Formulae (as provided on the formula sheet):

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

They can be expanded into four formulae, covering addition and subtraction for each of sin and cos:

$$\sin(A + B) = \sin A \cos B + \cos A \sin B$$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B$$

$$\sin(A - B) = \sin A \cos B - \cos A \sin B$$

$$\cos(A - B) = \cos A \cos B + \sin A \sin B$$

They can be used to expand the usage of known *exact values* ($0^\circ, 30^\circ, 45^\circ, 60^\circ, 90^\circ, 180^\circ, 270^\circ, 360^\circ$).

Example 12.1.1

Evaluate $\sin 105^\circ$.

$$\sin 105^\circ = \sin(60^\circ + 45^\circ)$$

← Relate to known exact values

$$= \sin 60^\circ \cos 45^\circ + \cos 60^\circ \sin 45^\circ$$

← Expand using formula

$$= \frac{\sqrt{3}}{2} \times \frac{1}{\sqrt{2}} + \frac{1}{2} \times \frac{1}{\sqrt{2}}$$

← Substitute exact values

$$= \frac{\sqrt{3}}{2\sqrt{2}} + \frac{1}{2\sqrt{2}}$$

$$= \frac{\sqrt{3} + 1}{2\sqrt{2}}$$

← Simplify

Example 12.1.2


Evaluate $\sin 40^\circ \cos 10^\circ - \cos 40^\circ \sin 10^\circ$.

$$\begin{aligned}\sin 40^\circ \cos 10^\circ - \cos 40^\circ \sin 10^\circ &= \sin (40^\circ - 10^\circ) && \leftarrow \text{Recognise expanded form of } \sin(A - B) \\ &= \sin 30^\circ \\ &= \frac{1}{2} && \leftarrow \text{State exact value}\end{aligned}$$

12.2 Applications to Right-Angled Triangles

Given a right-angled triangle containing an angle x , values of $\sin x$, $\cos x$ and $\tan x$ can be stated using **SOHCA TOA**. For example with Pythagoras' Theorem used to find a missing side length if needed.

By Pythagoras: $\sqrt{5^2 + 3^2} = \sqrt{34}$



$$\sin x = \frac{\text{opp}}{\text{hyp}} = \frac{3}{\sqrt{34}}$$


$$\cos x = \frac{\text{adj}}{\text{hyp}} = \frac{5}{\sqrt{34}}$$

$$\tan x = \frac{\text{opp}}{\text{adj}} = \frac{3}{5}$$

These skills are often required in combination with applications of the addition formulae.

Example 12.2.1

The diagram below shows angles p and q contained within right-angled. Find $\sin(p + q)$.



Opposite side for $p = 2$ \leftarrow By Pythagoras: $\sqrt{\sqrt{20}^2 - 4^2} = \sqrt{20 - 16} = \sqrt{4} = 2$

Hypotenuse for $q = \sqrt{40}$ \leftarrow By Pythagoras: $\sqrt{6^2 + 2^2} = \sqrt{36 + 4} = \sqrt{40}$

Once all side length have been calculated, values of $\sin p$, $\cos p$, $\sin q$, $\cos q$ can be obtained.

$$\sin(p + q) = \sin p \cos q + \cos p \sin q$$

\leftarrow Expand addition formula

$$= \frac{2}{\sqrt{20}} \times \frac{6}{\sqrt{40}} + \frac{4}{\sqrt{20}} \times \frac{2}{\sqrt{40}}$$

\leftarrow Substitute using SOHCAHTOA

$$= \frac{12}{\sqrt{800}} + \frac{8}{\sqrt{800}}$$

$$= \frac{20}{20\sqrt{2}}$$

$$= \frac{1}{\sqrt{2}}$$

\leftarrow Simplify (and rationalise if required)

Other expansions such as $\cos(p + q)$ can be calculated similarly, and note that $\tan(p + q) = \frac{\sin(p + q)}{\cos(p + q)}$.

12.3 The Double Angle Formulae

Expansions for $\sin 2x$ and $\cos 2x$ in terms of $\sin x$ and $\cos x$ can be obtained from the addition formulae:


$$\begin{aligned}\sin 2x &= \sin(x + x) \\ &= \sin x \cos x + \cos x \sin x \\ &= 2 \sin x \cos x\end{aligned}$$

$$\begin{aligned}\cos 2x &= \cos(x + x) \\ &= \cos x \cos x - \sin x \sin x \\ &= \cos^2 x - \sin^2 x \\ &= 2 \cos^2 x - 1 \\ &= 1 - 2 \sin^2 x\end{aligned}$$

The other arrangements of the $\cos 2x$ expansion come from the trigonometric identity $\sin^2 x + \cos^2 x = 1$. These expansions are provided in the formula sheet. Care is needed to determine *whether* to expand a *double angle* trig term and, in the case of $\cos 2x$, *which* expansion to use.

Example 12.3.1

The diagram below shows angle p contained within a right-angled triangle. Find $\sin(2p)$.




Example 12.3.2

Given $\sin q = \frac{4}{11}$, where $0 < q < \frac{\pi}{2}$, find $\cos 2q$.

Review Exercise

The circle a fundamental mathematical shape, and in this chapter its equation will be introduced.


For all points (x, y) on a circle, the distance to the *centre* is the same, and that distance is the *radius*.



In order to fully define a circle lying on an x - y Cartesian plane, only two properties are required:

- The **radius** of the circle.
- The **centre** of the circle.

For any point (x, y) on a circle with **radius** r and **centre** (a, b) , a right-angle triangle can be constructed:



By Pythagoras' Theorem, the set of points (x, y) which lie on the circle satisfy the relationship:

$$(x - a)^2 + (y - b)^2 = r^2$$

This is known as the **equation of the circle**.

13.1 The Equation of the Circle

The equation of the circle, introduced on the previous page, is given in the formula list:

The Equation of the Circle (as provided on the formula sheet):

The circle with centre (a, b) and radius r has equation: $(x - a)^2 + (y - b)^2 = r^2$

When stating the equation of a circle, it is not necessary to expand the brackets unless specified.

Example

State the equation of the circle with centre $(5, -2)$ and diameter $6\sqrt{2}$.

$$\text{centre} = (5, -2)$$

← State centre (a, b)

$$\text{radius} = 3\sqrt{2}$$

← State radius r

$$\text{equation: } (x - 5)^2 + (y - (-2))^2 = (3\sqrt{2})^2$$

← Substitute into equation

$$(x - 5)^2 + (y + 2)^2 = 18$$

← Simplify

The radius and centre of a circle may be quickly stated given its equation:

Example

State the radius and centre of the circle with equation $(x + 4)^2 + (y - 1)^2 = 25$.

$$\text{centre} = (-4, 1)$$

← State centre (a, b)

$$\text{radius} = 5$$

← State radius r

Knowledge from the earlier chapters such as *Straight Lines* and *Vectors* may be required, such as:

The Distance Formula:

The distance between (x_1, y_1) and (x_2, y_2) :

$$\text{distance} = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

The Midpoint Formula:

The midpoint between (x_1, y_1) and (x_2, y_2) :

$$\text{midpoint} = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$

Example

Find the equation of the circle with line PQ as its diameter, where $P(-1, 2)$ and $Q(7, -4)$.

$$\text{centre} = \left(\frac{-1 + 7}{2}, \frac{2 + (-4)}{2} \right) = (3, -1)$$

← Find centre as midpoint of diameter

$$\text{radius} = \sqrt{(3 - (-1))^2 + (-1 - 2)^2} = \sqrt{25} = 5$$

← Calculate radius using distance formula

$$(x - 3)^2 + (y + 1)^2 = 25$$

← Substitute into equation and simplify

Note that **concentric** circles have the same centre and **congruent** circles have the same radius.

Exercise 13.1

13.2 The General Equation of the Circle

The equation of the circle with centre $(3, -4)$ and radius 7 can be expanded:

$$\begin{aligned}(x - 3)^2 + (y + 4)^2 &= 49 \\(x^2 - 6x + 9) + (y^2 + 8y + 16) &= 49 \\x^2 + y^2 - 6x + 8y + 9 + 16 - 49 &= 0 \\x^2 + y^2 - 6x + 8y - 24 &= 0\end{aligned}$$

This expanded form is called the **general equation of the circle**.

General Equation of the Circle (as provided on the formula sheet):

The circle with centre $(-g, -f)$ and radius $\sqrt{g^2 + f^2 - c}$ has equation:

$$x^2 + y^2 + 2gx + 2fy + c = 0$$

Unless specified, there is no requirement to expand the equation of the circle into the general form.

It is necessary, however, to be able to determine the radius and centre of a circle given its general equation.

By equating x and y coefficients, and constant terms, the radius and centre can be found:

Example

A circle has equation $x^2 + y^2 + 14x - 12y + 9 = 0$. Determine its centre and radius.

$$x^2 + y^2 + 2gx + 2fy + c = 0 \quad \leftarrow \text{Write down general equation in } g, f, c$$

$$x^2 + y^2 + 14x - 12y + 9 = 0 \quad \leftarrow \text{Align equation of circle given}$$

$$g = 7 \quad f = -6 \quad c = 9 \quad \leftarrow \text{Find the values of } g, f, c$$

$$\text{centre} = (-g, -f) \quad \leftarrow \text{Find centre}$$

$$= (-7, -(-6))$$

$$= (-7, 6)$$

$$\text{radius} = \sqrt{g^2 + f^2 - c} \quad \leftarrow \text{Find radius}$$

$$= \sqrt{7^2 + (-6)^2 - 9}$$

$$= \sqrt{49 + 36 - 9}$$

$$= \sqrt{76}$$

$$= 2\sqrt{19}$$


Note that the equation $x^2 + y^2 + 2gx + 2fy + c = 0$ only defines a circle where $g^2 + f^2 - c > 0$.

Exercise 13.2

13.3 The Intersection of a Line and a Circle


Given a circle and a line lying on the x - y Cartesian plane, one of three possible relationships must apply:

the line and circle
intersect at two points




$$b^2 - 4ac > 0$$

the line meets the
circle at a tangent



$$b^2 - 4ac = 0$$

the line does not
intersect the circle



$$b^2 - 4ac < 0$$

Whether seeking to *find points of intersection*, *find the point of contact for a tangent*, or *show that the line and circle do not intersect*, the process should begin with preparing to solve simultaneous equations.

Since the equation of a circle is difficult to write in the form “ $y = \dots$ ”, solving simultaneous equations to find points of intersection between a line and a circle will require the **substitution method**.

Example

Show that the line with the equation $y - x = -7$ is a tangent to the circle with the equation $x^2 + y^2 - 4x + 2y - 3 = 0$, and find the point of contact.

$$y = x - 7$$

← Prepare the equation of the line

$$x^2 + (x - 7)^2 - 4x + 2(x - 7) - 3 = 0$$

← Substitute into the circle's equation

$$x^2 + (x^2 - 14x + 49) - 4x + 2x - 14 - 3 = 0$$

← Expand the brackets

$$2x^2 - 16x + 32 = 0$$

← Collect like terms

$$x^2 - 8x + 16 = 0$$

← Divide through by common factor of

$$(x - 4)(x - 4) = 0$$

← Factorise

repeated factor \Rightarrow tangency

← “Show that...”

$$x = 4$$

← Solve to find x -coordinate

$$y = 4 - 7$$

← Substitute to find y -coordinate

$$= -3$$

Point of contact is $(4, -3)$

Tangency could also be shown by considering the *discriminant* ($b^2 - 4ac$) of the equation $x^2 - 8x + 16 = 0$. However, when the point of contact is required it is quicker to use the *repeated factor* instead.

To show that a line and circle do not intersect, the discriminant must be considered, as it is not sufficient to make a conclusion based on being unable to solve through factorising.


Exercise 13.3

13.4 Finding the Equation of a Tangent to a Circle

To find the equation of a tangent to any form of curve the following are generally required:

- A **point** which the tangent passes through.
- The **gradient** of the tangent.

In chapter 3 (Differentiation I), the gradients of tangents to curves were found through differentiation. Since the form of the equation of the circle cannot be differentiated using methods covered in this course, a geometric approach is used to find the tangent to a circle.



The radius and tangent are perpendicular:

$$m_{\text{radius}} \times m_{\text{tangent}} = -1$$

Given the point of contact (or *point of tangency*) at which a tangent meets a circle, the gradient of the radius from the centre to that point should be first determined. A tangent is **perpendicular to the radius** which meets it at the point of contact.

Example

Find the equation of the line which meets the circle with equation $x^2 + y^2 - 4x + 6y + 3 = 0$ at a tangent at the point P(5, -2).

$$x^2 + y^2 - 4x + 26y + 3 = 0$$

$$g = -2 \quad f = 3 \quad c = 3$$

$$\text{centre} = (-g, -f)$$

$$= (2, -3)$$

← Find the centre of the circle

$$m_{\text{radius}} = \frac{-3 - (-2)}{2 - 5}$$

← Find the gradient of the radius

$$= \frac{-1}{-3}$$

$$= \frac{1}{3}$$

$$m_{\text{tangent}} = -3$$

← State the gradient of the tangent

$$y - b = m(x - a)$$

$$y - (-2) = -3(x - 5)$$

← Substitute point of contact and m_{tangent}

$$y + 2 = -3x + 15$$

$$y = -3x + 17$$


Exercise 13.4

1.

13.5 Points Inside and Outside Circles


Given a circle and a point lying on the x - y Cartesian plane, one of three possible relationships must apply, depending on the *distance from the centre of the circle to the point* and the *radius*:

the point is outside the circle




distance $>$ radius

the point lies on the circle



distance $=$ radius

the point lies outside the circle



distance $<$ radius

Example

Show that the point $Q(-1, 7)$ lies outside of the circle with equation $(x - 3)^2 + (y + 5)^2 = 17$.

$$\text{centre} = (3, -5)$$

← Find the centre of the circle

$$\text{radius} = \sqrt{17}$$

← Find the radius of the circle

$$\text{distance} = \sqrt{(-1 - 3)^2 + (7 - (-5))^2}$$

← Find the distance from centre to point

$$= \sqrt{(-4)^2 + (12)^2}$$

$$= \sqrt{16 + 144}$$

$$= \sqrt{160}$$

Since $\sqrt{160} > \sqrt{17}$ the point lies outside the circle ← State conclusion

As with the equation of any curve, substituting the coordinates of a point into the equation of a circle may be used to check whether it lies on the circle or not. This can be useful when trying to determine the values of unknown constants.

Example

Given that the point $P(10, -3)$ lies on the circle with equation $x^2 + y^2 + kx + 6y - 11 = 0$, find the value of k .

$$10^2 + (-3)^2 + k(10) + 6(-3) - 11 = 0$$

← Substitute $(10, -3)$

$$100 + 9 + 10k - 18 - 11 = 0$$

← Simplify

$$10k + 80 = 0$$

$$10k = -80$$

$$k = -8$$


← Solve for k

Exercise 13.5

1.

13.6 The Intersections of Circles

Given two circles with *distance between the their centres* d and *radii* r_1 and r_2 (where $r_1 \geq r_2$), one of five possible relationships must exist between the circles:



Example

Show that circles C_1 and C_2 , with equations $(x-3)^2 + (y+6)^2 = 18$ and $(x-2)^2 + (y-1)^2 = 8$ respectively, touch externally.

$$r_1 = \sqrt{18} = 3\sqrt{2}$$

← State the radius of circle C_1

$$r_2 = \sqrt{8} = 2\sqrt{2}$$

← State the radius of circle C_2

$$\text{centre}_1 = (3, -6)$$

← State the centre of circle C_1

$$\text{centre}_2 = (2, 1)$$

← State the centre of circle C_2

$$\text{distance} = \sqrt{(3-2)^2 + (-6-1)^2}$$

← Find distance between centres

$$= \sqrt{1 + 49}$$

$$= \sqrt{50}$$

$$= 5\sqrt{2}$$

Since $3\sqrt{2} + 2\sqrt{2} = 5\sqrt{2}$ the circles touch externally

← State conclusion


Exercise

Review Exercise


In this chapter, the sums and differences of *equal-angled* trigonometric operations will be explored.

e.g. $3 \sin x^\circ + 4 \cos x^\circ$

The graphs of basic trigonometric functions such as $y = 3 \sin x^\circ$ and $y = 4 \cos x^\circ$ should be familiar:



The graph of $y = 3 \sin x^\circ + 4 \cos x^\circ$ is the same as that of $y = 5 \cos (x - 36.9)^\circ$.



Determining that $3 \sin x^\circ + 4 \cos x^\circ = 5 \cos (x - 36.9)^\circ$ allows the maximum values and minimum values of $3 \sin x^\circ + 4 \cos x^\circ$ to be determined and the values of x which produce them, and allow equations such as $3 \sin x^\circ + 4 \cos x^\circ = 2$ to be solved.

Any trigonometric expression $k_1 \sin x \pm k_2 \cos x$ can be written as either $k \sin (x \pm a)$ or $k \cos (x \pm a)$.

This chapter will cover how to do this and explore the various applications of this property.

14.1 The Wave Function using $k \cos(x - a)$

Whilst any of $k \sin(x \pm a)$ or $k \cos(x \pm a)$ may be used in general, any Higher exam question is likely to specify a particular form to use. The simplest is often $k \cos(x - a)$.

To express $3 \sin x^\circ + 4 \cos x^\circ$ in the form $k \cos(x - a)^\circ$, two key *trigonometric identities* are needed:

$$\sin^2 x + \cos^2 x = 1 \quad \text{and} \quad \tan x = \frac{\sin x}{\cos x}$$

First, $k \cos(x - a)^\circ$ can be expanded using the formula $\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$:

$$k \cos(x - a)^\circ = k \cos x^\circ \cos a^\circ + k \sin x^\circ \sin a^\circ$$

It is required that this expansion is equal to the original expression:

$$3 \sin x^\circ + 4 \cos x^\circ = k \cos x^\circ \cos a^\circ + k \sin x^\circ \sin a^\circ$$

Equating the coefficients of $\sin x^\circ$ and $\cos x^\circ$ gives a set of simultaneous equations in k and a :

$$\begin{aligned} k \sin a^\circ &= 3 && \leftarrow \text{Coefficients of } \sin x^\circ \\ k \cos a^\circ &= 4 && \leftarrow \text{Coefficients of } \cos x^\circ \end{aligned}$$

Squaring both sides of each equation and adding allows k to be determined using $\sin^2 x + \cos^2 x = 1$:

$$\begin{aligned} k^2 \sin^2 a^\circ + k^2 \cos^2 a^\circ &= 3^2 + 4^2 \\ k^2 (\sin^2 x^\circ + \cos^2 x^\circ) &= 3^2 + 4^2 \\ k^2 (1) &= 3^2 + 4^2 \\ k &= \sqrt{3^2 + 4^2} \\ k &= \sqrt{25} \\ k &= 5 \end{aligned}$$

Dividing $k \sin a^\circ$ by $k \cos a^\circ$ allows a to be calculated, noting $k \sin a^\circ$ and $k \cos a^\circ$ are both positive:

$$\begin{aligned} \frac{k \sin a^\circ}{k \cos a^\circ} &= \frac{3}{4} \\ \tan a^\circ &= \frac{3}{4} \\ a_{\text{acute}}^\circ &= \tan^{-1} \left(\frac{3}{4} \right) = 36.9^\circ \\ a^\circ &= 36.9^\circ \end{aligned} \quad \begin{array}{c|c} \checkmark \text{S} & \checkmark \text{A}^\circ \checkmark \\ \hline \text{T} & \checkmark \text{C} \checkmark \end{array}$$

Hence $3 \sin x^\circ + 4 \cos x^\circ = 5 \cos(x - 36.9)^\circ$. Here it has been assumed that $k > 0$ and $0 < a < 360$.

Some abbreviations of the working shown on the previous page are routinely permitted in the Higher exam. The following example demonstrates an appropriate level of detail in its solution.

Example

Express $3 \cos x - \sin x$ in the form $k \cos(x - a)$ where $k > 0$ and $0 < a < 2\pi$.

$$3 \cos x - \sin x = k \cos(x - a)$$

$$3 \cos x - \sin x = k \cos x \cos a + k \sin x \sin a \quad \leftarrow \text{Expand } k \cos(x - a)$$

$$k \sin a = -1$$

\leftarrow Equate $\sin x$ coefficients

$$k \cos a = 3$$

\leftarrow Equate $\cos x$ coefficients

$$k = \sqrt{(-1)^2 + 3^2}$$

\leftarrow Calculate k

$$= \sqrt{10}$$

$$\tan a = \frac{-1}{3}$$

\leftarrow Find $\tan a$

$$\text{acute}^\circ = \tan^{-1}\left(\frac{1}{3}\right) = 18.4^\circ$$

\leftarrow Find acute angle first

| | |
|-----|------|
| S | A ✓ |
| ✓ T | C ✓✓ |

$360^\circ - \text{acute}^\circ$

\leftarrow Negative $\sin a$: quadrants T and C

\leftarrow Positive $\cos a$: quadrants A and C

$$a^\circ = 341.6^\circ$$

\leftarrow Using $360^\circ - 18.4^\circ$ (Double-ticked quadrant)

$$a = 5.96$$

\leftarrow Convert to radians: $341.6 \times \frac{\pi}{180}$

$$3 \cos x - \sin x = \sqrt{10} \cos(x - 5.96)$$

\leftarrow State solution

Exercise 14.1

1. Express each of the following in the form $k \cos(x - a)^\circ$, where $k > 0$ and $0 < a < 360$.

(a) $5 \cos x^\circ + 12 \sin x^\circ$

(b) $4 \sin x^\circ + 5 \cos x^\circ$

(c) $6 \cos x^\circ + \sin x^\circ$

2. Express each of the following in the form $k \cos(x - a)^\circ$, where $k > 0$ and $0 < a < 360$.

(a) $8 \sin x^\circ - 6 \cos x^\circ$

(b) $\cos x^\circ - 3 \sin x^\circ$

(c) $-2 \sin x^\circ - \cos x^\circ$

3. Express each of the following in the form $k \cos(x - a)^\circ$, where $k > 0$ and $0 < a < 360$.

(a) $\sin x^\circ + \cos x^\circ$

(b) $\cos x^\circ - \sqrt{3} \sin x^\circ$

(c) $-\sin x^\circ - \sqrt{3} \cos x^\circ$

14.2 Other Forms of the The Wave Function

As well as $k \cos(x - a)$, three other forms may be used, shown below including their expansions:

$$k \cos(x + a) = k \cos x \cos a - k \sin x \sin a$$

$$k \sin(x + a) = k \sin x \cos a + k \cos x \sin a$$

$$k \sin(x - a) = k \sin x \cos a - k \cos x \sin a$$

Example

Express $12 \cos x^\circ - 5 \sin x^\circ$ in the form $k \sin(x - a)^\circ$ where $k > 0$ and $0 < a < 360$.

$$12 \cos x^\circ - 5 \sin x^\circ = k \sin(x - a)^\circ$$

$$12 \cos x^\circ - 5 \sin x^\circ = k \sin x^\circ \cos a^\circ - k \cos x^\circ \sin a^\circ \quad \leftarrow \text{Expand } k \sin(x - a)^\circ$$

$$-k \sin a^\circ = 12 \implies k \sin a^\circ = -12 \quad \leftarrow \text{Equate } \sin x^\circ \text{ coefficients}$$

$$k \cos a^\circ = -5 \quad \leftarrow \text{Equate } \cos x^\circ \text{ coefficients}$$

$$k = \sqrt{(-12)^2 + (-5)^2} \quad \leftarrow \text{Calculate } k$$

$$= 13$$

$$\tan a^\circ = \frac{-12}{-5} = \frac{12}{5} \quad \leftarrow \text{Find } \tan a^\circ$$

$$\text{acute}^\circ = \tan^{-1}\left(\frac{12}{5}\right) = 67.4^\circ \quad \leftarrow \text{Find acute angle first}$$

$$180^\circ + \text{acute}^\circ$$

\leftarrow Negative $\sin a^\circ$: quadrants T and C

\leftarrow Negative $\cos a^\circ$: quadrants S and T

$$a^\circ = 247.4^\circ \quad \leftarrow \text{Using } 180^\circ + 67.4^\circ \text{ (Double-ticked)}$$

$$12 \cos x^\circ - 5 \sin x^\circ = 13 \sin(x - 247.4)^\circ \quad \leftarrow \text{State solution}$$

Exercise 14.2

1. Express each of the following in the form $k \sin(x + a)^\circ$, where $k > 0$ and $0 < a < 360$.

(a) $3 \cos x^\circ + 2 \sin x^\circ$

(b) $4 \sin x^\circ + 3 \cos x^\circ$

(c) $7 \cos x^\circ - \sin x^\circ$

2. Express each of the following in the form $k \cos(x + a)$, where $k > 0$ and $0 < a < 2\pi$.

(a) $3 \sin x + 8 \cos x$

(b) $-3 \cos x - 4 \sin x$

(c) $-6 \sin x + 2 \cos x$

3. Express each of the following in the form $k \sin(x - a)$, where $k > 0$ and $0 < a < 2\pi$.



(a) $\sqrt{3} \sin x + \cos x$

(b) $\cos x - \sin x$

(c) $2\sqrt{3} \sin x - 2 \cos x$

14.3 Maximum and Minimum Values using the Wave Function

The graphs of $y = \sin x^\circ$ and $y = \cos x^\circ$, including their turning points, should already be familiar:




Graphs of the form $k \sin(x \pm a)$ and $k \cos(x \pm a)$ have maximum/minimum values of $\pm k$, and their *horizontal translation* is described by $\pm a$ following the rules covered in the Graph Transformations chapter.

Example

Given that $3 \sin x^\circ + 4 \cos x^\circ$ can be expressed as $5 \cos(x - 36.9)^\circ$, state the minimum value of $f(x) = 3 \sin x^\circ + 4 \cos x^\circ$ and the value of x at which it occurs for $0 < x < 360$.

Sketch the graph of $y = 5 \cos x^\circ$ translated 36.9° to the right:



Use the graph to state the answer:

Hence the minimum value of $f(x)$ is -5 , which occurs when $x = 216.9$.

Exercise 14.3

1. Sketch each for $0 < x < 360$, showing the coordinates of any roots and turning points:

(a) $y = 7 \cos(x - 35)^\circ$ (b) $y = 3 \sin(x - 23)^\circ$ (c) $y = \sqrt{5} \cos(x + 41)^\circ$

2. Find the maximum value of each and the value(s) of x for which they occur for $0 < x < 360$:

(a) $8 \sin(x - 50)^\circ$ (b) $\sqrt{2} \cos(x + 27)^\circ$ (c) $7 \sin(x + 42.3)^\circ$

3. Find the minimum value of each and the value(s) of x for which they occur for $0 < x < 2\pi$:

(a) $3 \cos\left(x - \frac{\pi}{6}\right)$ (b) $5 \sin\left(x + \frac{\pi}{3}\right) + 2$ (c) $-12 \sin\left(x + \frac{\pi}{4}\right)$

4. (a) Express $6 \sin x^\circ - 7 \cos x^\circ$ in the form $k \sin(x - a)^\circ$ where $k > 0$ and $0 < a < 360$.

(b) Hence state the coordinates of the turning points of $12 \sin x^\circ - 14 \cos x^\circ$ for $0 < x < 360$.

14.4 Solving Equations using the Wave Function

Solving an equation like $3 \sin x^\circ + 4 \cos x^\circ = 2$ can be approached by using the wave function to rewrite it as $5 \cos(x - 36.9)^\circ = 2$, before solving it in the manner covered in Chapter 6.

Example

It can be shown that $5 \sin x^\circ - 2 \cos x^\circ$ can be expressed as $\sqrt{29} \sin(x - 21.8)^\circ$.
Hence, solve the equation $5 \sin x^\circ - 2 \cos x^\circ = 4$ where $0 < x < 360$.

$$5 \sin x^\circ - 2 \cos x^\circ = 4$$

$$\sqrt{29} \sin(x - 21.8)^\circ = 4$$

← Substitute the wave function form

$$\sin(x - 21.8)^\circ = \frac{4}{\sqrt{29}}$$

← Rearrange to $\sin(\dots) = \dots$

$$a = \sin^{-1}\left(\frac{4}{\sqrt{29}}\right) = 48.0^\circ$$

← Calculate acute angle

$$\begin{array}{c|c} 180^\circ - a^\circ & a^\circ \\ \hline \checkmark S & A \checkmark \\ T & C \end{array}$$

← Positive $\sin a^\circ$: quadrants A and S

$$x - 21.8^\circ = 48.0^\circ, 180^\circ - 48.0^\circ$$

← Apply ticked quadrants


$$x - 21.8^\circ = 48.0^\circ, 132.0^\circ$$

$$x^\circ = 69.8^\circ, 153.8^\circ$$

← Add 21.8° to both sides

Note that any solutions outwith the domain (often $0 < x < 360$) should have 360° added or subtracted to bring it back within the domain, where possible.

Exercise 14.4

1. Solve each equation for $0 < x < 360$: 

(a) $7 \sin(x - 18)^\circ = 4$

(b) $3 \cos(x + 34)^\circ = -2$

(c) $\sqrt{5} \sin(x - 106)^\circ + 1 = 0$

2. Given $6 \sin x^\circ - 8 \cos x^\circ = 10 \sin(x - 53.1)^\circ$, solve $6 \sin x^\circ - 8 \cos x^\circ = 5$ where $0 < x < 360$.

3. (a) Express $\sqrt{3} \sin x^\circ + \cos x^\circ$ in the form $k \cos(x - a)^\circ$ where $k > 0$ and $0 < a < 360$.


(b) Hence solve the equation $\sqrt{3} \sin x^\circ + \cos x^\circ = -1$ where $0 < x < 360$.

4. Solve each equation for $0 < x < 2\pi$: 

(a) $4 \sin(x + 0.31) + 2 = 1$

(b) $9 \cos(x + 1.24) = 5$

(c) $2\sqrt{3} \sin(x - 0.82) = \sqrt{5}$

5. (a) Express $3 \cos x + 2 \sin x$ in the form $k \sin(x + a)$ where $k > 0$ and $0 < a < 2\pi$. 

(b) Hence solve the equation $2 + 6 \cos x + 4 \sin x = 5$ where $0 < x < 2\pi$. 

14.5 Multiple Angles and Different Variables

The techniques covered in this chapter can be applied to trigonometric expressions beyond those containing only $\sin x$ and $\cos x$; they work for any sum or difference of *equal-angled* trigonometric operations.

e.g. $3 \sin t^\circ + 4 \cos t^\circ$ or $2 \sin 2x - 5 \cos 2x$

Example

Express $5 \cos 2t^\circ - 3 \sin 2t^\circ$ in the form $k \sin(2t + a)^\circ$ where $k > 0$ and $0 < a < 360$.

$$5 \cos 2t^\circ - 3 \sin 2t^\circ = k \sin(2t + a)^\circ$$

$$\underline{5 \cos 2t^\circ} - \underline{3 \sin 2t^\circ} = k \underline{\sin 2t^\circ} \cos a^\circ + k \underline{\cos 2t^\circ} \sin a^\circ \quad \leftarrow \text{Expand } k \sin(2t + a)^\circ$$

$$k \sin a^\circ = 5$$

\leftarrow Equate $\sin 2t^\circ$ coefficients

$$k \cos a^\circ = -3$$

\leftarrow Equate $\cos 2t^\circ$ coefficients

$$\begin{aligned} k &= \sqrt{(5)^2 + (-3)^2} \\ &= \sqrt{34} \end{aligned}$$

\leftarrow Calculate k

$$\tan a^\circ = \frac{5}{-3} = -\frac{5}{3}$$

\leftarrow Find $\tan a^\circ$

$$\text{acute}^\circ = \tan^{-1}\left(\frac{5}{3}\right) = 59.0^\circ$$

\leftarrow Find acute angle first

$$\begin{array}{c|c} 180^\circ - \text{acute}^\circ & \\ \hline \checkmark \checkmark \text{S} & \text{A} \checkmark \\ \checkmark \text{T} & \text{C} \end{array}$$

\leftarrow Positive $\sin a^\circ$: quadrants A and S

\leftarrow Negative $\cos a^\circ$: quadrants S and T

$$a^\circ = 121.0^\circ$$

\leftarrow Using $180^\circ - 59.0^\circ$ (Double-ticked)

$$5 \cos 2t^\circ - 3 \sin 2t^\circ = \sqrt{34} \sin(2t + 121.0)^\circ$$


\leftarrow State solution

Exercise 14.5

- Express $4 \cos t^\circ - 3 \sin t^\circ$ in the form $k \sin(t - a)^\circ$, where $k > 0$ and $0 < a < 360$.
- Express $2 \sin 2x - \cos 2x$ in the form $k \cos(2x - a)$ where $k > 0$ and $0 < a < 2\pi$.
- (a) Express $12 \cos t^\circ + 5 \sin t^\circ$ in the form $k \sin(t + a)^\circ$, where $k > 0$ and $0 < a < 360$.
 (b) Hence state:
 - The maximum value of the function $f(x) = 12 \cos t^\circ + 5 \sin t^\circ$, $0 < t < 360$.
 - The value(s) of t for which it occurs.
- (a) Express $\sin 2x - \sqrt{3} \cos 2x$ in the form $k \cos(2x - a)$ where $k > 0$, $0 < a < 2\pi$.
 (b) Hence solve $\sin 2x - \sqrt{3} \cos 2x - 1 = 0$, $0 < x < 2\pi$.

Wave Function Review Exercise

- Express $8 \sin x^\circ + 7 \cos x^\circ$ in the form $k \sin(x + a)^\circ$ where $k > 0$ and $0 < a < 360$.
- Express $\sqrt{5} \sin x + \cos x$ in the form $k \cos(x - a)$ where $k > 0$ and $0 < a < 2\pi$.
- Express $\sqrt{3} \cos t^\circ - \sin t^\circ$ in the form $k \sin(t + a)^\circ$ where $k > 0$ and $0 < a < 360$.
- Part of the graphs of $y = 3 \cos x^\circ - 5 \sin x^\circ$ and $y = -4$ are shown in the diagram below:



Points P and Q are points of intersection.

- Express $y = 3 \cos x^\circ - 5 \sin x^\circ$ in the form $k \cos(x + a)^\circ$ where $k > 0$ and $0 < a < 360$.
 - Hence determine the coordinates of P and Q.
- Express $2 \sin x^\circ - 4 \cos x^\circ$ in the form $k \sin(x - a)^\circ$ where $k > 0$ and $0 < x < 360$.
 - Hence state:
 - The maximum value of $2 \sin x^\circ - 4 \cos x^\circ$.
 - The value of x for which it occurs, where $0 < x < 360$.
 - Express $\cos x + \sqrt{3} \sin x$ in the form $k \cos(x - a)$ where $k > 0$ and $0 < a < 2\pi$.
 - Hence sketch the graph of $y = 2 \cos x + 2\sqrt{3} \sin x$ for $0 < x < 2\pi$.
 - Express $6 \cos t^\circ - 3 \sin t^\circ$ in the form $k \cos(t + a)^\circ$ where $k > 0$ and $0 < a < 360$.
 - Hence solve $2 \cos t^\circ - \sin t^\circ + 2 = 1$ where $0 < t < 360$.
 - Express $\sin 2x^\circ - \cos 2x^\circ$ in the form $k \sin(2x - a)$ where $k > 0$ and $0 < a < 360$.
 - Hence solve the equation $\sin 2x^\circ = \cos 2x^\circ$ where $0 < x < 360$.

Each part of an operation of the form a^n can be described using the following terminology:



Power functions take the form $f(x) = x^n$, with x as the base and a constant power, n .

e.g. $f(x) = x^3$

Exponential functions take the form $f(x) = a^x$, with x as the exponent and a constant base, $a > 0, a \neq 1$.

e.g. $f(x) = 3^x$

An example of an exponential function in everyday life is that of something *appreciating* by a percentage of its value, such as an antique vase of value £4000 increasing by 20% each year. Its value after 0 years, 1 years, 2 years, and so on, can be calculated as follows:

| | |
|-------------------------------|-----------------|
| $4000 \times 1.2^0 = 4000$ | ← After 0 years |
| $4000 \times 1.2^1 = 4800$ | ← After 1 year |
| $4000 \times 1.2^2 = 5760$ | ← After 2 years |
| $4000 \times 1.2^3 = 6912$ | ← After 3 years |
| $4000 \times 1.2^4 = 8294.40$ | ← After 4 years |

The function to describe its value V after x years is given by: $V(x) = 4000 \times 1.2^x$

One advantage of defining this function is the ability to calculate the value at times other than after a whole number of years. For example, the value after three and a half years can be calculated as:


$$V(3.5) = 4000 \times 1.2^{3.5} = 7571.72$$

This chapter will introduce a range of skills required when working with exponential functions.

15.1 Graphs of Exponential Functions


Where $a > 1$:

$y = a^x$ is *strictly increasing* on $x \in \mathbb{R}$:



Where $0 < a < 1$:

$y = a^x$ is *strictly decreasing* on $x \in \mathbb{R}$:



This describes **exponential growth**.

This describes **exponential decay**.

Since $a^0 = 1$, any graph of the form $y = a^x$ will pass through the point $(0, 1)$ for all $a \neq 0$.


Since $a^1 = a$, any graph of the form $y = a^x$ will pass through the point $(1, a)$ for all a .

The graph of $y = a^x$ has the x -axis as a *horizontal asymptote* - a line that it approaches but never meets.

Determining the equation of the graph of an exponential function can typically be achieved using substitution or consideration of graph transformations, along with knowledge of points $(0, 1)$ and $(1, a)$.

Example

The graph of $y = a^x + b$ is below. Find the values of a and b , and state the range of $f(x) = a^x + b$.



$$2 = a^0 + b$$

← Substitute $(0, 2)$

$$2 = 1 + b$$

$$1 = b$$

← Solve to obtain b

$$5 = a^2 + 1$$

← Substitute $(2, 5)$ and $b = 1$

$$4 = a^2$$

$$2 = a$$

← Solve to obtain a

$$y = 2^x + 1$$

← State equation


$$f(x) > 1$$

← State range


Exercise 15.1

1. Find the equation of each exponential graph using the form given, then determine the value of k .


(a) $y = a^x$




(b) $y = a^x$




(c) $y = a^x$




(d) $y = 4^x + b$




(e) $y = a^x + 2$




(f) $y = a^x + b$



(g) $y = a^x + b$



(h) $y = a^x + b$



2. Sketch the graph of each of the following, and state the range of $y = f(x)$:

(a) $y = 3^x$

(b) $y = 3^x + 1$

(c) $y = 3^x - 2$

(d) $y = 3^{x+2}$

(e) $y = (\frac{1}{2})^x$

(f) $y = (\frac{1}{2})^x - 1$

(g) $y = (\frac{1}{2})^{x-2}$

(h) $y = (\frac{1}{2})^{1-x}$

15.2 Evaluating Logarithms

The function $f(x) = \log_a x$, where $a > 0$ and $x > 0$, is defined as the inverse function to $f(x) = a^x$.

$$\begin{array}{ll} \text{If} & x = a^y \\ \text{then} & \log_a x = y \quad (\text{where } a > 0, x > 0) \end{array}$$

A numerical example will help build an understanding of how to interpret a logarithm:

$$\begin{array}{ll} \text{Since} & 8 = 2^3 \\ \text{then} & \log_2 8 = 3 \end{array}$$

Hence $\log_2 8$ ("log to the base 2 of 8") can be read as: "2 raised to *which* power gives a value of 8?"

Example

Evaluate $\log_3 81$.

$$\log_3 81 = 4 \leftarrow \text{Since } 3^4 = 81$$

Example

Evaluate $\log_5 1$.

$$\log_5 1 = 0 \leftarrow \text{Since } 5^0 = 1$$

Example

Evaluate $\log_7 49$.

$$\log_7 49 = 2 \leftarrow \text{Since } 7^2 = 49$$

Example

Evaluate $\log_3 \frac{1}{9}$.

$$\log_3 \frac{1}{9} = -2 \leftarrow \text{Since } 3^{-2} = \frac{1}{3^2} = \frac{1}{9}$$


Exercise 15.2

1. State the value of each of the following logarithms:

- | | | | | |
|-----------------|------------------|---------------------|------------------|------------------|
| (a) $\log_6 36$ | (b) $\log_5 125$ | (c) $\log_4 16$ | (d) $\log_2 16$ | (e) $\log_9 81$ |
| (f) $\log_3 81$ | (g) $\log_2 32$ | (h) $\log_{10} 100$ | (i) $\log_8 64$ | (j) $\log_4 64$ |
| (k) $\log_7 7$ | (l) $\log_3 9$ | (m) $\log_8 1$ | (n) $\log_5 25$ | (o) $\log_5 5$ |
| (p) $\log_2 64$ | (q) $\log_4 4$ | (r) $\log_2 1$ | (s) $\log_6 216$ | (t) $\log_2 128$ |

2. By considering negative and fractional indices, evaluate each:

- | | | | | |
|--------------------------|--------------------------|---------------------------|--------------------------|--------------------------|
| (a) $\log_9 3$ | (b) $\log_{64} 8$ | (c) $\log_{25} 5$ | (d) $\log_{100} 10$ | (e) $\log_8 2$ |
| (f) $\log_{27} 3$ | (g) $\log_4 2$ | (h) $\log_{16} 2$ | (i) $\log_{81} 3$ | (j) $\log_{11} 121$ |
| (k) $\log_5 \frac{1}{5}$ | (l) $\log_3 \frac{1}{3}$ | (m) $\log_6 \frac{1}{36}$ | (n) $\log_2 \frac{1}{8}$ | (o) $\log_9 \frac{1}{3}$ |

3. Calculate the value of each to three significant figures: 

- | | | | | |
|-----------------|-----------------|--------------------|------------------|------------------|
| (a) $\log_2 31$ | (b) $\log_7 50$ | (c) $\log_{10} 98$ | (d) $\log_5 120$ | (e) $\log_2 0.9$ |
|-----------------|-----------------|--------------------|------------------|------------------|

15.3 Laws of Logs

There are two key identities that can be obtained from the definition of a logarithm. For all $a > 0$:

$$\log_a 1 = 0$$

and

$$\log_a a = 1$$

There also three “log laws” that can be applied for logarithms of the *same base*. For all $a, x, y > 0$:

The **product law**:

$$\log_a (xy) = \log_a x + \log_a y$$

The **quotient law**:

$$\log_a \left(\frac{x}{y} \right) = \log_a x - \log_a y$$

The **power law**:

$$\log_a (x^y) = y \log_a x$$

Example

Evaluate $\log_6 8 + 2 \log_6 3 - \log_6 2$.

$$\begin{aligned} & \log_6 8 + 2 \log_6 3 - \log_6 2 && \leftarrow \text{Consider order of operations} \\ &= \log_6 8 + \log_6 (3^2) - \log_6 2 && \leftarrow \text{Apply power law} \\ &= \log_6 8 + \log_6 9 - \log_6 2 \\ &= \log_6 (8 \times 9) - \log_6 2 && \leftarrow \text{Apply product law} \\ &= \log_6 72 - \log_6 2 \\ &= \log_6 \left(\frac{72}{2} \right) && \leftarrow \text{Apply product law} \\ &= \log_6 36 \\ &= 2 && \leftarrow \text{Evaluate} \end{aligned}$$

Exercise 15.3

1. Evaluate each of the following:

(a) $\log_6 9 + \log_6 4$

(b) $\log_{12} 2 + \log_{12} 6$

(c) $\log_2 3 + \log_2 \frac{1}{3}$

(d) $\log_3 54 - \log_3 2$

(e) $\log_5 10 - \log_5 2$

(f) $\log_5 100 - \log_5 4$

(g) $2 \log_4 6 - \log_4 9$

(h) $\log_{12} 4 + 2 \log_{12} 6$

(i) $3 \log_6 2 + \log_6 9 - \log_6 2$

(j) $2 \log_6 3 + \log_6 \frac{2}{3}$

(k) $\log_5 \frac{1}{4} + 2 \log_5 10$

(l) $2 \log_2 6 - \log_2 12 + \log_2 \frac{1}{6}$

2. Simplify each, giving an answer in the form $\log_p q$ where p and q are positive integers:

(a) $\log_7 3 + \log_7 6 - \log_7 9$

(b) $\frac{1}{2} \log_6 9 + \log_6 5$

(c) $2 \log_2 9 - 3 \log_2 3$

3. Simplify each, giving an answer in the form $\log_a p$ where p is a positive integer:

(a) $\log_a 3 + 2 \log_a 4$

(b) $\log_a 12 - \frac{1}{3} \log_a 8$

(c) $\log_a 120 - \log_a 4 + \log_a \frac{1}{15}$

15.4 Solving Log Equations

To solve a *log equation*, it is typically desirable to have each side within a logarithm of the *same base*:

If $\log_a x = \log_a y$
 then $x = y$ (where $a > 0, x > 0, y > 0$)

“Cancelling the logs”, mathematically, is raising each side to an exponential base a , using $a^{\log_a b} = b$.

Example

Solve the equation $\log_3 x + \log_3 5 = \log_3 20$.

$$\log_3 x + \log_3 5 = \log_3 20$$

$$\log_3 5x = \log_3 20$$

← Apply product law

$$5x = 20$$

← “Cancel” \log_3 from both sides

$$x = 4$$

← Solve

To express any term as a logarithm, multiply it by $\log_a a$ (which equals 1) and then apply the power law:

Example

Solve the equation $\log_3 x - \log_3 2 = 2$.

$$\log_3 x - \log_3 2 = 2 \log_3 3$$

← Multiply the 2 by $\log_3 3$

$$\log_3 \left(\frac{x}{2} \right) = \log_3 3^2$$

← Apply quotient and power laws

$$\frac{x}{2} = 9$$

← “Cancel” \log_3 from both sides

$$x = 18$$

← Multiply both sides by 2 to solve

Exercise 15.4

1. Solve:

(a) $\log_5 3 + \log_5 x = \log_5 18$

(b) $\log_7 2x + \log_7 4 = \log_7 56$

(c) $\log_6 5 + \log_6 (x - 1) = \log_6 30$

(d) $\log_a x - 2 \log_a 3 = \log_a 18$

(e) $\log_8 2x - \log_8 3 = \log_8 1$

(f) $\log_3 8 - \log_3 x = \log_3 2$

2. Solve each:

(a) $\log_6 x + \log_6 12 = 2$

(b) $\log_4 x + \log_4 8 = 2$

(c) $\log_2 10 + \log_2 x = 4$

(d) $\log_5 x - \log_5 2 = 1$

(e) $\log_3 x - 2 \log_3 2 = 2$

(f) $\log_4 2 + 3 \log_4 x = 2$

(g) $\log_2 5 + 3 = \log_2 (x + 2)$

(h) $\log_3 (2x) - 1 = \log_3 8$

(i) $2 - \log_6 (x - 3) = 2 \log_6 3$

Some algebraic manipulations used to solve an equation may produce solutions do not, in fact, satisfy the original equation. These **extraneous solutions** can arise when solving log equations.

Since $\log_a x$ is only defined for $a > 0$ and $x > 0$, any solutions which contradict this must be discarded.

Example

Solve the equation $\log_3 x + \log_3 (x - 2) = \log_3 8$.

$$\log_3 x + \log_3 (x - 2) = \log_3 8 \quad \leftarrow \text{Note } x > 0 \text{ and } (x - 2) > 0 \text{ therefore } x > 2$$

$$\log_3 (x(x - 2)) = \log_3 8 \quad \leftarrow \text{Apply product law}$$

$$x^2 - 2x = 8 \quad \leftarrow \text{"Cancel" } \log_3 \text{ and expand}$$

$$x^2 - 2x - 8 = 0 \quad \leftarrow \text{Equate quadratic equation to zero}$$

$$(x + 2)(x - 4) = 0 \quad \leftarrow \text{Factorise}$$

$$x + 2 = 0, x - 4 = 0$$

$$x = -2, x = 4 \quad \leftarrow \text{Discard } x = -2 \text{ since } x > 2$$

All solutions for x must be checked carefully to verify that they satisfy the *original* log equation.

3. Solve for x :

(a) $\log_a x + \log_a (x + 1) = \log_a 6$

(b) $\log_4 x + \log_4 (x - 3) = \log_4 10$

(c) $\log_6 (x - 1) + \log_6 x = 1$

(d) $\log_2 (x + 6) - \log_2 x = 2$

(e) $\log_5 (x + 2) + \log_5 (x - 3) = \log_5 6$

(f) $\log_a 3 = \log_a (x + 5) + \log_a 2x$

4. Solve each:

(a) $\log_5 x^2 + \log_5 4 = \log_5 36$

(b) $\log_6 2 + \log_6 x^2 = \log_6 50$

(c) $\log_a 7 + \log_a x^2 = \log_a 14$

(d) $2 \log_a x + \log_a 3 = \log_a 48$

(e) $\log_9 4 + 2 \log_9 x = 2 \log_9 8$

(f) $2 \log_7 3x + \log_7 2 = \log_7 18$

5. Solve each for a :

(a) $\log_a 3 + \log_a 2 = 1$

(b) $\log_a 18 - \log_a 2 = 1$

(c) $\log_a 4 + \log_a 2 = 3$

(d) $\log_a 12 - \log_a 4 = \frac{1}{2}$


(e) $\log_a 16 + \log_a 4 = 2$

(f) $\log_a 50 - \log_a 2 = 2$

15.5 Graphs of Logarithmic Functions

Given a function f , defined on a suitable domain such that an inverse function exists, the graph of its inverse function $y = f^{-1}(x)$ can be obtained by **reflecting** the graph of $y = f(x)$ in the line $y = x$.

Since the inverse to the exponential function $f(x) = a^x$ is the **logarithmic function** $f^{-1}(x) = \log_a x$, knowledge of exponential graphs allows log graphs to be sketched:



The graph of $y = a^x$ passes through...

$(0, 1)$ and $(1, a)$

...and has a **horizontal** asymptote of $x = 0$.

The graph of $y = \log_a x$ passes through...

$(1, 0)$ and $(a, 1)$.

..and has a **vertical** asymptote of $y = 0$.


In general, if $y = f(x)$ passes through (x, y) then $y = f^{-1}(x)$ passes through (y, x) .

Knowledge of the graph of $y = \log_a x$ may be combined with an understanding of graph transformations.

Example

Sketch the graph of $y = \log_3(x - 1) + 2$.

Consider $y = \log_3(x - 1) + 2$ as a $f(x - 1) + 3$ transformation of $y = \log_3 x$:



Exercise 15.5

1. Sketch each of the following:

(a) $y = \log_5 x$

(b) $y = \log_2 x$

(c) $y = \log_9 x$

2. Sketch each of the following, showing clearly where any asymptote is not along an axis:

(a) $y = \log_4 x + 1$

(b) $y = \log_7 x - 1$

(c) $y = \log_5 x + 2$

(d) $y = \log_8(x + 5)$

(e) $y = \log_2(x + 3)$

(f) $y = \log_3(x - 2)$

(g) $y = \log_7(x + 3) + 2$


(h) $y = \log_5(x - 3) + 1$

(i) $y = \log_6(x + 2) - 1$

As in section 15.1 (Graphs of Exponential Functions), substitution of known coordinates on the graph of a log function into its equation can allow unknown constants it contains to be determined.

Example

The graph of $y = \log_a(x + b)$ is below. Find the values of a and b .



$$0 = \log_a(-2 + b)$$

← Substitute $(-2, 0)$

$$-2 + b = 1$$

← Since $\log_a 1 = 0$ for all a

$$b = 3$$

← Solve to obtain b

$$1 = \log_a(2 + 3)$$

← Substitute $(2, 1)$ and $b = 3$

$$a = 5$$


← Since $\log_a a = 1$ for all a

$$y = \log_5(x + 3)$$


← State equation

3. Determine the equation of each, using the form given:


(a) $y = \log_4(x + a)$




(b) $y = \log_a(x - 2)$



(c) $y = \log_a(x + b)$



(d) $y = \log_a(bx)$




15.6 Solving Exponential Equations

To solve an *exponential equation* such as $10 = 2^x$, which has a non-integer solution, the relationship between a log and an exponential of the same base, stated in section 15.2, can be used as follows:

$$\begin{aligned} \text{If} \quad & 10 = 2^x \\ \text{then} \quad & \log_2 10 = x \end{aligned}$$

The solution, using a calculator and correct to three significant figures, is $x = 3.32$. This makes use of the property $\log_a a^x = x$, which follows from the inverse relationship between a log and exponential.

Example

Solve the equation $14 = 3 \times 5^x$, giving the answer correct to three significant figures. 

$$14 = 3 \times 5^x$$

$$\frac{14}{3} = 5^x$$

← Divide both sides by 3


$$\log_5 \left(\frac{14}{3} \right) = x$$

← Take \log_5 of both sides

$$x = 0.957 \text{ (to 3sf)}$$

← Calculate the solution

Exercise 15.6

1. Solve, giving the answers to three significant figures: 

(a) $30 = 5^x$

(b) $6 = 3^x$

(c) $13 = 8^x$

(d) $3 = 7^x$

(e) $2^x = 11$

(f) $10^x = 500$

(g) $6^x = 3$


(h) $4^x = 65$

(i) $19 = 2.1^t$

(j) $1.3^{2y} = 0.7$

(k) $7^{4p} = \frac{1}{2}$

(l) $\frac{2}{3} = 4^{0.7k}$

2. Solve, giving the answers to three significant figures: 

(a) $12 = 4 \times 8^x$

(b) $24 = 6 \times 7^x$

(c) $2 = 7 \times 5^x$

(d) $21 = 4 \times 3^x$

(e) $6 \times 8^x = 5$

(f) $3 \times 4^x = 45$

(g) $0.03 \times 8^x = 0.7$


(h) $18 \times 5^x = 0.7$

(i) $5 = 3 \times 4^q$

(j) $7 \times 6^{0.03t} = 5$

(k) $10 \times 10^{-3y} = 9$

(l) $10 = 9 \times 10^{-0.037k}$

3. Solve, giving the answer to three significant figures: 

(a) $20 = 4 \times 7^x + 3$

(b) $7 = \frac{1}{2} \times 3^x - 6$

(c) $5 + 3 \times 4^x = 23$

(d) $3 \times \left(\frac{1}{2}\right)^x = 7$

(e) $6 \times 1.4^x - 2.3 = 31.4$

(f) $0.4 = 3 \times 0.3^x - 0.5$

(g) $3 \times 4^{0.0723k} = 21$

(h) $11 \times 3^{-0.104x} = 6$

(i) $3 = 5 \times 2^{-0.00562k}$

15.7 Natural Logarithms

Some logarithms with particular *bases* are routinely used in various fields, such as the *binary logarithm* (\log_2) in computer science, or the *decimal* or *common logarithm* (\log_{10} or just \log) in engineering.

The most useful logarithm in mathematics is the **natural logarithm** (\log_e or more often just **ln**), which takes Euler's Number e as its base, where $e \approx 2.719$. (It lies in the ease which which exponentials in base e can be used together with calculus, which is beyond the scope of this course.)

The *laws of logs* apply to natural logarithms since they apply to *all* \log_a where $a > 0$:


Example

Simplify $\ln 48 - 3 \ln 2$, giving the answer in the form $\ln p$.

$$\begin{aligned}
 & \ln 48 - 3 \ln 2 \\
 &= \ln 48 - \ln (2^3) && \leftarrow \text{Apply the power law} \\
 &= \ln 48 - \ln 8 \\
 &= \ln \left(\frac{48}{8} \right) && \leftarrow \text{Apply the quotient law} \\
 &= \ln 6 && \leftarrow \text{Simplify}
 \end{aligned}$$


Exponential equations in base e are solved using the natural logarithm **ln**, since $\ln(e^x) = x$:

Example

Given $37 = 21e^{-0.48x}$, find the value of k to three significant figures. 

$$\begin{aligned}
 & 37 = 21e^{-0.48k} \\
 & \frac{37}{21} = e^{-0.48k} && \leftarrow \text{Divide both sides by 21} \\
 & \ln \left(\frac{37}{21} \right) = -0.48k && \leftarrow \text{Take ln of both sides} \\
 & \frac{\ln \left(\frac{37}{21} \right)}{-0.48} = k && \leftarrow \text{Divide both sides by } -0.48 \\
 & k = 1.18 \text{ (to 3sf)} && \leftarrow \text{Calculate the solution}
 \end{aligned}$$

Exercise 15.7

1. For each, find the value of k to three significant figures: 

- | | | | |
|---------------------|--------------------|----------------------|----------------------|
| (a) $0.7 = e^k$ | (b) $1.8 = e^k$ | (c) $3e^k = 24$ | (d) $4 = 7e^x$ |
| (e) $125 = 5e^{2k}$ | (f) $50 = 7e^{4k}$ | (g) $10 = 12e^{-3k}$ | (h) $17 = 100^{-5k}$ |

2. Solve each for x , giving the answers to three significant figures: 


- | | | | |
|--------------------------|--------------------------|--------------------------|------------------------|
| (a) $120 = 85e^{0.475x}$ | (b) $50 = 32e^{0.0273x}$ | (c) $14 = 23e^{-0.381x}$ | (d) $3P = Pe^{0.045x}$ |
|--------------------------|--------------------------|--------------------------|------------------------|

15.8 Exponential Growth and Decay


The behaviour of many real-world things are described by **exponential growth** or **exponential decay**.

Time, t , is a common variable for such a function in place of x , and it typically describes the behaviour of some real-world variable which is named suitably in place of y of $f(x)$, such as V for “value”, P for “population size” or M for “mass”. Where $k > 0$:

$V = V_0 e^{kt}$ describes the **exponential growth** of V from some initial value V_0 as t increases.



$V = V_0 e^{-kt}$ describes the **exponential decay** of V from some initial value V_0 as t increases.



Example

80 micrograms of a radioactive substance, which decays over time, is obtained. The mass M micrograms of the substance remaining after t hours can be modelled by the equation:

$$M = M_0 e^{-kt}$$

(a) State the value of M_0 .

$$M_0 = 80$$

← M_0 is the initial value

Constant k may be found through substitution if initial value V_0 is known and the value of V at time t .

(b) After 5 hours, only 50.7 micrograms of the substance remains.

Determine the value of k , rounded to three significant figures. 

$$50.7 = 80e^{-k \times 5}$$

← Substitute $M_0 = 80$, $M = 50.7$ and $t = 5$

$$\frac{50.7}{80} = e^{-5k}$$

← Divide by 80

$$\ln\left(\frac{50.7}{80}\right) = -5k$$

← Take natural log of both sides


$$\frac{\ln\left(\frac{50.7}{80}\right)}{-5} = k$$

← Divide both sides by (-5)

$$k = 0.0912$$

← Calculate and round as required

A feature of exponential functions is that the time taken for the value to grow or decay by any given proportion is always constant. For example the time taken for an exponential growth function to double a value from 20 to 40 will be the same time it will take to double from 50 to 100.

(c) Find the time it will take for only half of the original substance to remain (its *half-life*). 

$$40 = 80e^{-0.0912t} \quad \leftarrow \text{Substitute } M = 40 \text{ and } k = 0.0912 \text{ is now known}$$

$$\frac{1}{2} = e^{-0.0912t} \quad \leftarrow \text{Divide by 80 and simplify}$$

$$\ln\left(\frac{1}{2}\right) = -0.0912t \quad \leftarrow \text{Take natural log of both sides}$$

$$\frac{\ln\left(\frac{1}{2}\right)}{-0.0912} = t \quad \leftarrow \text{Divide both sides by } (-0.0912)$$

$$t = 7.60 \text{ hours} \quad \leftarrow \text{Calculate}$$

Exercise 15.8

All questions: 

1. For each *exponential growth* function, find the value of t it will take for the value to double:

(a) $V = 40e^{0.3t}$

(b) $P(t) = 25e^{0.08t}$

(c) $M = M_0e^{0.00572t}$

2. The number of bacteria in a population, B , can be modelled using the equation $B = 400e^{0.0376t}$ where t is time measured in minutes.

(a) State the number of bacteria in the population initially.

(b) Find, to the nearest thousand, the number of bacteria after 45 minutes.

(c) Determine the time it takes for the population of bacteria to double in size.

3. A population of deer is growing exponentially in an area, and the size of the population, P , can be described by the equation $P = 860e^{0.173t}$, where t is the number of years from 2026.

(a) State the number of deer in the population in 2026.

(b) Find, to the nearest ten, the number of deer expected to be in the population in 2028.

(c) Determine the time it will take for the population of deer to double in size.

4. For each *exponential decay* function, find the value of t it will take for the value to halve:

(a) $T = 20e^{-0.8t}$

(b) $V(t) = 360e^{-0.072t}$

(c) $P = P_0e^{-0.00197t}$

5. The temperature T (in $^{\circ}\text{C}$) of a substance t minutes after the end of an experiment can be modelled by the equation $T = T_0e^{-kt}$. After 7 minutes, the substance has cooled from 140°C to 55°C .

(a) Determine the value of k to three significant figures.

(b) Find the time taken for the substance to cool to 30°C .

6. The concentration C of a drug in a patient's blood t hours after being given a dose can be modelled by the equation $C = 28e^{-kt}$.


(a) If the concentration is 23 after 90 minutes, calculate the value of k to three significant figures.

(b) Find the time taken for the concentration to reduce by half.


(c) Find the time taken for the concentration to be only 15% of its original value.

15.9 Experimental Data of the form $y = ax^b$

Where a relationship between variables x and y is a **power** function in the form $y = ax^b$, taking any log of both sides will reveal the **linear relationship** that therefore exists between $\log x$ and $\log y$.




$$\begin{aligned}
 y &= ax^b \\
 \log y &= \log(ax^b) \\
 \log y &= \log x^b + \log a \\
 \log y &= b \log x + \log a \\
 &\quad \downarrow \quad \downarrow \\
 \log y &= m \log x + c
 \end{aligned}$$



Drawing such a plot of $\log y$ against $\log x$ and obtaining a straight line can be used to check for a power relationship. Where a relationship is of the form $y = ax^b$, determining the values of a and b is possible by first finding the *gradient* and *y-intercept* on the $\log y \sim \log x$ graph.

Example

Two variables, x and y , are connected by the equation $y = kx^n$. The graph of $\log_3 y$ against $\log_3 x$ is a straight line as shown. Determine the values of k and n .



$$\log_3 y = \log_3 (kx^n)$$

← Take \log_3 of both sides

$$\log_3 y = \log_3 x^n + \log_3 k$$

← Apply product law

$$\log_3 y = n \log_3 x + \log_3 k$$

← Apply power law

$$\text{gradient} = \frac{6 - 3}{2 - 0} = \frac{3}{2}$$

← Find gradient

$$n = \frac{3}{2}$$

← Equate to n

$$y\text{-intercept} = 3$$

← Find y -intercept

$$\log_3 k = 3$$

← Equate to $\log_3 k$

$$\log_3 k = 3 \log_3 3$$

← Solve for k





$$\log_3 k = \log_3 3^3$$

$$k = 27$$





Therefore the relationship between x and y is $y = 27x^{\frac{3}{2}}$.

Exercise 15.9

1. For each graph where the relationship between x and y is $y = kx^n$, determine the values of k and n :




2. For each graph where the relationship between x and y is $y = ax^b$, determine the values of a and b :




15.10 Experimental Data of the form $y = ab^x$

Where a relationship between variables x and y is an **exponential** function in the form $y = ab^x$, taking any log of both sides will reveal the **linear relationship** that therefore exists between x and $\log y$.




$$\begin{aligned}
 y &= ab^x \\
 \log y &= \log(ab^x) \\
 \log y &= \log b^x + \log a \\
 \log y &= x \log b + \log a \\
 &\quad \downarrow \quad \downarrow \\
 \log y &= xm + c
 \end{aligned}$$



Drawing such a plot of $\log y$ against x and obtaining a straight line can be used to check for an exponential relationship. Where a relationship is of the form $y = ab^x$, determining the values of a and b is possible by first finding the *gradient* and *y-intercept* on the $\log y \sim x$ graph.

Example

Two variables, x and y , are connected by the equation $y = ab^x$. The graph of $\log_9 y$ against x is a straight line as shown. Determine the values of a and b .



$$\log_9 y = \log_9 (ab^x)$$

← Take \log_9 of both sides

$$\log_9 y = \log_9 b^x + \log_9 a$$

← Apply product law

$$\log_9 y = x \log_9 b + \log_9 a$$

← Apply power law

$$\text{gradient} = \frac{4 - 2}{4 - 0} = \frac{2}{4} = \frac{1}{2}$$

← Find gradient

$$\log_9 b = \frac{1}{2}$$

← Equate to $\log_9 b$

$$b = 3$$

← Solve for b

$$y\text{-intercept} = 2$$

← Find y -intercept

$$\log_9 a = 2$$

← Equate to $\log_9 a$

$$a = 81$$





← Solve for a

Therefore the relationship between x and y is $y = 81 \times 3^x$.





Note the the process of solving for a and solving for b may require solving each log equation step-by-step.

Exercise 15.10

1. For each graph where the relationship between x and y is $y = ab^x$, determine the values of a and b :



2. For each graph where the relationship between x and y is $y = ka^x$, determine the values of a and k :



Logs and Exponentials Review Exercise

1. Sketch each of the following:

(a) $y = 4^x + 2$

(b) $y = 5^{x+2}$

(c) $\log_3(x-2) + 4$

2. Express $2 \log_p 10 + \log_p 2 - \log_p 50$ in the form $\log_p q$, where q is a positive integer.

3. Evaluate the following:

(a) $\log_8 + \log_8 - \log_8$

(b) $\log_3 \frac{1}{4} + 2 \log_3 6$

(c) $\frac{1}{2} \log_5 64 - \log_5 40$

4. Solve each equation:

(a) $\ln x - \ln 5 = \ln 2$

(b) $\log_6 x + \log_6 4 = 2$

(c) $\log_2 x + \log_2(x+7) = 3$

5. Given that $\log_p 4 + 3 = \log_p 32$, find the value of p .

6. A population of mice is discovered living in some woodland, and number of mice can be modelled by:

$$M = 80e^{kt}$$



where M is the estimated number of mice in the population, t is the number of weeks since the population was discovered and k is a constant.

(a) State the estimated number of mice in the population when it was first discovered.


(b) It is estimated there will be 120 mice in the population after 3 weeks. Find the value of k .

(c) Calculate the time taken for the population to double in size.

7. Variables x and y are linked by the equations given below. Find the values of a and b for each:



8. The graph of $y = f(x)$ is shown below where $f(x) = \log_3(x+a) + b$.



(a) State the domain of $f(x)$.

(b) Find the values of a and b .

(c) Sketch the graph of $y = f^{-1}(x)$.

16.1 Differentiating Trig Functions

Exercise

16.2 Integrating Trig Functions

Exercise

16.3 The Chain Rule for Differentiation

Exercise

16.4 Integrating $(ax + b)^n$

Exercise

Review Exercise

CHALLENGE PROBLEMS

The following problems **do not** represent the kind of question expected to feature in a Higher Mathematics exam, either in the way they are presented or the level of difficulty. Instead, they aim to encourage a flexible approach towards problem-solving and an understanding that the skills covered in the course have applications beyond those featured in any typical exam. *Some questions may be solvable without using the skills covered in this chapter, and some questions may be unrelated to this chapter.*

Warm-Up Problems

Chapter 1 Problems

Chapter 2 Problems

Chapter 3 Problems

Chapter 4 Problems

Chapter 5 Problems

Chapter 6 Problems

Chapter 7 Problems

Chapter 8 Problems

Chapter 9 Problems

Chapter 10 Problems

Chapter 11 Problems

Chapter 12 Problems

Chapter 13 Problems

Chapter 14 Problems

Chapter 15 Problems

Chapter 16 Problems

End of Course Problems

ANSWERS

Chapter 1 Answers

Chapter 2 Answers

Chapter 3 Answers

Chapter 4 Answers

Chapter 5 Answers

Chapter 6 Answers

Chapter 7 Answers

Chapter 8 Answers

Chapter 9 Answers

Chapter 10 Answers

Chapter 11 Answers

Chapter 12 Answers

Chapter 13 Answers

Chapter 14 Answers

Exercise 14.1


1. (a) $13 \cos(x - 67.4)^\circ$ (b) $\sqrt{41} \cos(x - 38.7)^\circ$ (c) $\sqrt{37} \cos(x - 9.5)^\circ$
2. (a) $10 \cos(x - 126.9)^\circ$ (b) $\sqrt{10} \cos(x - 288.4)^\circ$ (c) $\sqrt{5} \cos(x - 243.4)^\circ$
3. (a) $\sqrt{2} \cos(x - 45)^\circ$ (b) $2 \cos(x - 300)^\circ$ (c) $2 \cos(x - 210)^\circ$

Exercise 14.2

1. (a) $\sqrt{13} \sin(x + 56.3)^\circ$ (b) $5 \sin(x + 36.9)^\circ$ (c) $5\sqrt{2} \sin(x + 98.1)^\circ$
2. (a) $\sqrt{73} \cos(x + 5.92)$ (b) $5 \cos(x + 2.21)$ (c) $2\sqrt{10} \cos(x + 1.25)$
3. (a) $2 \sin(x - \frac{5\pi}{6})$ (b) $\sqrt{2} \sin(x - \frac{5\pi}{4})$ (c) $4 \sin(x - \frac{\pi}{6})$


Exercise 14.3

1. Sketches showing turning points and roots:




a) $y = 7 \cos(x - 35)^\circ$:

- maximum turning point at $(35^\circ, 7)$
- minimum turning point at $(215^\circ, -7)$
- x -intercepts at $(125^\circ, 0)$ and $(305^\circ, 0)$



b) $y = 3 \sin(x - 23)^\circ$:

- maximum turning point at $(113^\circ, 3)$
- minimum turning point at $(293^\circ, -3)$
- x -intercepts at $(23^\circ, 0)$ and $(203^\circ, 0)$



c) $y = \sqrt{5} \cos(x + 41)^\circ$:

- maximum turning point at $(319^\circ, \sqrt{5})$
- minimum turning point at $(139^\circ, -\sqrt{5})$
- x -intercepts at $(49^\circ, 0)$ and $(229^\circ, 0)$

2. (a) max value 8 when $x^\circ = 140^\circ$ (b) max value $\sqrt{2}$ when $x^\circ = 333^\circ$ (c) max value 7 when $x^\circ = 47.7^\circ$
3. (a) min value -3 when $x = \frac{7\pi}{6}$ (b) min value -3 when $x = \frac{7\pi}{6}$ (c) min value -12 when $x = \frac{\pi}{4}$
4. (a) $\sqrt{85} \sin(x - 49.4)^\circ$ (b) max TP at $(139.4^\circ, 2\sqrt{85})$, min TP at $(219.4^\circ, -2\sqrt{85})$

Exercise 14.4


1. (a) $x^\circ = 52.8^\circ, 163.2^\circ$ (b) $x^\circ = 97.8^\circ, 194.2^\circ$ (c) $x^\circ = 79.4^\circ, 312.6^\circ$
2. $x^\circ = 83.1^\circ, 203.1^\circ$
3. (a) $2 \cos(x - 60)^\circ$ (b) $x^\circ = 180^\circ, 300^\circ$
4. (a) $x = 3.08, 5.72$ (b) $x = 4.06, 6.02$ (c) $x = 1.52, 3.26$
5. (a) $\sqrt{13} \sin(x + 0.98)$ (b) $x = 1.73, 5.73$

Exercise 14.5

1. $5 \sin(t - 233.1)^\circ$
2. $\sqrt{5} \cos(2x - 2.04)$
3. (a) $13 \sin(t + 67.4)^\circ$ (b) max value of 13 occurs when $t = 22.6$
4. (a) $2 \cos(2x - \frac{5\pi}{6})$ (b) $x = \frac{\pi}{4}, \frac{7\pi}{12}, \frac{5\pi}{4}, \frac{19\pi}{12}$

Review Exercise

1. $\sqrt{113} \sin(x + 41.2)^\circ$
2. $\sqrt{6} \cos(x - 1.15)$
3. $2 \sin(x + \frac{2\pi}{3})$
4. (a) $\sqrt{34} \cos(x + 59.0)^\circ$ (b) P($74.3^\circ, -4$) and Q($167.7^\circ, -4$)
5. (a) $2\sqrt{5} \sin(x - 63.4)^\circ$ (b) i. max value is $2\sqrt{5}$ ii. occurs when $x = 153.4$
6. (a) $2 \cos(x - \frac{\pi}{3})$
(b) Sketch of $y = \cos x + \sqrt{3} \sin x$:





7. (a) $3\sqrt{5} \cos(x + 26.6)^\circ$ (b) $x = 90, 216.9$
8. (a) $\sqrt{2} \sin(2x - 45)^\circ$ (b) $x = 22.5, 112.5, 202.5, 292.5$

Chapter 15 Answers



Exercise 15.1

1. (a) $y = 3^x$
 $k = 9$ (b) $y = \left(\frac{1}{2}\right)^x$
 $k = 1$ (c) $y = 2^x$
 $k = 8$ (d) $y = 4^x - 5$
 $k = -5$
- (e) $y = 2^x + 2$
 $k = 2$ (f) $y = 2^x + 3$
 $k = 7$ (g) $y = 4^x - 1$
 $k = 1$ (h) $y = 3^x - 2$
 $k = -\frac{5}{3}$
2. (a) $y = 3^x$, range $f(x) > 0$ (b) $y = 3^x + 1$, range $f(x) > 1$





(c) $y = 3^x - 2$, range $f(x) > -2$

(d) $y = 3^{x+2}$, range $f(x) > 0$





(e) $y = \left(\frac{1}{2}\right)^x$, range $f(x) > 0$

(f) $y = \left(\frac{1}{2}\right)^x - 1$, range $f(x) > -1$



(g) $y = \left(\frac{1}{2}\right)^{x-2}$, range $f(x) > 0$

(h) $y = \left(\frac{1}{2}\right)^{1-x}$, range $f(x) > 0$



Exercise 15.2

1. (a) 2 (b) 3 (c) 2 (d) 4 (e) 2
 (f) 4 (g) 5 (h) 2 (i) 2 (j) 3
 (k) 1 (l) 2 (m) 0 (n) 2 (o) 1
 (p) 6 (q) 1 (r) 0 (s) 3 (t) 7
2. (a) $\frac{1}{2}$ (b) $\frac{1}{2}$ (c) $\frac{1}{2}$ (d) $\frac{1}{2}$ (e) $\frac{1}{3}$
 (f) $\frac{1}{3}$ (g) $\frac{1}{2}$ (h) $\frac{1}{4}$ (i) $\frac{1}{4}$ (j) 2
 (k) -1 (l) -1 (m) -2 (n) -3 (o) $-\frac{1}{2}$
3. (a) 4.95 (b) 2.01 (c) 1.99 (d) 2.97 (e) -0.152

Exercise 15.3


1. (a) 2 (b) 1 (c) 0
 (d) 3 (e) 1 (f) 2
 (g) 1 (h) 2 (i) 2
 (j) 1 (k) 2 (l) -1
2. (a) $\log_7 2$ (b) $\log_6 15$ (c) $\log_2 3$
3. (a) $\log_a 48$ (b) $\log_a 6$ (c) $\log_a 2$

Exercise 15.4


1. Solve:
 (a) $x = 6$ (b) $x = 7$ (c) $x = 7$ (d) $x = 54$ (e) $x = \frac{3}{2}$ (f) $x = 4$
2. Solve each:
 (a) $x = 3$ (b) $x = 2$ (c) $x = 1.6$ (d) $x = 10$ (e) $x = 36$
 (f) $x = 2$ (g) $x = 38$ (h) $x = 12$ (i) $x = 7$
3. (a) $x = 2$ (b) $x = 5$ (c) $x = 3$ (d) $x = 2$ (e) $x = 4$ (f) $x = \frac{1}{2}$
4. (a) $x = 3$ (b) $x = 5$ (c) $x = \sqrt{2}$ (d) $x = 4$ (e) $x = 4$ (f) $x = 1$
5. (a) $a = 6$ (b) $a = 9$ (c) $a = 2$ (d) $a = 9$ (e) $a = 8$ (f) $a = 5$

Exercise 15.5


1. (a) $y = \log_5 x$




(b) $y = \log_2 x$




(c) $y = \log_9 x$




2. (a) $y = \log_4 x + 1$




(b) $y = \log_7 x - 1$




(c) $y = \log_5 x + 2$




(d) $y = \log_8(x - 5)$




(e) $y = \log_2(x + 3)$




(f) $y = \log_3(x - 2)$




(g) $y = \log_7(x + 3) + 2$



(h) $y = \log_5(x - 3) + 1$



(i) $y = \log_6(x + 2) - 1$



3. (a) $y = \log_4(x + 2)$

(b) $y = \log_3(x - 2)$

(c) $y = \log_5(x - 1)$

(d) $y = \log_6 2x$

Exercise 15.6

1. (a) $x =$ (b) $x =$ (c) $x =$ (d) $x =$ (e) $x =$ (f) $x =$

(g) $x =$ (h) $x =$ (i) $x =$ (j) $x =$ (k) $x =$ (l) $x =$

2. (a) $x =$ (b) $x =$ (c) $x =$ (d) $x =$ (e) $x =$ (f) $x =$

(g) $x =$ (h) $x =$ (i) $x =$ (j) $x =$ (k) $x =$ (l) $x =$

3. (a) $x =$ (b) $x =$ (c) $x =$ (d) $x =$ (e) $x =$

(f) $x =$ (g) $x =$ (h) $x =$ (i) $x =$

Exercise 15.7

Exercise 15.8

Exercise 15.9

Exercise 15.10

Review Exercise

Chapter 16 Answers