

Contents

Introduction	1
1 The Straight Line	2
2 Recurrence Relations	3
3 Differentiation I	4
4 Quadratic Theory	5
5 Sets and Functions	6
6 Trigonometry	7
7 Graph Transformations	8
8 Vectors	9
9 Differentiation II	10
10 Polynomials	11
11 Integration	12
11.1 Indefinite Integrals	13
11.2 Differential Equations	15
11.3 Definite Integrals	16
11.4 The Area Under a Curve	17
11.5 The Area Between Two Curves	21


Review Exercise	23
12 Addition Formulae	24
13 The Circle	25
14 The Wave Function	26
14.1 The Wave Function using $k \cos(x - a)$	27
14.2 Other Forms of the The Wave Function	29
14.3 Maximum and Minimum Values using the Wave Function	30
14.4 Solving Equations using the Wave Function	31
14.5 Multiple Angles and Different Variables	32
Wave Function Review Exercise	33
15 Logs and Exponentials	34
15.1 Graphs of Exponential Functions	35
15.2 Logarithms	37
16 Further Calculus	38
Answers	39
Challenge Problems	40

Introduction

Intro

Introduction and Overview

In mathematics, *integration* can be thought of as the process of calculating some kind of “*sum*”, such as the total area under a curve, by breaking it down into sums of increasingly small parts. This technique was known about and used thousands of years ago in both Ancient Greece and other parts of the world.



In the 17th century both Gottfried Wilhelm Leibniz and Isaac Newton independently discovered the *fundamental theorem of calculus*, showing that *integration* can be performed using *antidifferentiation*, which is the ‘reverse’ of differentiation. Now, *integration* is used to refer to both the idea of calculating such a ‘sum’, and to the process of antidifferentiation.



This chapter will first introduce the use of integration to calculate an *antiderivative*, before moving on to using this concept to calculating area enclosed using curves.

Chapter Contents

- 11.1 Indefinite Integrals
- 11.2 Differential Equations
- 11.3 Definite Integrals
- 11.4 The Area Under a Curve
- 11.5 The Area Between Two Curves

Integration skills covered in this chapter will be extended upon in the later chapter of **Further Calculus**.

11.1 Indefinite Integrals

The *derivative* of the function $f(x) = x^3 - 4x^2 - 7x + 3$ is obtained by *multiplying by the power* and then *reducing the power by one* for each term.

$$\begin{aligned}\text{If } f(x) &= x^3 - 4x^2 + 3 \\ \text{Then } f'(x) &= 3x^2 - 8x\end{aligned}$$

Integration can be considered the *inverse* of differentiation, with the aim of taking the *derivative* $f'(x)$ and determining the *original function* $f(x)$. However, given only the derivative $f'(x) = 3x^2 - 8x$ it would not be possible to know any constant term the original function $f(x)$ contained. Finding the *indefinite integral* of a function requires the inclusion of a **Constant of Integration, C** :

$$\begin{aligned}\text{If } f'(x) &= 3x^2 - 8x \\ \text{Then } f(x) &= x^3 - 4x^2 + C\end{aligned}$$

Indefinite Integrals: "The (indefinite) integral of x^n with respect to x ..."

$$\int (x^n) dx = \frac{x^{n+1}}{n+1} + C$$

In other words, **increase the power by 1** then **divide by the new power**.

The *integral sign*, \int , should always appear accompanied by dx , for a function in x .

Note: $\int (f(x) + g(x)) dx = \int f(x) dx + \int g(x) dx$

Example 11.1.1

Find $\int (6x^2 + 8x) dx$.

$$\begin{aligned}\int (6x^2 + 8x) dx \\ &= \frac{6x^3}{3} + \frac{8x^2}{2} + C && \leftarrow \text{Constant of integration} \\ &= 2x^3 + 4x^2 + C && \leftarrow \text{Simplify}\end{aligned}$$

As with differentiation, *preparation for integration* may be needed.

Example 11.1.2

Find $\int \left(4\sqrt{x} - \frac{3}{x^2}\right) dx, x > 0$.

$$\begin{aligned}
 & \int \left(4\sqrt{x} - \frac{3}{x^2}\right) dx \\
 &= \int \left(4x^{\frac{1}{2}} - 3x^{-2}\right) dx && \leftarrow \text{Preparing to integrate} \\
 &= \frac{4x^{\frac{3}{2}}}{\frac{3}{2}} - \frac{3x^{-1}}{-1} + C && \leftarrow \text{Integrate: } \frac{1}{2} + 1 = \frac{1}{2} + \frac{2}{2} = \frac{3}{2} \\
 &= \frac{8}{3}x^{\frac{3}{2}} + 3x^{-1} + C && \leftarrow \text{Simplify: } 4 \div \frac{3}{2} = \frac{4}{1} \times \frac{2}{3} = \frac{8}{3}
 \end{aligned}$$

The *derivative* of a *linear* term is a *constant*, so the integral of a *constant* is *linear*.

Example 11.1.3

Find $\int ((2x - 1)(x + 3)) dx$.

$$\begin{aligned}
 & \int ((2x - 1)(x + 3)) dx \\
 &= \int (2x^2 - 5x - 3) dx && \leftarrow \text{Preparing to integrate} \\
 &= \frac{2x^3}{3} - \frac{5x^2}{2} - 3x + C && \leftarrow \text{Integral of } -3 \text{ is } -3x
 \end{aligned}$$

Integration *with respect to variables other than x* should not use dx , and the notation used should match the variable of the function instead.

Example 11.1.4

Find $\int (3t^2 - 5) dt$.

$$\begin{aligned}
 & \int (3t^2 - 5) dt \\
 &= \frac{3t^3}{3} - 5t + C \\
 &= t^3 - 5t + C
 \end{aligned}$$

11.2 Differential Equations

11.3 Definite Integrals

The indefinite integral of the function $f(x)$ can be notated as $F(x)$:

$$\int f(x) \, dx = F(x)$$

The *definite integral* of $f(x)$ from a to b is the difference between $F(b)$ and $F(a)$:

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

Together, they form a core part of the **Fundamental Theorem of Calculus**:

Fundamental Theorem of Calculus:

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

where

$$F(x) = \int f(x) \, dx$$

Since any *constant of integration* within $F(x)$ will cancel through subtraction ($C - C$), it is *not included* when calculating a *definite integral*.


Example 11.3.1

Find $\int_1^2 (3x^2 + 6x - 2) \, dx$.

$$\begin{aligned} & \int_1^2 (3x^2 + 6x - 2) \, dx \\ &= \left[\frac{3x^3}{3} + \frac{6x^2}{2} - 2x \right]_1^2 && \leftarrow \text{Integrate} \\ &= [x^3 + 3x^2 - 2x]_1^2 && \leftarrow \text{Simplify} \\ &= ((2)^3 + 3(2)^2 - 2(2)) - ((1)^3 + 3(1)^2 - 2(1)) && \leftarrow \text{Substitute} \\ &= 16 - 4 \\ &= 12 && \leftarrow \text{Evaluate} \end{aligned}$$


11.4 The Area Under a Curve

The area shaded below, with each "bar" having width 1 unit, is given by the *sum* $1 + 2 + 3 + 1 + 2$.



An integral, \int , can be seen as "sum", but for a continuous function. Given a function $f(x)$, the area enclosed between a section of the curve $y = f(x)$ and the x -axis can be calculated using a definite integral.

Area Under a Curve:




$$\text{Area} = \int_a^b f(x) \, dx = F(b) - F(a)$$

The values of a and b can be referred to as the *bounds* of the integration.


Example 11.4.1

Part of the graph of $y = 5 + 2x - x^2$ is shown. Calculate the shaded area.



$$\begin{aligned} & \int_{-1}^2 (5 + 2x - x^2) dx && \leftarrow \text{Area} \\ &= \left[5x + x^2 - \frac{x^3}{3} \right]_{-1}^2 && \leftarrow \text{Integrate} \\ &= \left(5(2) + (2)^2 - \frac{(2)^3}{3} \right) - \left(5(-1) + (-1)^2 - \frac{(-1)^3}{3} \right) && \leftarrow \text{Substitute} \\ &= \frac{34}{3} - \left(-\frac{11}{3} \right) && \leftarrow \text{Evaluate} \\ &= 15 \text{ square units} \end{aligned}$$

Where the area between a curve and the x -axis lies *under* the x -axis, the definite integral will be negative.




To avoid "*positive*" and "*negative*" areas cancelling each other out, such sections must be calculated as separate integrals, and their *absolute values* added.

$$\text{Area} = A_1 + A_2$$

Example 11.4.2

Part of the graph of $y = x^3 - 10x^2 + 27x - 18$ is shown. Calculate the shaded area.



$$\int_1^3 (x^3 - 10x^2 + 27x - 18) dx$$

$$= \left[\frac{x^4}{4} - \frac{10x^3}{3} + \frac{27x^2}{2} - 18x \right]_1^3$$

$$= \left(\frac{(3)^4}{4} - \frac{10(3)^3}{3} + \frac{27(3)^2}{2} - 18(3) \right)$$

$$- \left(\frac{(1)^4}{4} - \frac{10(1)^3}{3} + \frac{27(1)^2}{2} - 18(1) \right)$$

$$= -\frac{9}{4} - \left(-\frac{91}{12} \right)$$

$$= \frac{16}{3}$$

$$\int_3^6 (x^3 - 10x^2 + 27x - 18) dx$$

$$= \left[\frac{x^4}{4} - \frac{10x^3}{3} + \frac{27x^2}{2} - 18x \right]_3^6$$

$$= \left(\frac{(6)^4}{4} - \frac{10(6)^3}{3} + \frac{27(6)^2}{2} - 18(6) \right)$$

$$- \left(\frac{(3)^4}{4} - \frac{10(3)^3}{3} + \frac{27(3)^2}{2} - 18(3) \right)$$


$$= -18 - \left(-\frac{9}{4} \right)$$

$$= -\frac{63}{4}$$

$$\therefore \text{Area} = \frac{16}{3} + \frac{63}{4} = \frac{253}{12} \text{ square units}$$


11.5 The Area Between Two Curves

The area **between** two curves can be calculated using the subtraction of one definite integral from another:



Note that $\int_a^b f(x) dx - \int_a^b g(x) dx = \int_a^b (f(x) - g(x)) dx$, leading to the following formula:


Area Between Two Curves:



$$\text{Area} = \int_a^b (f(x) - g(x)) dx \quad \text{or} \quad \text{Area} = \int_a^b (\text{Upper} - \text{Lower}) dx$$

Example 11.5.1

Part of the graphs of $y = -3x$ and $y = 4 - x^2$ are shown. Calculate the shaded area.



$$\int_{-1}^4 (4 - x^2 - (-3x)) dx$$

← Upper – Lower

$$\int_{-1}^4 (4 - x^2 + 3x) dx$$

← Simplify

$$= \left[4x - \frac{x^3}{3} + \frac{3x^2}{2} \right]_{-1}^4$$

← Integrate

$$= \left(4(4) - \frac{(4)^3}{3} + \frac{3(4)^2}{2} \right) - \left(4(-1) - \frac{(-1)^3}{3} + \frac{3(-1)^2}{2} \right)$$

← Substitute

$$= \frac{56}{3} - \left(-\frac{13}{6} \right)$$

← Evaluate


$$= \frac{125}{6} \text{ square units}$$

Review Exercise


In this chapter, the sums and differences of *equal-angled* trigonometric operations will be explored.

e.g. $3 \sin x^\circ + 4 \cos x^\circ$

The graphs of basic trigonometric functions such as $y = 3 \sin x^\circ$ and $y = 4 \cos x^\circ$ should be familiar:



The graph of $y = 3 \sin x^\circ + 4 \cos x^\circ$ is the same as that of $y = 5 \cos (x - 36.9)^\circ$.



Determining that $3 \sin x^\circ + 4 \cos x^\circ = 5 \cos (x - 36.9)^\circ$ allows the maximum values and minimum values of $3 \sin x^\circ + 4 \cos x^\circ$ to be determined and the values of x which produce them, and allow equations such as $3 \sin x^\circ + 4 \cos x^\circ = 2$ to be solved.

Any trigonometric expression $k_1 \sin x \pm k_2 \cos x$ can be written as either $k \sin (x \pm a)$ or $k \cos (x \pm a)$.

This chapter will cover how to do this and explore the various applications of this property.

14.1 The Wave Function using $k \cos(x - a)$

Whilst any of $k \sin(x \pm a)$ or $k \cos(x \pm a)$ may be used in general, any Higher exam question is likely to specify a particular form to use. The simplest is often $k \cos(x - a)$.

To express $3 \sin x^\circ + 4 \cos x^\circ$ in the form $k \cos(x - a)^\circ$, two key *trigonometric identities* are needed:

$$\sin^2 x + \cos^2 x = 1 \quad \text{and} \quad \tan x = \frac{\sin x}{\cos x}$$

First, $k \cos(x - a)^\circ$ can be expanded using the formula $\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$:

$$k \cos(x - a)^\circ = k \cos x^\circ \cos a^\circ + k \sin x^\circ \sin a^\circ$$

It is required that this expansion is equal to the original expression:

$$3 \sin x^\circ + 4 \cos x^\circ = k \cos x^\circ \cos a^\circ + k \sin x^\circ \sin a^\circ$$

Equating the coefficients of $\sin x^\circ$ and $\cos x^\circ$ gives a set of simultaneous equations in k and a :

$$\begin{aligned} k \sin a^\circ &= 3 && \leftarrow \text{Coefficients of } \sin x^\circ \\ k \cos a^\circ &= 4 && \leftarrow \text{Coefficients of } \cos x^\circ \end{aligned}$$

Squaring both sides of each equation and adding allows k to be determined using $\sin^2 x + \cos^2 x = 1$:

$$\begin{aligned} k^2 \sin^2 a^\circ + k^2 \cos^2 a^\circ &= 3^2 + 4^2 \\ k^2 (\sin^2 x^\circ + \cos^2 x^\circ) &= 3^2 + 4^2 \\ k^2 (1) &= 3^2 + 4^2 \\ k &= \sqrt{3^2 + 4^2} \\ k &= \sqrt{25} \\ k &= 5 \end{aligned}$$

Dividing $k \sin a^\circ$ by $k \cos a^\circ$ allows a to be calculated, noting $k \sin a^\circ$ and $k \cos a^\circ$ are both positive:

$$\begin{aligned} \frac{k \sin a^\circ}{k \cos a^\circ} &= \frac{3}{4} \\ \tan a^\circ &= \frac{3}{4} \\ a_{\text{acute}}^\circ &= \tan^{-1} \left(\frac{3}{4} \right) = 36.9^\circ \\ a^\circ &= 36.9^\circ \end{aligned} \quad \begin{array}{c|c} \checkmark \text{S} & \checkmark \text{A}^\circ \checkmark \\ \hline \text{T} & \checkmark \text{C} \checkmark \end{array}$$

Hence $3 \sin x^\circ + 4 \cos x^\circ = 5 \cos(x - 36.9)^\circ$. Here it has been assumed that $k > 0$ and $0 < a < 360$.

Some abbreviations of the working shown on the previous page are routinely permitted in the Higher exam. The following example demonstrates an appropriate level of detail in its solution.

Example

Express $3 \cos x - \sin x$ in the form $k \cos(x - a)$ where $k > 0$ and $0 < a < 2\pi$.

$$3 \cos x - \sin x = k \cos(x - a)$$

$$3 \cos x - \sin x = k \cos x \cos a + k \sin x \sin a \quad \leftarrow \text{Expand } k \cos(x - a)$$

$$k \sin a = -1$$

\leftarrow Equate $\sin x$ coefficients

$$k \cos a = 3$$

\leftarrow Equate $\cos x$ coefficients

$$k = \sqrt{(-1)^2 + 3^2}$$

\leftarrow Calculate k

$$= \sqrt{10}$$

$$\tan a = \frac{-1}{3}$$

\leftarrow Find $\tan a$

$$\text{acute}^\circ = \tan^{-1}\left(\frac{1}{3}\right) = 18.4^\circ$$

\leftarrow Find acute angle first

S	A✓
✓T	C✓✓
$360^\circ - \text{acute}^\circ$	

\leftarrow Negative $\sin a$: quadrants T and C

\leftarrow Positive $\cos a$: quadrants A and C

$$a^\circ = 341.6^\circ$$

\leftarrow Using $360^\circ - 18.4^\circ$ (Double-ticked quadrant)

$$a = 5.96$$

\leftarrow Convert to radians: $341.6 \times \frac{\pi}{180}$

$$3 \cos x - \sin x = \sqrt{10} \cos(x - 5.96)$$

\leftarrow State solution

Exercise 14.1

1. Express each of the following in the form $k \cos(x - a)^\circ$, where $k > 0$ and $0 < a < 360$.

(a) $5 \cos x^\circ + 12 \sin x^\circ$

(b) $4 \sin x^\circ + 5 \cos x^\circ$

(c) $6 \cos x^\circ + \sin x^\circ$

2. Express each of the following in the form $k \cos(x - a)$, where $k > 0$ and $0 < a < 2\pi$.

(a) $8 \sin x - 6 \cos x$

(b) $\cos x - 3 \sin x$

(c) $-2 \sin x - \cos x$

3. Express each of the following in the form $k \cos(x - a)^\circ$, where $k > 0$ and $0 < a < 360$.

(a) $\sin x^\circ + \cos x^\circ$

(b) $\cos x^\circ - \sqrt{3} \sin x^\circ$

(c) $-\sin x^\circ - \sqrt{3} \cos x^\circ$

14.2 Other Forms of the The Wave Function

As well as $k \cos(x - a)$, three other forms may be used, shown below including their expansions:

$$k \cos(x + a) = k \cos x \cos a - k \sin x \sin a$$

$$k \sin(x + a) = k \sin x \cos a + k \cos x \sin a$$

$$k \sin(x - a) = k \sin x \cos a - k \cos x \sin a$$

Example

Express $12 \cos x^\circ - 5 \sin x^\circ$ in the form $k \sin(x - a)^\circ$ where $k > 0$ and $0 < a < 360$.

$$12 \cos x^\circ - 5 \sin x^\circ = k \sin(x - a)^\circ$$

$$12 \cos x^\circ - 5 \sin x^\circ = k \sin x^\circ \cos a^\circ - k \cos x^\circ \sin a^\circ \quad \leftarrow \text{Expand } k \sin(x - a)^\circ$$

$$-k \sin a^\circ = 12 \implies k \sin a^\circ = -12 \quad \leftarrow \text{Equate } \sin x^\circ \text{ coefficients}$$

$$k \cos a^\circ = -5 \quad \leftarrow \text{Equate } \cos x^\circ \text{ coefficients}$$

$$k = \sqrt{(-12)^2 + (-5)^2} \quad \leftarrow \text{Calculate } k$$

$$= 13$$

$$\tan a^\circ = \frac{-12}{-5} = \frac{12}{5} \quad \leftarrow \text{Find } \tan a^\circ$$

$$\text{acute}^\circ = \tan^{-1}\left(\frac{12}{5}\right) = 67.4^\circ \quad \leftarrow \text{Find acute angle first}$$

$$180^\circ + \text{acute}^\circ$$

\leftarrow Negative $\sin a^\circ$: quadrants T and C

\leftarrow Negative $\cos a^\circ$: quadrants S and T

$$a^\circ = 247.4^\circ \quad \leftarrow \text{Using } 180^\circ + 67.4^\circ \text{ (Double-ticked)}$$

$$12 \cos x^\circ - 5 \sin x^\circ = 13 \sin(x - 247.4)^\circ \quad \leftarrow \text{State solution}$$

Exercise 14.2

1. Express each of the following in the form $k \sin(x + a)^\circ$, where $k > 0$ and $0 < a < 360$.

(a) $3 \cos x^\circ + 2 \sin x^\circ$

(b) $4 \sin x^\circ + 3 \cos x^\circ$

(c) $7 \cos x^\circ - \sin x^\circ$

2. Express each of the following in the form $k \cos(x + a)$, where $k > 0$ and $0 < a < 2\pi$.

(a) $3 \sin x + 8 \cos x$

(b) $-3 \cos x - 4 \sin x$

(c) $-6 \sin x + 2 \cos x$

3. Express each of the following in the form $k \sin(x - a)$, where $k > 0$ and $0 < a < 2\pi$.



(a) $\sqrt{3} \sin x + \cos x$

(b) $\cos x - \sin x$

(c) $2\sqrt{3} \sin x - 2 \cos x$

14.3 Maximum and Minimum Values using the Wave Function

The graphs of $y = \sin x^\circ$ and $y = \cos x^\circ$, including their turning points, should already be familiar:




Graphs of the form $k \sin(x \pm a)$ and $k \cos(x \pm a)$ have maximum/minimum values of $\pm k$, and their *horizontal translation* is described by $\pm a$ following the rules covered in the Graph Transformations chapter.

Example

Given that $3 \sin x^\circ + 4 \cos x^\circ$ can be expressed as $5 \cos(x - 36.9)^\circ$, state the minimum value of $f(x) = 3 \sin x^\circ + 4 \cos x^\circ$ and the value of x at which it occurs for $0 < x < 360$.

Sketch the graph of $y = 5 \cos x^\circ$ translated 36.9° to the right:



Use the graph to state the answer:

Hence the minimum value of $f(x)$ is -5 , which occurs when $x = 216.9$.

Exercise 14.3

- Sketch each for $0 < x < 360$, showing the coordinates of any roots and turning points:
 - $7 \cos(x - 10)^\circ$
 - $3 \sin(x - 20)^\circ$
 - $\sqrt{5} \cos(x + 40)^\circ$
- Find the maximum value of each and the value(s) of x for which they occur for $0 < x < 360$:
 - $8 \sin(x - 50)^\circ$
 - $\sqrt{2} \cos(x + 27)^\circ$
 - $7 \sin(x + 42.3)^\circ$
- Find the minimum value of each and the value(s) of x for which they occur for $0 < x < 2\pi$:
 - $3 \cos\left(x - \frac{\pi}{6}\right)$
 - $5 \sin\left(x + \frac{\pi}{3}\right) + 2$
 - $-12 \sin\left(x + \frac{\pi}{4}\right)$
- Express $6 \sin x^\circ - 7 \cos x^\circ$ in the form $k \sin(x - a)^\circ$ where $k > 0$ and $0 < a < 360$.
 - Hence state the coordinates of the turning points of $12 \sin x^\circ - 14 \cos x^\circ$ for $0 < x < 360$.

14.4 Solving Equations using the Wave Function

Solving an equation like $3 \sin x^\circ + 4 \cos x^\circ = 2$ can be approached by using the wave function to rewrite it as $5 \cos(x - 36.9)^\circ = 2$, before solving it in the manner covered in Chapter 6.

Example

It can be shown that $5 \sin x^\circ - 2 \cos x^\circ$ can be expressed as $\sqrt{29} \sin(x - 21.8)^\circ$.
Hence, solve the equation $5 \sin x^\circ - 2 \cos x^\circ = 4$ where $0 < x < 360$.

$$5 \sin x^\circ - 2 \cos x^\circ = 4$$

$$\sqrt{29} \sin(x - 21.8)^\circ = 4$$

← Substitute the wave function form

$$\sin(x - 21.8)^\circ = \frac{4}{\sqrt{29}}$$

← Rearrange to $\sin(\dots) = \dots$

$$a = \sin^{-1}\left(\frac{4}{\sqrt{29}}\right) = 48.0^\circ$$

← Calculate acute angle

$$\begin{array}{c|c} 180^\circ - a^\circ & a^\circ \\ \hline \checkmark S & A \checkmark \\ T & C \end{array}$$

← Positive $\sin a^\circ$: quadrants A and S

$$x - 21.8^\circ = 48.0^\circ, 180^\circ - 48.0^\circ$$

← Apply ticked quadrants

$$x - 21.8^\circ = 48.0^\circ, 132.0^\circ$$

$$x^\circ = 69.8^\circ, 153.8^\circ$$

← Add 21.8° to both sides

Note that any solutions outwith the domain (often $0 < x < 360$) should have 360° added or subtracted to bring it back within the domain, where possible.

Exercise 14.4

1. Solve each equation for $0 < x < 360$: 

$$(a) 7 \sin(x - 18)^\circ = 4 \quad (b) 3 \cos(x + 34.1)^\circ = -2 \quad (c) \sqrt{5} \sin(x - 106)^\circ + 1 = 0$$

2. Given $6 \sin x^\circ - 8 \cos x^\circ = 10 \sin(x - 53.1)^\circ$, solve $6 \sin x^\circ - 8 \cos x^\circ = 5$ where $0 < x < 360$.


3. (a) Express $\sqrt{3} \sin x^\circ + \cos x^\circ$ in the form $k \sin(x - a)^\circ$ where $k > 0$ and $0 < a < 360$.

(b) Hence solve the equation $\sqrt{3} \sin x^\circ + \cos x^\circ = 1$ where $0 < x < 360$.

4. Solve each equation for $0 < x < 2\pi$: 

$$(a) 4 \sin(x + 0.31) + 2 = 1 \quad (b) 9 \cos(x + 1.24) = 5 \quad (c) 2\sqrt{3} \sin(x - 0.82) = \sqrt{5}$$

5. (a) Express $3 \cos x + 2$ in the form $k \cos(x + a)$ where $k > 0$ and $0 < a < 2\pi$. 

(b) Hence solve the equation $2 + 6 \sin x + 4 \cos x = 5$ where $0 < x < 2\pi$. 

14.5 Multiple Angles and Different Variables

The techniques covered in this chapter can be applied to trigonometric expressions beyond those containing only $\sin x$ and $\cos x$; they work for any sum or difference of *equal-angled* trigonometric operations.

e.g. $3 \sin t^\circ + 4 \cos t^\circ$ or $2 \sin 2x - 5 \cos 2x$

Example

Express $5 \cos 2t^\circ - 3 \sin 2t^\circ$ in the form $k \sin(2t + a)^\circ$ where $k > 0$ and $0 < a < 360$.

$$5 \cos 2t^\circ - 3 \sin 2t^\circ = k \sin(2t + a)^\circ$$

$$\underline{5 \cos 2t^\circ} - \underline{3 \sin 2t^\circ} = k \underline{\sin 2t^\circ} \cos a^\circ + k \underline{\cos 2t^\circ} \sin a^\circ \quad \leftarrow \text{Expand } k \sin(2t + a)^\circ$$

$$k \sin a^\circ = 5$$

\leftarrow Equate $\sin 2t^\circ$ coefficients

$$k \cos a^\circ = -3$$

\leftarrow Equate $\cos 2t^\circ$ coefficients

$$k = \sqrt{(5)^2 + (-3)^2}$$

$$= \sqrt{34}$$

\leftarrow Calculate k

$$\tan a^\circ = \frac{5}{-3} = -\frac{5}{3}$$

\leftarrow Find $\tan a^\circ$

$$\text{acute}^\circ = \tan^{-1}\left(\frac{5}{3}\right) = 59.0^\circ$$

\leftarrow Find acute angle first

$$\begin{array}{c|c} 180^\circ - \text{acute}^\circ & \\ \hline \checkmark \checkmark \text{S} & \text{A} \checkmark \\ \checkmark \text{T} & \text{C} \end{array}$$

\leftarrow Positive $\sin a^\circ$: quadrants A and S

\leftarrow Negative $\cos a^\circ$: quadrants S and T

$$a^\circ = 121.0^\circ$$

\leftarrow Using $180^\circ - 59.0^\circ$ (Double-ticked)

$$5 \cos 2t^\circ - 3 \sin 2t^\circ = \sqrt{34} \sin(2t + 121.0)^\circ$$


\leftarrow State solution

Exercise 14.5

- Express $4 \cos t^\circ - 3 \sin t^\circ$ in the form $k \sin(t - a)^\circ$, where $k > 0$ and $0 < a < 360$.
- Express $2 \sin 2x - \cos 2x$ in the form $k \cos(2x - a)$ where $k > 0$ and $0 < a < 2\pi$.
- (a) Express $12 \cos t^\circ + 5 \sin t^\circ$ in the form $k \sin(t + a)^\circ$, where $k > 0$ and $0 < a < 360$.
 (b) Hence state:
 - The maximum value of the function $f(x) = 12 \cos t^\circ + 5 \sin t^\circ$, $0 < t < 360$.
 - The value(s) of t for which it occurs.
- (a) Express $\sin 2x - \sqrt{3} \cos 2x$ in the form $k \cos(2x - a)$ where $k > 0$, $0 < a < 2\pi$.
 (b) Hence solve $\sin 2x - \sqrt{3} \cos 2x - 1 = 0$, $0 < x < 2\pi$.
 (c) Sketch $y = -\sin 2x + \sqrt{3} \cos 2x - 1 = 0$ for $0 \leq x \leq 2\pi$.

Wave Function Review Exercise

- Express $8 \sin x^\circ + 7 \cos x^\circ$ in the form $k \sin(x - a)^\circ$ where $k > 0$ and $0 < a < 360$.
- Express $\sqrt{5} \sin x + \cos x$ in the form $k \cos(x - a)$ where $k > 0$ and $0 < a < 2\pi$.
- Express $\sqrt{3} \cos t^\circ - \sin t^\circ$ in the form $k \sin(t + a)^\circ$ where $k > 0$ and $0 < a < 360$.
- Part of the graphs of $y = 3 \cos x^\circ - 5 \sin x^\circ$ and $y = -4$ are shown in the diagram below:



Points P and Q are points of intersection.

- Express $y = 3 \cos x^\circ - 5 \sin x^\circ$ in the form $k \cos(x + a)^\circ$ where $k > 0$ and $0 < a < 360$.
 - Hence determine the coordinates of P and Q.
- Express $2 \sin x^\circ - 4 \cos x^\circ$ in the form $k \sin(x - a)^\circ$ where $k > 0$ and $0 < x < 360$.
 - Hence sketch the graph of $y = 2 \sin x^\circ - 4 \cos x^\circ$ for $0 < x < 360$.
 - Express $\cos x + \sqrt{3} \sin x$ in the form $k \cos(x - a)$ where $k > 0$ and $0 < a < 2\pi$.
 - Hence sketch the graph of $y = 2 \cos x + 2\sqrt{3} \sin x$ for $0 < x < 2\pi$.
 - Express $6 \cos t - 3 \sin t$ in the form $k \cos(t + a)$ where $k > 0$ and $0 < a < 2\pi$.
 - Hence solve $2 \cos t - \sin t + 2 = 1$ where $0 < t < 2\pi$.
 - Express $\sin 2x^\circ - \cos 2x^\circ$ in the form $k \sin(2x - a)$ where $k > 0$ and $0 < a < 360$.
 - Hence solve the equation $\sin 2x^\circ = \cos 2x^\circ$ where $0 < x < 360$.

Each part of an operation of the form a^n can be described using the following terminology:



Power functions take the form $f(x) = x^n$, with x as the base and a constant power, n .

e.g. $f(x) = x^3$

Exponential functions take the form $f(x) = a^x$, with x as the exponent and a constant base, $a > 0, a \neq 1$.

e.g. $f(x) = 3^x$

An example of an exponential function in everyday life is that of something *appreciating* by a percentage of its value, such as an antique vase of value £4000 increasing by 20% each year. Its value after 0 years, 1 years, 2 years, and so on, can be calculated as follows:

$4000 \times 1.2^0 = 4000$	← After 0 years
$4000 \times 1.2^1 = 4800$	← After 1 year
$4000 \times 1.2^2 = 5760$	← After 2 years
$4000 \times 1.2^3 = 6912$	← After 3 years
$4000 \times 1.2^4 = 8294.40$	← After 4 years

The function to describe its value V after x years is given by: $V(x) = 4000 \times 1.2^x$

One advantage of defining this function is the ability to calculate the value at times other than after whole years. For example, the value after three and a half years can be calculated as:


$$V(3.5) = 4000 \times 1.2^{3.5} = 7571.72$$

This chapter will introduce a range of skills required when working with exponential functions.

15.1 Graphs of Exponential Functions

Where $a > 1$:


$y = a^x$ is *strictly increasing* on $x \in \mathbb{R}$:



This describes **exponential growth**.

Where $0 < a < 1$:

$y = a^x$ is *strictly decreasing* on $x \in \mathbb{R}$:



This describes **exponential decay**.


Since $a^0 = 1$, any graph of the form $y = a^x$ will pass through the point $(0, 1)$ for all $a \neq 0$.

Since $a^1 = a$, any graph of the form $y = a^x$ will pass through the point $(1, a)$ for all a .

Determining the equation of the graph of an exponential function can typically be achieved using substitution or consideration of graph transformations, along with knowledge of points $(0, 1)$ and $(1, a)$.

Example

The graph of $y = a^x + b$ is shown below.



$$2 = a^0 + b$$

← Substitute $(0, 2)$

$$2 = 1 + b$$

$$1 = b$$

← Solve to obtain b

$$5 = a^2 + 1$$

← Substitute $(2, 5)$ and $b = 1$

$$4 = a^2$$

$$2 = a$$

← Solve to obtain a


$$y = 2^x + 1$$

← State equation


Exercise 15.1

1. Find the equation of each exponential graphs using the form given.


(a) $y = a^x$




(b) $y = a^x$




(c) $y = a^x$




(d) $y = a^x$




(e) $y = a^x$




(f) $y = a^x$



(g) $y = a^x$



(h) $y = a^x$



15.2 Logarithms

ANSWERS

CHALLENGE PROBLEMS

The following problems **do not** represent the kind of question expected to feature in a Higher Mathematics exam, either in the way they are presented or the level of difficulty. Instead, they aim to encourage a flexible approach towards problem-solving and an understanding that the skills covered in the course have applications beyond those featured in any typical exam. *Some questions may be solveable without using the skills covered in this chapter, and some questions may be unrelated to this chapter.*