

Signaling to the Actin Cytoskeleton During Cell Morphogenesis and Patterning

Alison Sinclair, Mike Schenkel, and Jaideep Mathur

Abstract The plant is a supracellular organism whose cells are locked in position through shared walls but maintain apoplastic and symplastic connectivity. Their fixed position places each cell into a unique niche within the organism. Consequently, every environmental cue is perceived slightly differently by each cell. The response of each plant cell varies accordingly. Thus, plant growth and development reflect a progression of accommodative arrangements reached between constituent cells. In recent years the actin cytoskeleton, through its direct involvement in subcellular compartmentation, organelle and vesicle trafficking, and structural reinforcement, has emerged as a key player during accommodative growth and development. Here, using salient actin-cytoskeleton-associated cellular phenotypes, we elaborate upon the molecular-cell biological machinery involved in organizing the actin cytoskeleton during cell shape development in plants.

1 Introduction

A series of division, expansion, differentiation, and cell death events are involved in the fabrication of a multicellular plant. These events occur within a genetically defined framework in response to environmental cues and result in the precise patterns that provide each plant species with a unique, recognizable identity. However, if the millions of cells that make up a plant are considered individually, it becomes apparent that each wall-encased plant cell occupies a unique niche with

A. Sinclair

Laboratory of Plant Development & Interactions, Department of Molecular and Cellular Biology, CBS, University of Guelph, 50 Stone Road, Guelph, ON N1G 2W1, Canada

M. Schenkel

Laboratory of Plant Development & Interactions, Department of Molecular & Cellular Biology, CBS, University of Guelph, 50 Stone Road, Guelph, ON N1G 2W1, Canada

J. Mathur (✉)

Department of Molecular and Cellular Biology, 4463 Science Complex, University of Guelph, 488 Gordon Street, Guelph, ON N1G 2W1, Canada
e-mail: jmathur@uoguelph.ca

respect to the environment. Owing to its fixed location within the plant, each cell perceives and responds to environmental cues in a manner slightly different from that of its neighbors. Further, each cell has only limited response options for contributing to the genetically determined architectural master plan: it can divide to add more adjustable units (cells); it can expand in a polar fashion to fill out spaces; or it can die and thus vacate space to help reorganize the overall shape. In addition, as components of a supracellular organism, individual cells can differentiate into specialized forms for fulfilling specific functions. The creation of a precise form by numerous plant cells thus reflects an innate ability to accommodate cell structure and function to that of neighboring cells. How is this cooperation and coordination of developmental activities achieved by wall-locked plant cells?

A number of signaling cascades, including $\text{Ca}^{2+}/\text{H}^+$, reactive oxygen species (ROS), heterotrimeric G-proteins, phospholipases, phosphoinositides, mitogen-activated protein kinases, cyclic AMP, and GTPases, are recognized in plants (Hirt 2000; Hong-Bo et al. 2008; Sumimoto 2008; Shao et al. 2008; Chen 2008; Cheung and Wu 2008; Kost 2008). In recent years the actin cytoskeleton has emerged as a central responder to many of these signaling channels (Brenner et al. 2006; Nibau et al. 2006; Kost 2008). The actin cytoskeleton also constitutes the signaling hub through which numerous subcellular activities are regulated. These include subcellular compartmentation, organelle and vesicle motility, as well as structural reinforcement in co-operation with microtubules. With each new study it becomes increasingly apparent that almost every actin-associated activity has an impact on cell growth and morphogenesis. Interestingly, seemingly different cues appear to produce similar responses from the actin cytoskeleton. Moreover, as we follow molecular cascades elicited in response to diverse signals, we become aware of the considerable gaps in our ability to directly link environmental cues to the far-downstream behavior of the actin cytoskeleton and the subsequent elicitation of a developmental response. While one can easily envision a kind of inverted pyramid of subcellular events whereby diverse cues get parsed into increasingly narrower response categories, the molecular-cell biological mechanisms behind this thinking applied to the actin cytoskeleton are only beginning to be understood.

As detailed and excellent reviews on different aspects of signaling to the actin cytoskeleton continue being published, here we have adopted a broad definition for the term “signaling.” The actin cytoskeleton has been placed at a hub: on one hand we explore the putative molecular-cell biological events that organize actin within the plant cell, and on the other we link information on actin organization to the morphogenesis and development of model cell types in different organisms and assorted mutants of *Arabidopsis thaliana*.

2 Actin Organization in Plant Cells

Our understanding of signaling to the actin cytoskeleton first requires a characterization of the typical actin cytoskeleton organization in plants. The actin gene family in plants is diverse and built up of multiple isoforms with unique patterns of regulation (Meagher et al. 2000). The actin cytoskeleton, comprising actin monomers (G-actin)

and polymerized filamentous actin (F-actin) can change rapidly through cue-mediated depolymerization and repolymerization cycles. In living cells, F-actin strands twist around each other to produce helical arrangements of microfilaments possessing an approximate diameter of 7 nm. Immunocytochemical methods have informed us a lot about the actin organization in different cell types in plants (Vitha et al. 2000; Staiger and Schliwa 1987). More recent advances in live imaging of F-actin arrays have greatly shaped our present views on the dynamic actin cytoskeleton, notably, the introduction of fluorescent phalloidin into plant cells through micro injection (Schmit and Lambert 1990), microabrasion (Ramachandran et al. 2000), or after enzymatic digestion and freeze shattering (Wasteney et al. 1997) has been pursued successfully. Synthetic fluorescent fusion proteins created using the filamentous actin binding domain of the Talin gene (Kost et al. 1998) and the actin binding domain 2 of the Fimbrin 1 gene (Sheahan et al. 2004; Wang et al. 2008) are the most used live probes for visualizing F-actin arrays in plants. G-actin pools in plant cells appear predominantly cytosolic and show up as diffuse fluorescence. Three morphologically distinguishable, but readily interchangeable states characterize the polymerized actin. These are seen as very fine meshworks comprising actin filaments, thick F-actin bundles, and loose F-actin strands (Staiger et al. 2000; Mathur et al. 1999; Wang et al. 2008). Since at the light-microscopy level it is not possible to resolve individual filaments of actin, the states probably reflect different levels of actin bundling. Actin patches as described for yeasts (Young et al. 2004) have not been observed in plant cells; however, brightly fluorescent regional aggregates of F-actin can be clearly observed in a variety of plant cells. Whether these represent true actin patches comprising interwoven actin filaments or reflect subcellular compartmentation is still an open question. Actin cages around large organelles have been described (Kandasamy and Meagher 1999). In addition to the above, short-lived arrays of actin filaments that contribute to the preprophase band and to stages of phragmoplast formation have been observed in dividing cells (Kakimoto and Shibaoka 1987; Sano et al. 2005).

Occasionally, 3–5- μm -long, brightly fluorescent bundles of F-actin might be encountered in cells that are experiencing subcellular stress or have been physically injured. All arrays are readily convertible to each other and reflect the dynamic nature of the actin cytoskeleton.

2.1 Different Actin Regulators Organize the Actin Cytoskeleton

A large number of actin genes and actin-interacting proteins have been identified and cloned from plants (Staiger et al. 2000; Meagher and Fehcheimer 2003; Ketelaar et al. 2004; Staiger and Blanchoin 2006; Grunt et al. 2008). Since many actin interactors are conserved between eukaryotes, *in vitro* studies have been very instructive in ascribing roles for different proteins involved in actin cytoskeleton organization (Ayscough 1998). The direct visualization of actin arrays in plant cells through fluorescent imaging (Miller et al. 1996; Kost et al. 1998; Mathur et al. 1999; Blancaflor, 2000; Kovar et al. 2000b; Chen et al. 2002) has reinforced many of the *in vitro* observations.

As expected, cells expressing fluorescent protein fused to actin monomer interacting proteins such as profilin (Kovar et al. 2000a) exhibit a diffuse fluorescence (Ramachandran et al. 2000; Fu et al. 2002). However, overexpression of fluorescent protein fusions with different actin depolymerizing factors, also actin monomer interacting proteins, labels filamentous actin (Dong et al. 2001; Chen et al. 2002). Labeling experiments have confirmed that the proteins fimbrin (Kovar et al. 2000b) and villin (Klahre et al. 2000; Tominaga et al. 2000) bind to and bundle F-actin at different strengths. The ARP2/3 complex, an efficient enhancer of actin polymerization, is known to initiate side branches on actin filaments, creating a dendritic actin network (Mullins et al. 1998; Amann and Pollard 2001). Though immunological detection of ARP3 (one of the large subunits of the ARP2/3 complex) has been carried out (Van Gestel et al. 2003), and despite the demonstrated presence of its seven subunits (Mathur 2005a; Szymanski 2005), the complex has not been isolated from plants thus far. Inferences on ARP2/3 complex activity in relation to the actin cytoskeleton in plants are thus largely based on observations made in heterologous systems and actin organization observed in *arp2/3* mutants.

More than 20 formins are recognized in plants, and as profilin-binding poly(L-proline)-containing proteins these have important roles in cytoskeletal organization (Cheung and Wu 2004; Deeks et al. 2005; Michelot et al. 2005; Ingouff et al. 2005). A major regulator of actin nucleation and polymerization through its control of the pool of unpolymerized ATP-actin is the adenyl cyclase associated protein (AtCAP1; Chaudhry et al. 2007; Deeks et al. 2007). A gelsolin-related actin binding protein PrABP80 has been identified in poppy pollen tubes (Huang et al. 2004). Biochemical assays indicate that the protein can nucleate actin polymerization from monomers, block the assembly of profilin-actin complex onto actin filament ends, enhance profilin-mediated actin depolymerization, and also sever actin filaments (Huang et al. 2004).

Since the number of proteins that can interact and affect actin organization continues to increase, clearly the key to actin's pivotal role as a subcellular responder lies in the way it is organized through the combinatorial and cumulative activities of its various regulatory proteins.

2.2 Local Ionic Interactions Modulate Actin Regulators

The plant cell functions by maintaining a basal level of ionic homeostasis. Any extrinsic or intrinsic cues that perturb this homeostasis are dealt with by the release or sequestration of relevant ionic species (Foreman et al. 2003; Shabala et al. 2006). This provides the cell with a fine-tuned mechanism for sensing environmental cues as well as responding to them. Since proteins affecting actin nucleation, polymerization, and subsequent organizational activities respond rapidly to alterations in their ionic environment (Day et al. 2002), they are direct mediators in the stimulus perception-response elicitation machinery. Many actin-regulating proteins are themselves regulated by ionic conditions in their environment (Bretscher and Weber

1980; Matsudaira and Burgess 1982; Harris and Weeds 1983; Kovar et al. 2000a; Huang et al. 2003; Drøbak et al. 2004; Gu et al. 2004; Jones et al. 2007). The best studied ionic pools playing a role in signaling to actin include Ca^{2+} (Hetherington and Brownlee 2004; Bothwell and Ng 2005), NADPH oxidases (Navazio et al. 2000; Kwak et al. 2003; Foreman et al. 2003) and ROS, such as hydrogen peroxide (H_2O_2), superoxide anion ($\text{O}_2^{\cdot-}$), hydroxyl ions (OH) and nitric oxide (NO) (Miller et al. 2008; Mori and Schroeder 2004). Low, basal levels of cytosolic free calcium ($[\text{Ca}^{2+}]_{\text{cyt}}$) are sustained by Ca^{2+} -ATPases, and through the activity of a $\text{Ca}^{2+}/\text{H}^+$ antiporter that removes Ca^{2+} from the cytosol. Elevation of $[\text{Ca}^{2+}]_{\text{cyt}}$ during signaling occurs through the opening of different classes of plasma membrane, vacuolar membrane, or endoplasmic reticulum localized Ca^{2+} -permeable ion channels. Whereas Ca^{2+} - and H^+ -mediated signaling in plants is usually achieved through storage and release mechanisms, ROS signaling is usually controlled through rapid production and scavenging.

A major approach adopted for linking subcellular ionic distribution to actin organization and cell growth characteristics has been to observe and quantify ionic gradients and site-specific localizations (Felle and Hepler 1997). This approach has been successfully applied to single-celled, tip-growing cells such as pollen tubes (Cárdenas et al. 2008; Kroeger et al. 2008) and root hairs (Carol and Dolan 2006; Monshausen et al. 2007). In *Arabidopsis*, ROS stimulate the entry of Ca^{2+} into the cell through a calcium plasma membrane hyperpolarization-activated Ca^{2+} channel (Foreman et al. 2003). Recently, Takeda et al. (2008) have elucidated an elegant mechanism involving local positive feedback, where ROS derived through an RHD2 NADPH oxidase stimulates a Ca^{2+} influx into the cytoplasm. Though the targeted actin regulators were not identified in this study, the regulatory loop is clearly involved in cell shape determination of root hair cells. Similar interdependent local loops between ROS and Ca^{2+} gradients have been uncovered during polarized growth of *Fucus serratus* zygotes (Coelho et al. 2008).

Details pertaining to integrated calcium-mediated signaling (see “Integrated Calcium Signaling in Plants” by Narendra Tuteja, this volume), the role of apoplastic and cell-wall mediation in signaling events (see “Signaling and Cell Walls” by Ewelina Rodakowska et al., this volume), and signaling during plant-pathogen interactions utilizing localized ROS production (see “Signaling via Plant Peroxidases” by Claude Penel and Christophe Dunand, this volume) are not further elaborated upon here.

2.3 *Rho-like GTPases of Plants: Master Switches Responding to Environmental Cues*

In tracing back many of the ionic alterations brought about in response to environmental cues, a conserved set of proteins constantly comes up. Rho-like GTPases of plants (ROPs) localize to the cell membrane (Bischoff et al. 2000; Molendijk et al. 2001; Cheung et al. 2003) and are involved in multiple signal transduction events in plants. In *Arabidopsis*, 11 ROPs have been described and extensive reviews have

been published on signaling mediated by ROPs and their regulators (Zheng and Yang 2000; Gu et al. 2004; Xu and Scheres 2005; Nibau et al. 2006; Kost 2008). Direct links between ROP activity and Ca^{2+} spiking (Li et al. 1999; Jones et al. 2002), and ROP mediation in localized ROS activation (Jones et al. 2007) have been demonstrated. ROP-mediated signaling is also one of the best elucidated chains of molecular links directed to the actin cytoskeleton in plants (Fu et al. 2005; Fig. 1).

Specific ROPs and their regulators (RICs, ROP-GEFs/ROP-GDI) have an effect on the actin cytoskeleton but these proteins, high up in the response hierarchy of




Fig. 1 A generalized depiction of events involved in signaling to the actin cytoskeleton in plants. The leaf epidermal pavement cells drawn here are considered to represent all cell types since the vast majority of actin organizations and actin-mediated processes can be observed in them. Moreover, whereas many different signal cascades might operate in different plant cells, presently pavement cells provide one of the best characterized systems for dissecting signals to the actin cytoskeleton (Fu et al. 2005). An inverted pyramid of events suggests the gradual parsing of signals through the molecular apparatus to produce an impact on the actin cytoskeleton. Diverse environmental cues perceived at the cell boundary (cell wall and plasma membrane) initiate feedback loops of Rho-like GTPases of plants (ROP)-GTPases and ionic activity, which subsequently affect several proteins that bind directly to actin (such as actin monomer interacting, nucleating, bundling, and severing proteins) and bring about changes in its organization. Local actin organization has implications for subcellular motility, exocytosis-endocytosis, cytomembrane organization, and structural reinforcement. The cumulative results of these activities lead to cell differentiation, morphogenesis, and functional diversification during plant development. *Solid lines and arrows* depict experimentally demonstrated links between protein complexes, whereas *broken lines* imply putative interactions

the plant cell, do not cause their effects through direct binding to actin. Their effects impact multiple proteins and processes, influencing both the microtubule and the actin cytoskeleton (Fu et al. 2005). From a phenotype-based viewpoint an interference with ROPs and associated machinery does not produce cell forms that can be solely linked to actin defects (e.g., root hair phenotypes depicted in Fig. 2).




Fig. 2 Salient cellular phenotypes in model cell types of *Arabidopsis thaliana*. The left side shows cell morphology displayed by wild-type plants, while the phenotypes drawn in **bold lines** are exhibited by different mutants. In general, cells with an aberrant actin cytoskeleton are smaller in size. Phenotypes displayed by ROPs and their regulators do not strictly follow the pattern obtained by an interference with the actin cytoskeleton only. These phenotypes suggest a strong disturbance in the microtubule cytoskeleton too and could reflect actin-microtubule interactivity during cell shape development

However, neither are the ROP-associated phenotypes indicative of microtubule cytoskeletal defects only. Live imaging of the actin and microtubule cytoskeletons in *Arabidopsis* plants possessing aberrant activity of different ROPs and their regulators (e.g., ROP-GDI) suggests that they exhibit both actin and microtubule defects (Mathur, unpublished data).

3 Developmental Consequences of Different Actin Organizations

Whereas it has been possible to draw correlations between a small region of growth and its ionic environment, such as the one provided by tip-growing cells (Gu et al. 2005; Ovecka et al. 2008; Kroeger et al. 2008), establishing similar correlations has been difficult for diffuse-growing cells that make up the bulk of higher plants. The very nature of diffuse growth involves widespread, possibly transient signaling events that preclude precise localization of ionic concentrations and gradients. Therefore, a reverse approach has been adopted for understanding molecular mechanisms involved in unraveling the signaling to and the roles of actin in diffuse-growing cells. This approach (Mathur et al. 1999; Szymanski et al. 1999; Fu et al. 2002) has largely been based on visualizing defects in actin organization in particular cell types and then tracing them to a molecular lesion involved in creating similar aberrant cellular activity. In this approach, the use of inhibitors such as cytochalasin D, latrunculin, phalloidin, and jasplakinolide has been very helpful in uncovering specific actin-associated cellular phenotypes (Mathur et al. 1999; Baluska et al. 2001; Collings et al. 2006; Foissner and Wasteneys 2007) and hinting at possible processes involved. The actin defective phenotype based approach, when combined with live imaging, has been successful in uncovering new protein classes and complexes involved in mediating signals to the actin cytoskeleton (Mathur et al. 1999, 2003a, b).

The occurrence of the actin organizations discussed above has been demonstrated in different model cell types in plants (representative images in publications: root hair – Ryan et al. 2001; pollen tube – Lovy-Wheeler et al. 2005; hypocotyl cells – Mathur et al. 2003b; pavement cells – Fu et al. 2002, 2005; trichomes – Mathur et al. 1999). Section 3.1 associates actin cytoskeleton organization to a developmental consequence in different cell types.

3.1 *Actin Patch Indicative of Cell Polarization*

Polar growth is an established facet of plant development. However, the analysis of early events leading to the establishment of polarity in the asymmetric environment inhabited by land plants is a difficult task. Observations on the early development of fucoid zygotes have been informative in highlighting a role for the actin cytoskeleton during polarity establishment in a nearly apolar cell (Kropf et al. 1989; Bisgrove 2007). Upon release from the mother frond, and following their fertilization, spherical fucoid zygotes rely on several vectorial inputs such as

blue light, gravity, electrical-ionic impulses, and temperature and osmotic gradients for subsequent development (Brownlee and Bouget 1998). A distinct actin patch has been associated with zygote polarization and creation of the rhizoidal pole (Kropf et al. 1989). A cortical localization of the actin modulator ARP2/3 appears to be intimately involved in actin patch formation (Hable and Kropf 2005). Actin depolymerizing substances disrupt cell polarization and subsequent development (Hable et al. 2003). In later stages of zonation of the zygote, asymmetric localization of ARP2 and actin results in distinct actin arrays (Hable and Kropf 2005).

3.2 Actin Organization Indicative of Focused Growth

Pollen tubes and root hairs have been the most extensively studied tip-growing cells and numerous publications have described the actin cytoskeleton in these cells and reviewed the implications of the polar F-actin observed in them (Geitmann and Emons 2000; Ryan et al. 2001; Ringli et al. 2005; Geitmann 2006). In recent years the moss *Physcomitrella patens* has emerged as an equally informative model system for understanding polarized tip growth.

In *Physcomitrella*, single apical cells of moss filaments undergo polarized tip growth by perceiving and aligning themselves to gravity and light inputs (Cove 2005). Several recent studies point to specialized roles for actin dynamics in modulating these events, prompting the hypothesis that the ARP2/3 complex is pivotal in translating environmental cues by acting as a downstream target for signals regulating polar tip growth (Perroud and Quatrano 2006). Subsequently, there has been a strong focus on evaluating the roles of both the ARP2/3 complex and its upstream regulators Scar/WAVE complexes in polar growth of the apical cell of *P. patens* filament cells. When the *arpc4* subunit of the ARP2/3 complex is deleted, the resulting null mutant is viable. However, while tissue morphogenesis in the mutant proceeds normally from filamentous growth to leafy shoot formation, it displays inefficient perception of polarized white light and a marked reduction in apical cell growth (Perroud and Quatrano 2006). Additionally, RNA interference lines knocking down expression of another ARP2/3 member, ARPC1, result in a similar, albeit more severe, phenotype as displayed by the *arpc4* mutant. Overexpression of the respective ARP2/3 members (ARPC4, ARPC1) rescues the mutant phenotypes. Moreover, insertion of a yellow fluorescent protein–ARPC4 localizes ARPC4 exclusively to the site of polarized extension in the apical cell tip (Perroud and Quatrano 2006). Moreover, the deletion of BRK1, an upstream regulator of the ARP2/3 complex (Frank and Smith 2002), results in a mutant that though viable and responsive to polarized white light shows significant reduction in apical cell growth (Perroud and Quatrano 2008). BRK1 also localizes to the apical tip. A role for BRK1 in ARP2/3 recruitment is supported as the BRK1 mutant of *physcomitrella* is unable to localize both ARPC4 and AGP (a cell wall proteoglycan) to the apical cell tip (Perroud and Quatrano 2008).

3.3 *Actin Organization Indicative of Diffuse Growth*

The majority of plant cells grow by diffuse growth. Different model cell types have been informative about the actin-associated molecular–cellular machinery involved in spreading growth occurring over large regions of the cell. These include trichomes (epidermal hair) that grow outward from the aerial epidermis, interconnected jigsaw puzzle shaped pavement cells, and axially elongating epidermal cells of the hypocotyl, cotyledon, and leaf petiole. Each of these epidermal cell types offers a different surface to the environment and their growth responses differ accordingly.

In *Arabidopsis*, trichomes are single-celled and display a rapid expansion phase in the later stage of their development (Mathur 2006b). As the trichome cell grows out of the epidermal plane, unencumbered by growth of neighboring cells, a large portion of it is exposed directly to the environment. In contrast, both pavement and hypocotyl/petiole cells achieve accommodative growth by maintaining connectivity with their neighbors. All three epidermal cell types exhibit long F-actin cables that stretch from end to end, as well as a fine cortical F-actin mesh. In addition, in both trichomes and pavement cells, areas of indentation (valleys) display bundled or aggregated F-actin. In general, the presence of bundled F-actin in plant cells has come to be associated with regions of low expansion, whereas the presence of a fine cortical F-actin meshwork suggests regions of increased expansion.

3.3.1 *Diffuse Growth in Trichomes*

The use of actin inhibitors resulted in random shape alterations in *Arabidopsis* trichomes (Mathur et al. 1999; Szymanski et al. 1999). The drug-induced trichome distortion is phenocopied by mutations in several *Arabidopsis* genes. Molecular characterization of these genes has identified the ARP2/3 complex, an efficient modulator of actin polymerization that initiates the formation of a fine dendritic F-actin mesh (reviewed in Mathur 2005a, b; Szymanski 2005; Hussey et al. 2006), and its upstream regulators, starting from ROPs, ROP-GEFs such as SPIKE1 (Qiu et al. 2002; Basu et al. 2008; Zhang et al. 2008), and including different SCAR/WAVE, HSPC300-like/BRICK proteins (Frank and Smith 2002; Frank et al. 2004; Zhang et al. 2005; Basu et al. 2005; Le et al. 2006; Djakovic et al. 2006), NAP125-like and PIR121-like proteins (reviewed by Szymanski 2005; Smith and Oppenheimer 2005). An elegant interaction map of SCAR genes and their interactions with ARP2/3 proteins obtained through mutant studies, yeast two-hybrid, and bimolecular fluorescence complementation proteins experiments has been presented (Uhrig et al. 2007). In addition to the ROP-SPIKE-SCAR/WAVE-ARP2/3 pathway mentioned above, several other signaling pathways to the actin cytoskeleton might be activated through the involvement of novel ROP-GEFs, termed “plant-specific ROP nucleotide exchanger (PRONE), which are exclusively active towards members of the ROP subfamily (Berken et al 2005).

3.3.2 Interdigitating Growth of Pavement Cells

The final “jigsaw puzzle” shape of epidermal pavement cells arises from multiple local projections (lobes) of a polygonal initial. During expansion, the lobes of one pavement cell fit into the indentions of its neighbors to produce an epidermal surface with an interdigitating pattern. Fu et al. (2005) revealed a ROP-GTPase signaling network underlying the creation of these lobes and indentions by pavement cells. Local activation of a ROP (AtROP2) activates ROP-interactive CRIB-motif-containing protein 4 (RIC4) to enhance actin dynamics and promote localized growth. However, ROP2 activity leads to the inactivation of another target, RIC1, that localizes to cortical microtubules and promotes their ordering into parallel arrays. RIC1-dependent microtubule organization not only inhibits cell outgrowth locally but also suppresses ROP2 activation in the indentation zone. Thus, while the ROP2–RIC4 interaction promotes cell outgrowth, the ROP2–RIC1 interaction restrains outgrowth. Such coordinated activity in epidermal cells creates the interdigitations between adjacent pavement cells.

3.3.3 Accommodative Growth of Cylindrical Cells

The growth of cells of the seedling hypocotyl and petioles relies on efficient actin-based communication between cell ends for producing elongation along the longitudinal axis (Baluska et al. 2003). These cells have a diffuse F-actin mesh but display cytoplasmic aggregation at the cell ends. Mutants in different subunits of the ARP2/3 complex, hypocotyl, and petiole cells often display increased F-actin aggregation at their ends and consequently lose cell-cell connectivity (Mathur et al. 2003a, b). The reduced actin polymerization activity in elongating cylindrical cells observed in the arp2/3 mutants has implications for cell-cell communication and polar transport of growth regulators. It thus offers a novel system to understand the initiation and maintenance of multicellularity in higher plants.

4 The Diverse Roles of the Actin Cytoskeleton in Plants

An appraisal of cellular phenotypes associated with different actin cytoskeleton organizations, the nature of actin regulatory proteins that can create the organizations, and the diversity of environmental cues that trigger subcellular activity suggests that actin can be involved in many different kinds of roles inside the plant cell. Some of these roles are discussed in Sect. 4.1.

4.1 *Actin as a Barrier*

The regional accumulation of actin correlates with increased aggregation and decreased motility of small organelles such as Golgi bodies, peroxisomes, and mitochondria

(Mathur et al. 2003a, b). Upon extrapolation, the observations could suggest that F-actin acts as an intracellular barrier for subcellular trafficking of vesicles. The notion was tested by looking for areas of actin aggregation versus those with fine cortical actin meshwork in different cell types (Mathur et al. 2003a). Regions with actin aggregation invariably displayed lesser growth as compared with regions with a fine meshwork of F-actin (Mathur et al. 2003a; Fu et al. 2005). According to the “actin barrier” concept, a fine F-actin meshwork resulting from increased actin dynamics mediated by the ARP2/3 complex and formins would allow and perhaps even aid the rapid motility of organelles and exocytotic vesicles (Mathur 2004). This leads to a promotion of local growth. In contrast, a dense web comprising F-actin aggregates and bundles in an intracellular locality might hinder organelle motility and vesicular trafficking and could thereby display a lower rate of regional growth. The idea of F-actin behaving as a flexible barrier is appealing when considering the phenotypes described above for pavement and hypocotyl cells impaired in their actin organization. The role of cortical F-actin as a “fence” has also been suggested through three-dimensional electron-tomographic reconstructions of the membrane cytoskeleton in normal rat kidney fibroblast cells (Morone et al. 2006).

4.2 *Actin as a Propulsive Force*

Asymmetric actin polymerization mediated through the ARP2/3 complex has been implicated in the rocketing motility of enteropathogenic organisms such as *Listeria monocytogenes* and *Shigella* species, as well as vesicles and organelles such as endosomes, phagosomes (Higgs and Pollard 2001; Gouin et al. 2005), and mitochondria (Boldogh et al. 2001). The polymerization of actin is also believed to act as the driving force at the leading edge of motile animal cells (Goley and Welch 2006). In plants, direct evidence for actin polymerization acting as a propulsive force for subcellular structures or as a protrusive force for cell membranes has not been observed. However, given the role of actin in vesicle movement during auxin efflux and translocation (Dhonukshe et al. 2007), the rapid movement of endocytic vesicles (Voigt et al. 2005), and cytoplasmic streaming, the possibility of actin polymerization mediated force generation within the plant cell cannot be totally ruled out (Mathur 2005b).

4.3 *Actin as an Interactive Support System*

The actin cytoskeleton interacts with microtubules. During root elongation in *Arabidopsis* the hypersensitivity to cytoskeletal antagonists has been used to demonstrate microtubule-microfilament cross-talk (Blancaflor 2000; Collings et al. 2006). One of the molecular mechanisms controlling these interactions has been traced to the antagonistic activities of ROP-RIC proteins (Fu et al. 2005) and elegantly described for the morphogenesis of pavement cells of *Arabidopsis*.

Interestingly, observations in trichome cells suggest that F-actin aggregates coincide with regional clusters of cytoplasmic microtubules (Saedler et al. 2004). The observation suggests that microtubules, with their own set of interactors and regulators, might act to confine F-actin-based regional expansion in a cell (Mathur 2004, 2006a). The role of F-actin in sculpting the cortical microtubule cytoskeleton has also been suggested (Schwab et al. 2003).

4.4 *Actin as a Track*

Observations on different organelles in plant cells have shown that their motility is actin-based (Ovecka et al. 2008) and involves interactions with myosin motors (Samaj et al. 2000; Reisen and Hanson 2007; Sparkes et al. 2008). In this situation, actin filaments act as subcellular tracks for the motor molecules. How cargo specificity and the choice of a specific actin track is decided by the different myosins in plant cells still remains an enigma. There also remains the question of interactivity and switching between myosin motors and microtubule-based motors such as kinesins.

5 Conclusions

While reviews on the minutiae of signaling to the actin cytoskeleton will continue being published, one of the long-term goals of the field is to uncover the mechanisms whereby this global perceiver of, and responder to, myriad environmental cues is able to create local influences within a cell. As described in this chapter, the key to actin's responsiveness lies in its interactivity with numerous regulatory proteins as well as with other components and compartments of cell. Signaling to actin thus becomes an exercise in attempting to understand and sum up everything that goes on inside a living cell. Realistically, we are far from the goal, but given the tremendous scope of actin interactions, the increased availability of novel mutants in model plant species, and the emergence of new tools and techniques for live imaging (Dhanoa et al. 2006), the prospects of unraveling the tangled web around actin looks very promising.

Acknowledgements We gratefully acknowledge funding by the Natural Sciences and Engineering Research Council of Canada (NSERC), Canada Foundation for Innovation (CFI), The Ministry of Research and Innovation (MRI), Ontario, and the Keefer Trust Fund, Guelph.

References

- Amann KJ, Pollard TD (2001) The Arp2/3 complex nucleates actin filament branches from the sides of pre-existing filaments. *Nat Cell Biol* 3:306–310
- Ayscough KR (1998) In vivo functions of actin-binding proteins. *Curr Opin Cell Biol* 10:102–111
- Baluska F, Jasik J, Edelmann HG, Salajová T, Volkmann D (2001) Latrunculin B-induced plant dwarfism: Plant cell elongation is F-actin-dependent. *Dev Biol* 231:113–124

- Baluska F, Wojtaszek P, Volkmann D, Barlow P (2003) The architecture of polarized cell growth: the unique status of elongating plant cells. *Bioessays* 25:569–576
- Basu D, Le J, El-Essal Sel-D, Huang S, Zhang C, Mallery EL, Koliantz G, Staiger CJ, Szymanski DB (2005) DISTORTED3/SCAR2 is a putative arabidopsis WAVE complex subunit that activates the Arp2/3 complex and is required for epidermal morphogenesis. *Plant Cell* 17:502–524
- Basu D, Le J, Zakharova T, Mallery EL, Szymanski DB (2008) A SPIKE1 signaling complex controls actin-dependent cell morphogenesis through the heteromeric WAVE and ARP2/3 complexes. *Proc Natl Acad Sci U S A* 105:4044–4049
- Berken A, Thomas C, Wittinghofer A (2005) A new family of RhoGEFs activates the Rop molecular switch in plants. *Nature* 436:1176–1178
- Bischoff F, Vahlkamp L, Molendijk A, Palme K (2000) Localization of AtROP4 and AtROP6 and interaction with the guanine nucleotide dissociation inhibitor AtRhoGDI1 from Arabidopsis. *Plant Mol Biol* 42:515–530
- Bisgrove SR (2007) Cytoskeleton and early development in Fucoid algae. *J Integr Plant Biol* 49:1192–1198
- Blancaflor EB (2000) Cortical actin filaments potentially interact with cortical microtubules in regulating polarity of cell expansion in primary roots of maize (*Zea mays* L.). *J Plant Growth Regul* 19:406–410
- Boldogh IR, Yang HC, Nowakowski WD, Karmon SL, Hays LG, Yates JR3rd, Pon LA (2001) ARP2/3 complex and actin dynamics are required for actin-based mitochondrial motility in yeast. *Proc Natl Acad Sci U S A* 98:3162–3167
- Bothwell JH, Ng CK (2005) The evolution of Ca²⁺ signalling in photosynthetic eukaryotes. *New Phytol* 166:21–38
- Brenner ED, Stahlberg R, Mancuso S, Vivanco J, Baluska F, Van Volkenburgh E (2006) Plant neurobiology: an integrated view of plant signaling. *Trends Plant Sci* 11:413–419
- Bretscher A, Weber K (1980) Villin is a major protein of the microvillus cytoskeleton which binds both G and F actin in a calcium-dependent manner. *Cell* 20:839–847
- Brownlee C, Bouget FY (1998) Polarity determination in *Fucus*: from zygote to multicellular embryo. *Semin Cell Dev Biol* 9:179–185
- Cárdenas L, Lovy-Wheeler A, Kunkel JG, Hepler PK (2008) Pollen tube growth oscillations and intracellular calcium levels are reversibly modulated by actin polymerization. *Plant Physiol* 146:1611–1621
- Carol RJ, Dolan L (2006) The role of reactive oxygen species in cell growth: lessons from root hairs. *J Exp Bot* 57:1829–1834
- Chaudhry F, Guérin C, von Witsch M, Blanchoin L, Staiger CJ (2007) Identification of Arabidopsis cyclase-associated protein 1 as the first nucleotide exchange factor for plant actin. *Mol Biol Cell* 18:3002–3014
- Chen JG (2008) Heterotrimeric G-proteins in plant development. *Front Biosci* 13:3321–3333
- Chen CY, Wong EI, Vidali L, Estavillo A, Hepler PK, Wu H, Cheung AY (2002) The regulation of actin organization by actin depolymerizing factor in elongating pollen tubes. *Plant Cell* 14:2175–2190
- Cheung AY, Wu HM (2004) Over-expression of an Arabidopsis formin stimulates supernumerary actin cable formation from pollen tube cell membrane. *Plant Cell* 16:257–269
- Cheung AY, Wu HM (2008) Structural and signaling networks for the polar cell growth machinery in pollen tubes. *Annu Rev Plant Biol* 59:547–572
- Cheung AY, Chen CY-, Tao L-, Andreyeva T, Twell D, Wu H- (2003) Regulation of pollen tube growth by Rac-like GTPases. *J Exp Bot* 54:73–81
- Coelho SM, Brownlee C, Bothwell JH (2008) A tip-high, Ca²⁺-interdependent, reactive oxygen species gradient is associated with polarized growth in *Fucus serratus* zygotes. *Planta* 227:1037–1046
- Collings DA, Lill AW, Himmelsbach R, Wasteneys GO (2006) Hypersensitivity to cytoskeletal antagonists demonstrates microtubule-microfilament cross-talk in the control of root elongation in *Arabidopsis thaliana*. *New Phytol* 170:275–290
- Cove D (2005) The moss *Physcomitrella patens*. *Annu Rev Genet* 39:339–358

- Day IS, Reddy VS, Shad Ali G, Reddy AS (2002) Analysis of EF-hand-containing proteins in Arabidopsis. *Genome Biol* 3(10): 1–24
- Deeks MJ, Cvrcková F, Machesky LM, Mikitová V, Ketelaar T, Zársky V, Davies B, Hussey PJ (2005) Arabidopsis group Ie formins localize to specific cell membrane domains, interact with actin-binding proteins and cause defects in cell expansion upon aberrant expression. *New Phytol* 168:529–540
- Deeks MJ, Rodrigues C, Dimmock S, Ketelaar T, Maciver SK, Malhó R, Hussey PJ (2007) Arabidopsis CAP1 – a key regulator of actin organisation and development. *J Cell Sci* 120:2609–2618
- Dhanoa PK, Sinclair AM, Mullen RT, Mathur J (2006) Illuminating subcellular structures and dynamics in plants: a fluorescent protein toolbox. *Can J Bot* 84:515–522
- Dhonukshe P, Aniento F, Hwang I, Robinson DG, Mravec J, Stierhof YD, Friml J (2007) Clathrin-mediated constitutive endocytosis of PIN auxin efflux carriers in Arabidopsis. *Curr Biol* 17:520–527
- Djakovic S, Dyachok J, Burke M, Frank MJ, Smith LG (2006) BRICK1/HSPC300 functions with SCAR and the ARP2/3 complex to regulate epidermal cell shape in Arabidopsis. *Development* 133:1091–1100
- Dong CH, Kost B, Xia G, Chua NH (2001) Molecular identification and characterization of the Arabidopsis AtADF1, AtADFS and AtADF6 genes. *Plant Mol Biol* 45:517–527
- Drøbak BK, Franklin-Tong VE, Staiger CJ (2004) The role of the actin cytoskeleton in plant cell signaling. *New Phytol* 163:13–30
- Felle HH, Hepler PK (1997) The cytosolic Ca^{2+} concentration gradient of *Sinapis alba* root hairs as revealed by Ca^{2+} -selective microelectrode tests and Fura-dextran ratio imaging. *Plant Physiol* 114:39–45
- Foissner I, Wasteneys GO (2007) Wide-ranging effects of eight cytochalasins and latrunculin A and B on intracellular motility and actin filament reorganization in characean internodal cells. *Plant Cell Physiol* 48:585–597
- Foreman J, Demidchik V, Bothwell JH, Mylona P, Miedema H, Torres MA, Linstead P, Costa S, Brownlee C, Jones JD, Davies JM, Dolan L (2003) Reactive oxygen species produced by NADPH oxidase regulate plant cell growth. *Nature* 422: 442–446
- Frank MJ, Smith LG (2002) A small, novel protein highly conserved in plants and animals promotes the polarized growth and division of maize leaf epidermal cells. *Curr Biol* 12:849–853
- Frank M, Egile C, Dyachok J, Djakovic S, Nolasco M, Li R, Smith LG (2004) Activation of Arp2/3 complex-dependent actin polymerization by plant proteins distantly related to Scar/WAVE. *Proc Natl Acad Sci U S A* 101:16379–16384
- Fu Y, Li H, Yang Z (2002) The ROP2 GTPase controls the formation of cortical fine F-actin and the early phase of directional cell expansion during Arabidopsis organogenesis. *Plant Cell* 14:777–794
- Fu Y, Gu Y, Zheng Z, Wasteneys G, Yang Z (2005) Arabidopsis interdigitating cell growth requires two antagonistic pathways with opposing action on cell morphogenesis. *Cell* 120:687–700
- Geitmann A (2006) Plant and fungal cytomchanics: quantifying and modeling cellular architecture. *Can J Bot* 84:581–593
- Geitmann A, Emons AM (2000) The cytoskeleton in plant and fungal cell tip growth. *J Microsc* 198:218–245
- Gouin E, Welch MD, Cossart P (2005) Actin-based motility of intracellular pathogens. *Curr Opin Microbiol* 8:35–45
- Goley ED, Welch MD (2006) The ARP2/3 complex: an actin nucleator comes of age. *Nat Rev Mol Cell Biol* 7:713–726
- Grunt M, Zarsky V, Cvrckova F (2008) Roots of Angiosperm formins: the evolutionary history of plant FH2 domain-containing proteins. *BMC Evol Biol* 8:115
- Gu Y, Wang Z, Yang Z (2004) ROP/RAC GTPase: an old new master regulator for plant signaling. *Curr Opin Plant Biol* 7:527–536
- Gu Y, Fu Y, Dowd P, Li S, Vernoud V, Gilroy S, Yang Z (2005) A Rho family GTPase controls actin dynamics and tip growth via two counteracting downstream pathways in pollen tubes. *J Cell Biol*. 169:127–138

- Hable WE, Kropf DL (2005) The Arp2/3 complex nucleates actin arrays during zygote polarity establishment and growth. *Cell Motil Cytoskel* 61:9–20
- Hable WE, Miller NR, Kropf DL (2003) Polarity establishment requires dynamic actin in fucoid zygotes. *Protoplasma* 221:193–204
- Harris HE, Weeds AG (1983) Plasma actin depolymerizing factor has both calcium-dependent and calcium-independent effects on actin. *Biochemistry* 22:2728–2741
- Hetherington AM, Brownlee C (2004) The generation of Ca^{2+} signals in plants. *Annu Rev Plant Biol* 55:401–427
- Higgs HN, Pollard TD (2001) Regulation of actin filament network formation through ARP2/3 complex: activation by a diverse array of proteins. *Annu Rev Biochem* 70:649–676
- Hirt H (2000) MAP kinases in plant signal transduction. *Results Probl Cell Differ* 27:1–9
- Hong-Bo S, Li-Ye C, Ming-An S (2008) Calcium as a versatile plant signal transducer under soil water stress. *Bioessays* 30:634–641
- Huang S, Blanchoin L, Kovar DR, Staiger CJ (2003) Arabidopsis capping protein (AtCP) is a heterodimer that regulates assembly at the barbed ends of actin filaments. *J Biol Chem* 278:44832–44842
- Huang S, Blanchoin L, Chaudhry F, Franklin-Tong VE, Staiger CJ (2004) A gelsolin-like protein from *Papaver rhoeas* pollen (PrABP80) stimulates calcium-regulated severing and depolymerization of actin filaments. *J Biol Chem* 279:23364–23375
- Hussey PJ, Ketelaar T, Deeks MJ (2006) Control of the actin cytoskeleton in plant cell growth. *Annu Rev Plant Biol* 57:109–112
- Ingouff M, Fitz Gerald JN, Guérin C, Robert H, Sørensen MB, Van Damme D, Geelen D, Blanchoin L, Berger F (2005) Plant formin AtFH5 is an evolutionarily conserved actin nucleator involved in cytokinesis. *Nat Cell Biol* 7:374–380
- Jones MA, Shen JJ, Fu Y, Li H, Yang Z, Grierson CS (2002) The Arabidopsis Rop2 GTPase is a positive regulator of both root hair initiation and tip growth. *Plant Cell* 14:763–776
- Jones MA, Raymond MJ, Yang Z, Smirnov N (2007) NADPH oxidase-dependent reactive oxygen species formation required for root hair growth depends on ROP GTPase. *J Exp Bot* 58:1261–1270
- Kakimoto T, Shibata H (1987) Actin filaments and microtubules in the preprophase band and phragmoplast of tobacco cells. *Protoplasma* 140:151–156
- Kandasamy MK, Meagher RB (1999) Actin-organelle interaction: association with chloroplast in Arabidopsis leaf mesophyll cells. *Cell Motil Cytoskel* 44:110–118
- Ketelaar T, Allwood EG, Anthony R, Voigt B, Menzel D, Hussey PJ (2004) The actin-interacting protein AIP1 is essential for actin organization and plant development. *Curr Biol* 14:145–149
- Klahre U, Friederich E, Kost B, Louvard D, Chua NH (2000) Villin-like actin-binding proteins are expressed ubiquitously in Arabidopsis. *Plant Physiol* 122:35–48
- Kost B (2008) Spatial control of Rho (Rac-Rop) signaling in tip-growing plant cells. *Trends Cell Biol* 18:119–127
- Kost B, Spielhofer P, Chua NH (1998) A GFP-mouse talin fusion protein labels plant actin filaments *in vivo* and visualizes the actin cytoskeleton in growing pollen tubes. *Plant J* 16:393–401
- Kovar DR, Drobak BK, Staiger CJ (2000a) Maize profilin isoforms are functionally distinct. *Plant Cell* 12:583–598
- Kovar DR, Staiger CJ, Weaver EA, McCurdy DW (2000b) AtFim1 is an actin filament crosslinking protein from *Arabidopsis thaliana*. *Plant J* 24:625–636
- Kroeger JH, Geitmann A, Grant M (2008) Model for calcium dependent oscillatory growth in pollen tubes. *J Theor Biol* 253:363–374
- Kropf DL, Berge SK, Quatrano RS (1989) Actin localization during Fucus embryogenesis. *Plant Cell* 1:191–200
- Kwak JM, Mori IC, Pei ZM, Leonhardt N, Torres MA, Dangl JL, Bloom RE, Bodde S, Jones JD, Schroeder JI (2003) NADPH oxidase AtrbohD and AtrbohF genes function in ROS-dependent ABA signaling in Arabidopsis. *EMBO J* 22:2623–2633
- Le J, Mallery EL, Zhang C, Brankle S, Szymanski DB (2006) Arabidopsis BRICK1/HSPC300 is an essential WAVE-complex subunit that selectively stabilizes the Arp2/3 activator SCAR2. *Curr Biol* 16:895–901

- Li H, Lin Y, Heath RM, Zhu MX, Yang Z (1999) Control of pollen tube tip growth by a Rop GTPase-dependent pathway that leads to tip-localized calcium influx. *Plant Cell* 11:1731–1742
- Lovy-Wheeler A, Wilsen KL, Baskin TI, Hepler PK (2005) Enhanced fixation reveals the apical cortical fringe of actin filaments as a consistent feature of the pollen tube. *Planta* 221:95–104
- Mathur J (2004) Cell shape development in plants. *Trends Plant Sci* 9:583–590
- Mathur J (2005a) The ARP2/3 complex: giving plant cells a leading edge. *Bioessays* 27:377–387
- Mathur J (2005b) Conservation of boundary extension mechanisms between plants and animals. *J Cell Biol* 168:679–682
- Mathur J (2006a) Local interactions shape plant cells. *Curr Opin Cell Biol* 18:40–46
- Mathur J (2006b) Trichome cell morphogenesis in *Arabidopsis*: a continuum of cellular decisions. *Can J Bot* 84:581–593
- Mathur J, Spielhofer P, Kost B, Chua N (1999) The actin cytoskeleton is required to elaborate and maintain spatial patterning during trichome cell morphogenesis in *Arabidopsis thaliana*. *Development* 126:5559–5568
- Mathur J, Mathur N, Kirik V, Kernebeck B, Srinivas BP, Hülskamp M (2003a) *Arabidopsis* CROOKED encodes for the smallest subunit of the ARP2/3 complex and controls cell shape by region specific fine F-actin formation. *Development* 130:3137–3146
- Mathur J, Mathur N, Kernebeck B, Hülskamp M (2003b) Mutations in actin-related proteins 2 and 3 affect cell shape development in *Arabidopsis*. *Plant Cell* 15:1632–1645
- Matsudaira PT, Burgess DR (1982) Partial reconstruction of the microvillus core bundle: characterization of villin as a Ca^{2+} -dependent, actin-bundling/depolymerizing protein. *J Cell Biol* 92:648–656
- Meagher RB, Fechtmeier M (2003) The *Arabidopsis* cytoskeletal genome. In: Somerville CR, Meyerowitz EM (eds) *The Arabidopsis book*. American Society of Plant Biologists, Rockville
- Meagher RB, McKinney EC, Kandasamy MK (2000) The significance of diversity in the plant actin gene family. In: Staiger CJ et al (eds) *Actin: a dynamic framework for multiple plant cell functions*. Kluwer, Dordrecht, pp 3–27
- Michelot A, Guérin C, Huang S, Ingouff M, Richard S, Rodiuc N, Staiger CJ, Blanchoin L (2005) The formin homology 1 domain modulates the actin nucleation and bundling activity of *Arabidopsis* FORMIN1. *Plant Cell* 17:2296–2313
- Miller DD, Lancelle SA, Hepler PK (1996) Actin microfilaments do not form a dense meshwork in *Lilium longiflorum* pollen tube tips. *Protoplasma* 195:123–132
- Miller G, Shulaev V, Mittler R (2008) Reactive oxygen signaling and abiotic stress. *Physiol Plant* 253:481–489
- Molendijk AJ, Bischoff F, Rajendrakumar CS, Friml J, Braun M, Gilroy S, Palme K (2001) *Arabidopsis thaliana* Rop GTPases are localized to tips of root hairs and control polar growth. *EMBO J* 20:2779–2788
- Monshausen GB, Bibikova TN, Messerli MA, Shi C, Gilroy S (2007) Oscillations in extracellular pH and reactive oxygen species modulate tip growth of *Arabidopsis* root hairs. *Proc Natl Acad Sci U S A* 104:20996–21001
- Mori IC, Schroeder JI (2004) Reactive oxygen species activation of plant Ca^{2+} channels. A signaling mechanism in polar growth, hormone transduction, stress signaling, and hypothetically mechanotransduction. *Plant Physiol* 135:702–708
- Morone N, Fujiwara T, Murase K, Kasai RS, Ike H, Yuasa S, Usukura J, Kusumi A (2006) Three-dimensional reconstruction of the membrane skeleton at the plasma membrane interface by electron tomography. *J Cell Biol* 174:851–862
- Mullins RD, Heuser JA, Pollard TD (1998) The interaction of Arp2/3 complex with actin: Nucleation, high affinity pointed end capping, and formation of branching networks of filaments. *Proc Natl Acad Sci U S A* 95:6181–6186
- Navazio L, Bewell MA, Siddiqua A, Dickinson GD, Galione A, Sanders D (2000) Calcium release from the endoplasmic reticulum of higher plants elicited by the NADP metabolite nicotinic acid adenine dinucleotide phosphate. *Proc Natl Acad Sci U S A* 97:8693–8698
- Nibau C, Wu HM, Cheung AY (2006) RAC/ROP GTPases: ‘hubs’ for signal integration and diversification in plants. *Trends Plant Sci* 11:309–315

- Ovecka M, Baluska F, Lichtscheidl I (2008) Non-invasive microscopy of tip-growing root hairs as a tool for study of dynamic and cytoskeleton-based vesicle trafficking. *Cell Biol Int* 32:549–553
- Perroud PF, Quatrano RS (2006) The role of ARPC4 in tip growth and alignment of the polar axis in filaments of *Physcomitrella patens*. *Cell Motil Cytoskel* 63:162–171
- Perroud PF, Quatrano RS (2008) BRICK1 is required for apical cell growth in filaments of the moss *Physcomitrella patens* but not for gametophore morphology. *Plant Cell* 20:411–422
- Qiu JL, Jilk R, Marks MD, Szymanski DB (2002) The Arabidopsis SPIKE1 gene is required for normal cell shape control and tissue development. *Plant Cell* 14:101–118
- Ramachandran S, Christensen HE, Ishimaru Y, Dong CH, Chao-Ming W, Cleary AL, Chua NH (2000) Profilin plays a role in cell elongation, cell shape maintenance, and flowering in Arabidopsis. *Plant Physiol* 124:1637–1647
- Reisen D, Hanson MR (2007) Association of six YFP-myosin XI-tail fusions with mobile plant cell organelles. *BMC Plant Biol* 7:6
- Ringli C, Baumberger N, Keller B (2005) The Arabidopsis root hair mutants der2-der9 are affected at different stages of root hair development. *Plant Cell Physiol* 46:1046–1053
- Ryan E, Steer M, Dolan L (2001) Cell biology and genetics of root hair formation in *Arabidopsis thaliana*. *Protoplasma* 215:140–149
- Saedler R, Mathur N, Srinivas BP, Kernebeck B, Hülskamp M, Mathur J (2004) Actin control over microtubules suggested by DISTORTED2 encoding the Arabidopsis ARPC2 subunit homolog. *Plant Cell Physiol* 45:813–822
- Samaj J, Peters M, Volkmann D, Baluska F (2000) Effects of myosin ATPase inhibitor 2,3-butanedione 2-monoxime on distributions of myosins, F-actin, microtubules, and cortical endoplasmic reticulum in maize root apices. *Plant Cell Physiol* 41:571–582
- Sano T, Higaki T, Oda Y, Hayashi T, Hasezawa S (2005) Appearance of actin microfilament ‘twin peaks’ in mitosis and their function in cell plate formation, as visualized in tobacco BY-2 cells expressing GFP-fimbrin. *Plant J* 44:595–605
- Schmit AC, Lambert AM (1990) Microinjected fluorescent phalloidin *in vivo* reveals the F-Actin dynamics and assembly in higher plant mitotic cells. *Plant Cell* 2:129–138
- Schwab B, Mathur J, Saedler R, Schwarz H, Frey B, Scheidegger C, Hülskamp M (2003) Regulation of cell expansion by the DISTORTED genes in *Arabidopsis thaliana*: actin controls the spatial organization of microtubules. *Mol Genet Genom* 269:350–360
- Shabala S, Shabala L, Gradmann D, Chen Z, Newman I, Mancuso S (2006) Oscillations in plant membrane transport: model predictions, experimental validation, and physiological implications. *J Exp Bot* 57:171–184
- Shao HB, Chu LY, Shao MA, Jaleel CA, Hong-Mei M (2008) Higher plant antioxidants and redox signaling under environmental stresses. *C R Biol* 331:433–441
- Sheahan MB, Staiger CJ, Rose RJ, McCurdy DW (2004) A green fluorescent protein fusion to actin-binding domain 2 of Arabidopsis fimbrin highlights new features of a dynamic actin cytoskeleton in live plant cells. *Plant Physiol* 136:3968–3978
- Smith LG, Oppenheimer DG (2005) Spatial control of cell expansion by the plant cytoskeleton. *Annu Rev Cell Dev Biol* 21:271–295
- Sparkes IA, Teanby NA, Hawes C (2008) Truncated myosin XI tail fusions inhibit peroxisome, Golgi, and mitochondrial movement in tobacco leaf epidermal cells: a genetic tool for the next generation. *J Exp Bot* 59:2499–2512
- Staiger CJ, Blanchoin L (2006) Actin dynamics: old friends with new stories. *Curr Opin Plant Biol* 9:554–562
- Staiger CJ, Schliwa M (1987) Actin localization and function in higher plants. *Protoplasma* 141:1–12
- Staiger CJ, Baluska F, Volkmann D, Barlow PW (2000) Actin: a dynamic framework for multiple plant cell functions. Springer, Berlin
- Sumimoto H (2008) Structure, regulation and evolution of Nox-family NADPH oxidases that produce reactive oxygen species. *FEBS J* 275:3249–3277
- Szymanski DB (2005) Breaking the WAVE complex: the point of Arabidopsis trichomes. *Curr Opin Plant Biol* 8:103–1012

- Szymanski DB, Marks MD, Wick SM (1999) Organized F-actin is essential for normal trichome morphogenesis in *Arabidopsis*. *Plant Cell* 11:2331–2347
- Takeda S, Gapper C, Kaya H, Bell E, Kuchitsu K, Dolan L (2008) Local positive feedback regulation determines cell shape in root hair cells. *Science* 319:1241–1244
- Tominaga M, Yokota E, Vidali L, Sonobe S, Hepler PK, Shimmen T (2000) The role of plant villin in the organization of the actin cytoskeleton, cytoplasmic streaming and the architecture of the transvacuolar strand in root hair cells of *Hydrocharis*. *Planta* 210:836–843
- Uhrig JF, Mutondo M, Zimmermann I, Deeks MJ, Machesky LM, Thomas P, Uhrig S, Rambke C, Hussey PJ, Hülskamp M (2007) The role of *Arabidopsis* SCAR genes in ARP2-ARP3-dependent cell morphogenesis. *Development* 134:967–977
- Van Gestel K, von Witsch M, Samaj J, Baluska F (2003) Immunological evidence for the presence of plant homologues of the actin-related protein Arp3 in tobacco and maize: subcellular localization to actin-enriched pit fields and emerging root hairs. *Protoplasma* 222:45–52
- Vitha S, Baluska F, Braun M, Šamaj J, Volkmann D, Barlow PW (2000) Comparison of cryofixation and aldehyde fixation for plant actin immunocytochemistry: aldehydes do not destroy F-actin. *Histochem J* 32:457–466
- Voigt B, Timmers AC, Samaj J, Hlavacka A, Ueda T, Preuss M, Nielsen E, Mathur J, Emans N, Stenmark H, Nakano A, Baluška F, Menzel D (2005) Actin-based motility of endosomes is linked to the polar tip growth of root hairs. *Eur J Cell Biol* 84:609–621
- Wang YS, Yoo CM, Blancaflor EB (2008) Improved imaging of actin filaments in transgenic *Arabidopsis* plants expressing a green fluorescent protein fusion to the C- and N-termini of the fimbrin actin-binding domain 2. *New Phytol* 177:525–536
- Wasteneys GO, Willingale-Theune J, Menzel D (1997) Freeze shattering: a simple and effective method for permeabilizing higher plant cell walls. *J. Microsc* 188:51–61
- Xu J, Scheres B (2005) Cell polarity: ROPing the ends together. *Curr Opin Plant Biol* 8:613–618
- Young ME, Cooper JA, Bridgman PC (2004) Yeast actin patches are networks of branched actin filaments. *J Cell Biol* 166:629–635
- Zhang C, Mallery EL, Schlueter J, Huang S, Fan Y, Brankle S, Staiger CJ, Szymanski DB (2008) *Arabidopsis* SCARs function interchangeably to meet actin-related protein 2/3 activation thresholds during morphogenesis. *Plant Cell* 20:995–1011
- Zhang X, Dyachok J, Krishnakumar S, Smith LG, Oppenheimer DG (2005) IRREGULAR TRICHOME BRANCH1 in *Arabidopsis* encodes a plant homolog of the actin-related protein2/3 complex activator Scar/WAVE that regulates actin and microtubule organization. *Plant Cell* 17:2314–2326
- Zheng Z, Yang Z (2000) The Rop GTPase switch turns on polar growth in pollen. *Trends Plant Sci* 5:298–303