

Analyzing and Visualizing Open Precipitation Data

Authors: Kostas Leptokaropoulos and Shubo Chakrabarti

Copyright 2024 - 2024 The MathWorks, Inc.

Table of Contents

Introduction.....	1
Data and Resources.....	1
Import Data from the web.....	3
Read and filter Geospatial Data.....	7
Filter Data in Time and Space.....	7
Import Filtered Data.....	8
Convert units from sec-1 to day-1.....	8
Calculate dates	8
Select a day to visualize precipitation data.....	9
Visualize and compare past and projected data.....	9
Some Stats!.....	11
Compare incremental and cumulative precipitation.....	11
Generate Daily/Cumulative Plots and Box Charts.....	11
Perform Statistical Tests.....	13
Publish reusable MATLAB code for reproducible results.....	14

Introduction

Public Data: Many public databases have been created for the purposes of making data freely accessible to the scientific community. A best practice is to assign a unique identifier to a dataset, so that it is discoverable. A common form of a unique identifier is a [Digital Object Identifier or DOI®](#) which points to the data.

Access Public Data: To access and process public data, you can use several routes.

- Download data files to your local machine and work with them in MATLAB®.
- Access data directly via an API. MATLAB's function that [import data from NetCDF files](#) from RESTful API used by many portals.
- If the portal offers only Python® bindings, [call Python from MATLAB](#).

Data formats: MATLAB supports a broad range of data formats

- There is a wide range of scientific data formats that can be [natively read in MATLAB](#). They include NetCDF, HDF5 and GRIB as well as more specialized data formats.
- In addition, the [Mapping Toolbox™](#) contains [built-in functions](#) to read data from many online data repositories in standard geo-data formats.
- Sometimes data import functions may be [written by the Geoscience community](#), and published on the [MATLAB File Exchange](#) - a portal for community contributions in MATLAB. All community contributions are covered by open source licenses, which means they can be re-used, modified or added to. Exact terms and conditions depend on the licenses used by the author.

Data and Resources

In this example, we use data covered by permissive license, (i.e., the [Creative Commons Attribution 4.0 International License](#) - CC BY 4.0). It is important for Open Science to have such data since under CC BY 4.0 license you are free to share, copy, redistribute and adapt the material as long as you give appropriate credit, provide a link to the license, and indicate if any changes were made.

We access global climate data from the WCRP CMIP6 (World Climate Research Programme Coupled Intercomparison Project - Phase 6), which are located at: <https://esgf-data.dkrz.de/search/cmip6-dkrz/>.

The Data Space is hosted by the [German Climate Computer Center](#) (DKRZ) and the [Infrastructure for the European Network for Earth System Modelling](#) (IS-ENES), as a part of the global [Earth System Grid Federation](#) (ESGF).

The user can apply multiple filters and preferences for a large variety of climate parameters, such as see ice thickness, air pressure, phytoplankton mass concentration, etc, provided by different sources. In our example we used the following filtering:

- CF Standard name: precipitation flux
- Frequency: day
- Nominal Resolution: 10km, 25km

Make sure to check the box *"Show all Replicas"*.

Example links for the data from [ESGF MetaGrid](#) ([Inl.gov](#)):

- data [Link](#) (1950 - 2014); e.g., [data 19500101-19501231](#)
- data [Link](#) (2015 - 2050); e.g., [data 20450101-20451231](#)

Data Acknowledgment Statement (see [CMIP6 terms of use](#)):

“We acknowledge the World Climate Research Programme, which, through its Working Group on Coupled Modelling, coordinated and promoted CMIP6. We thank the climate modeling groups for producing and making available their model output, the Earth System Grid Federation (ESGF) for archiving the data and providing access, and the multiple funding agencies who support CMIP6 and ESGF. We used the CMIP6 model data produced by the Research Center for Environmental Changes, Academia Sinica, Taiwan (AS-RCEC), which is licensed under a Creative Commons Attribution 4.0 International License (CC BY 4.0; <https://creativecommons.org/licenses/>).

Further information about this data can be found via the `further_info_url` (recorded as a "Global Attributes: License" in these .nc files). The data producers and data providers make no warranty, either express or implied, including, but not limited to, warranties of merchantability and fitness for a particular purpose. All liabilities arising from the supply of the information (including any liability arising in negligence) are excluded to the fullest extent permitted by law.”

Import Data from the web

```
clearvars; clc; close all
```

You can read any 2 data sets from the [example links](#) indicated above, one imported as "URLpast" and a subsequent one as "URLproj".

There are a bunch of [functions that you can use to interact with NetCDF files](#).

```
% read NETCDF data from the web
URLpast
= "https://esgf-data1.llnl.gov/thredds/dodsC/css03_data/CMIP6/HighResMIP/AS-
RCEC/HiRAM-SIT-HR/highresSST-present/r1i1p1f1/day/pr/gn/v20210713/pr_day_HiRAM-SIT-
HR_highresSST-present_r1i1p1f1_gn_19530101-19531231.nc";
URLproj
= "https://esgf-data1.llnl.gov/thredds/dodsC/css03_data/CMIP6/HighResMIP/AS-
RCEC/HiRAM-SIT-HR/highresSST-future/r1i1p1f1/day/pr/gn/v20210707/pr_day_HiRAM-SIT-
HR_highresSST-future_r1i1p1f1_gn_20450101-20451231.nc";
```

Read the precipitation flux as "Variable"

```
ncdisp(URLpast)
```

```
Source:      https://esgf-data1.llnl.gov/thredds/dodsC/css03_data/CMIP6/HighResMIP/AS-RCEC/HiRAM-SIT-HR/highresSST-pr
Format:      classic
Global Attributes:
Conventions      = 'CF-1.7 CMIP-6.2'
activity_id      = 'HighResMIP'
branch_method    = 'no parent'
contact          = 'Dr. Chia-Ying Tu (cytu@gate.sinica.edu.tw)'
creation_date    = '2021-07-01T18:11:53Z'
data_specs_version = '01.00.32'
experiment       = 'forced atmosphere experiment for 1950-2014'
experiment_id    = 'highresSST-present'
external_variables = 'areacella'
forcing_index    = 1
frequency        = 'day'
further_info_url = 'https://furtherinfo.es-doc.org/CMIP6.AS-RCEC.HiRAM-SIT-HR.highresSST-pr
grid            = 'primarily 0.25deg; 1536 x 768 longitude/latitude'
grid_label      = 'gn'
history          = '2021-07-01T18:11:53Z ; CMOR rewrote data to be consistent with CMIP6, C
initialization_index = 1
institution      = 'Research Center for Environmental Changes, Academia Sinica, Nankang, Ta
institution_id    = 'AS-RCEC'
mip_era          = 'CMIP6'
nominal_resolution = '25 km'
parent_activity_id = 'no parent'
parent_experiment_id = 'no parent'
parent_mip_era    = 'no parent'
parent_source_id  = 'no parent'
parent_time_units = 'no parent'
parent_variant_label = 'no parent'
physics_index    = 1
product          = 'model-output'
```

```

realization_index      = 1
realm                  = 'atmos'
source                 = 'HiRAM-SIT-HR (2018):
                        aerosol: none
                        atmos: GFDL-HiRAM (Cubed-sphere (c384) - 0.25 degree nominal horizontal
                        atmosChem: none
                        land: GFDL-LM3 (same grid as atmos)
                        landIce: none
                        ocean: SIT (1-D, tripolar - nominal 0.25 deg; 1440 x 1080 longitude/lat
                        ocnBgchem: none
                        seaIce: none'
source_id              = 'HiRAM-SIT-HR'
source_type            = 'AGCM AER'
sub_experiment         = 'none'
sub_experiment_id      = 'none'
table_id               = 'day'
table_info             = 'Creation Date:(28 May 2020) MD5:7d39246ef1034245efba3e04f5c9ac59'
title                  = 'HiRAM-SIT-HR output prepared for CMIP6'
variable_id            = 'pr'
variant_label          = 'r111p1f1'
license                = 'CMIP6 model data produced by REQUIRED is licensed under a Creative Com
cmor_version           = '3.5.0'
tracking_id            = 'hdl:21.14100/708b6a19-2ace-4e8f-b90c-23e3dcfc050f'
DODS_EXTRA.Unlimited_Dimension = 'time'

```

Dimensions:

```

time = 365      (UNLIMITED)
bnds = 2
lat  = 768
lon  = 1536

```

Variables:

```

time
  Size:          365x1
  Dimensions:    time
  Datatype:      double
  Attributes:
    bounds       = 'time_bnds'
    units        = 'days since 1948-1-1 00:00:00'
    calendar     = 'standard'
    axis         = 'T'
    long_name    = 'time'
    standard_name = 'time'
    _ChunkSizes  = 1

time_bnds
  Size:          2x365
  Dimensions:    bnds,time
  Datatype:      double
  Attributes:
    _ChunkSizes = [1 2]

lat
  Size:          768x1
  Dimensions:    lat
  Datatype:      double
  Attributes:
    bounds       = 'lat_bnds'
    units        = 'degrees_north'
    axis         = 'Y'
    long_name    = 'Latitude'
    standard_name = 'latitude'

lat_bnds
  Size:          2x768
  Dimensions:    bnds,lat
  Datatype:      double
  Attributes:
    _ChunkSizes = [768 2]

```

```

lon
  Size:      1536x1
  Dimensions: lon
  Datatype:  double
  Attributes:
    bounds      = 'lon_bnds'
    units       = 'degrees_east'
    axis        = 'X'
    long_name    = 'Longitude'
    standard_name = 'longitude'

lon_bnds
  Size:      2x1536
  Dimensions: bnds,lon
  Datatype:  double
  Attributes:
    _ChunkSizes = [1536      2]

pr
  Size:      1536x768x365
  Dimensions: lon,lat,time
  Datatype:  single
  Attributes:
    standard_name = 'precipitation_flux'
    long_name     = 'Precipitation'
    comment       = 'includes both liquid and solid phases'
    units         = 'kg m-2 s-1'
    original_name = 'precip'
    original_units = 'kg/m2/s'
    history       = '2021-07-01T18:11:53Z altered by CMOR: Converted units from 'kg/m2/s' to 'kg'
    cell_methods  = 'area: time: mean'
    cell_measures = 'area: areacella'
    missing_value = 1.000000020040877e+20
    _FillValue    = 1.000000020040877e+20
    _ChunkSizes   = [1  384  768]

```

ncdisp(URLproj)

```

Source:      https://esgf-data1.llnl.gov/thredds/dodsC/css03_data/CMIP6/HighResMIP/AS-RCEC/HiRAM-SIT-HR/highresSST-fu
Format:      classic
Global Attributes:
  Conventions      = 'CF-1.7 CMIP-6.2'
  activity_id      = 'HighResMIP'
  branch_method    = 'standard'
  branch_time_in_child = 24472
  branch_time_in_parent = 24472
  contact          = 'Dr. Chia-Ying Tu (cytu@gate.sinica.edu.tw)'
  creation_date    = '2021-06-29T11:29:22Z'
  data_specs_version = '01.00.32'
  experiment       = 'forced atmosphere experiment for 2015-2050 using SST/sea-ice derived fr
  experiment_id    = 'highresSST-future'
  external_variables = 'areacella'
  forcing_index     = 1
  frequency        = 'day'
  further_info_url  = 'https://furtherinfo.es-doc.org/CMIP6.AS-RCEC.HiRAM-SIT-HR.highresSST-fu
  grid             = 'primarily 0.25deg; 1536 x 768 longitude/latitude'
  grid_label       = 'gn'
  history          = '2021-06-29T11:29:22Z ; CMOR rewrote data to be consistent with CMIP6, C
  initialization_index = 1
  institution      = 'Research Center for Environmental Changes, Academia Sinica, Nankang, Ta
  institution_id    = 'AS-RCEC'
  mip_era          = 'CMIP6'
  nominal_resolution = '25 km'
  parent_activity_id = 'HighResMIP'

```

```

parent_experiment_id      = 'highresSST-present'
parent_mip_era            = 'CMIP6'
parent_source_id         = 'HiRAM-SIT-HR'
parent_time_units        = 'days since 1948-01-01 00:00:00'
parent_variant_label     = 'r1i1p1f1'
physics_index            = 1
product                  = 'model-output'
realization_index        = 1
realm                    = 'atmos'
source                   = 'HiRAM-SIT-HR (2018):
                           aerosol: none
                           atmos: GFDL-HiRAM (Cubed-sphere (c384) - 0.25 degree nominal horizontal
                           atmosChem: none
                           land: GFDL-LM3 (same grid as atmos)
                           landIce: none
                           ocean: SIT (1-D, tripolar - nominal 0.25 deg; 1440 x 1080 longitude/latitude)
                           ocnBgchem: none
                           seaIce: none'

source_id                = 'HiRAM-SIT-HR'
source_type              = 'AGCM AER'
sub_experiment            = 'none'
sub_experiment_id        = 'none'
table_id                 = 'day'
table_info                = 'Creation Date:(28 May 2020) MD5:7d39246ef1034245efba3e04f5c9ac59'
title                    = 'HiRAM-SIT-HR output prepared for CMIP6'
variable_id              = 'pr'
variant_label            = 'r1i1p1f1'
license                  = 'CMIP6 model data produced by REQUIRED is licensed under a Creative Commons
cmor_version              = '3.5.0'
tracking_id              = 'hdl:21.14100/bc166568-c398-40fb-9db3-391252149fdf'
DODS_EXTRA.Unlimited_Dimension = 'time'

```

Dimensions:

```

time = 365    (UNLIMITED)
bnds = 2
lat  = 768
lon  = 1536

```

Variables:

time

```

Size:          365x1
Dimensions:    time
Datatype:      double
Attributes:
    bounds      = 'time_bnds'
    units       = 'days since 1948-1-1 00:00:00'
    calendar    = 'standard'
    axis        = 'T'
    long_name    = 'time'
    standard_name = 'time'
    _ChunkSizes = 1

```

time_bnds

```

Size:          2x365
Dimensions:    bnds,time
Datatype:      double
Attributes:
    _ChunkSizes = [1 2]

```

lat

```

Size:          768x1
Dimensions:    lat
Datatype:      double
Attributes:
    bounds      = 'lat_bnds'
    units       = 'degrees_north'
    axis        = 'Y'
    long_name    = 'Latitude'

```

```

                standard_name = 'latitude'
lat_bnds
    Size:          2x768
    Dimensions:    bnds,lat
    Datatype:      double
    Attributes:
        _ChunkSizes = [768    2]
lon
    Size:          1536x1
    Dimensions:    lon
    Datatype:      double
    Attributes:
        bounds      = 'lon_bnds'
        units       = 'degrees_east'
        axis        = 'X'
        long_name    = 'Longitude'
        standard_name = 'longitude'
lon_bnds
    Size:          2x1536
    Dimensions:    bnds,lon
    Datatype:      double
    Attributes:
        _ChunkSizes = [1536    2]
pr
    Size:          1536x768x365
    Dimensions:    lon,lat,time
    Datatype:      single
    Attributes:
        standard_name = 'precipitation_flux'
        long_name      = 'Precipitation'
        comment        = 'includes both liquid and solid phases'
        units          = 'kg m-2 s-1'
        original_name   = 'precip'
        original_units  = 'kg/m2/s'
        history         = '2021-06-29T11:29:22Z altered by CMOR: Converted units from 'kg/m2/s' to 'kg'
        cell_methods    = 'area: time: mean'
        cell_measures    = 'area: areacella'
        missing_value    = 1.000000020040877e+20
        _FillValue      = 1.000000020040877e+20
        _ChunkSizes     = [1  384  768]

```

```
Variable = "pr";
```

Read and filter Geospatial Data

read longitude and latitude from the .nc file

```
lon = double(ncread(URLpast,'lon'));
lat = double(ncread(URLpast,'lat'));
```

Filter Data in Time and Space

Select boundaries of a rectangular area

```
%minimum and maximum longitude
minlon = 25;maxlon = 40; % set values from 0 to 360
%minimum and maximum latitude
minlat = 35;maxlat = 50; % set values from -90 to 90
```

Select first day and number of days after the first (*their sum must not exceed 365*)

```
starttime = 100;counttime = 190;% set values from 1 to 365
```

Find the indexes of longitude and latitude that correspond to the selected geographical boundaries

```
indlon = find(lon>=minlon & lon<=maxlon);
indlat = find(lat>=minlat & lat<=maxlat);

startlon=indlon(1);countlon=numel(indlon);
startlat=indlat(1);countlat=numel(indlat);
```

Import Filtered Data

Import longitude, latitude, time and precipitation values for past and future data, as constrained in the [previous section](#).

```
lat = double(ncread(URLpast,'lat',startlat,countlat));
lon = double(ncread(URLpast,'lon',startlon,countlon));
tpast = double(ncread(URLpast,'time',starttime,counttime));
tproj = double(ncread(URLproj,'time',starttime,counttime));
Varpast = double(ncread(URLpast,Variable,[startlon startlat starttime],[countlon
countlat counttime]));
Varproj = double(ncread(URLproj,Variable,[startlon startlat starttime],[countlon
countlat counttime]));
```

read the precipitation unit from the .nc file

```
unit = ncreadatt(URLpast,'pr','original_units');
```

Convert units from sec-1 to day-1

```
Varpast = Varpast*86400;
Varproj = Varproj*86400;
newunit = "kg m^-2 day^-1"
```

```
newunit =
"kg m^-2 day^-1"
```

Calculate dates

calculate date vectors, as days passed since the reference date in the .nc file

```
Tpast = datetime('1948-1-1')+tpast,Tproj = datetime('1948-1-1')+tproj
```

```
Tpast = 190x1 datetime
10-Apr-1953 12:00:00
11-Apr-1953 12:00:00
12-Apr-1953 12:00:00
13-Apr-1953 12:00:00
14-Apr-1953 12:00:00
15-Apr-1953 12:00:00
16-Apr-1953 12:00:00
```



```

17-Apr-1953 12:00:00
18-Apr-1953 12:00:00
19-Apr-1953 12:00:00
:
Tproj = 190x1 datetime
10-Apr-2045 12:00:00
11-Apr-2045 12:00:00
12-Apr-2045 12:00:00
13-Apr-2045 12:00:00
14-Apr-2045 12:00:00
15-Apr-2045 12:00:00
16-Apr-2045 12:00:00
17-Apr-2045 12:00:00
18-Apr-2045 12:00:00
19-Apr-2045 12:00:00
:

```

```

Tpast.Format = 'dd-MMM-uuuu';Tproj.Format = 'dd-MMM-uuuu';
Tpast_str = string(Tpast);
Tproj_str = string(Tproj);

```

Select a day to visualize precipitation data

```
close all
```

Select a day

```
t1 = Tpast_str(6);inx = Tpast_str == t1;
```

Create a grid with precipitation values

```

[lat1,lon1] = meshgrid(lat,lon);
Var1past = Varpast(:,:,inx);
Var1proj = Varproj(:,:,inx);
mV = min([min(Var1past(:)),min(Var1proj(:))])

```

```
mV = -9.0042e-16
```

```
MV = max([max(Var1past(:)),max(Var1proj(:))])
```

```
MV = 51.4354
```

Customize colormap limits to the selected data

```

CBmin = -9.00419266706534e-16;
CBmax = 51.43537279218435;

```

Visualize and compare past and projected data

Create a 2 frame figure. Plot the past data in the left frame and the projected data in the right frame. Add the coastlines in both figures. The text displayed in the figure (title, colorbar) is also directly imported from the .nc file metadata and the parameter values we have set.


```

tilayout(1,2)
nexttile % past
worldmap([minlat maxlat],[minlon maxlon])
geoshow(lat1,lon1,Var1past,'DisplayType','texturemap','FaceAlpha',0.5);
geoshow('landareas.shp','EdgeColor','black','FaceColor','none')
title([ncreadatt(URLpast,Variable,'long_name'),'on',t1])
    clim([CBmin CBmax])

nexttile %projection
worldmap([minlat maxlat],[minlon maxlon])
geoshow(lat1,lon1,Var1proj,'DisplayType','texturemap','FaceAlpha',0.5);
geoshow('landareas.shp','EdgeColor','black','FaceColor','none')
    cb = colorbar; % display colorbar
    cb.Location='eastoutside';
    cb.Label.String = (newunit);
title([ncreadatt(URLproj,Variable,'long_name'),'on',Tproj_str(inx)])
    clim([CBmin CBmax])

colormap turbo

```



Some Stats!

In this section a comparison is performed between the past and projected data.

- First, the precipitation time series (daily, solid lines; cumulative, dashed lines) are plotted in the same plot for direct comparison.
- Then, [box charts](#) are generated to show the daily sum average precipitation values distribution (including the outliers)
- Finally, 3 Hypothesis Tests are performed to evaluate the statistical significance of the results from the comparison of the past and projected precipitation:

a) The [Wilcoxon rank sum test](#): Tests the null hypothesis that past and projected data are samples from continuous distributions with equal medians, against the alternative that they are not.

b) The [2 sample t-test](#): Tests the null hypothesis that past and projected data come from independent random samples from normal distributions with equal means and equal but unknown variances. The alternative hypothesis is that the past and projected data comes from populations with unequal means.

c) The 2 sample [Kolmogorov - Smirnov test](#): Tests the null hypothesis that that past and projected data are from the same continuous distribution against the alternative hypothesis that they are from different continuous distributions.

Compare incremental and cumulative precipitation

```
N = numel(tpast)
```

```
N = 190
```

Select number of days (between 1 and N)

```
T1 = Tpast_str(1);inx1 = Tpast_str == T1;n1 = find(inx1);  
T2 = Tpast_str(190);inx2 = Tpast_str == T2;n2 = find(inx2);  
n = n2 - n1 +1;  
if n2<=n1;error('End date (n2) should be greater than start date (n1)');end
```

Generate Daily/Cumulative Plots and Box Charts

```
Boxp1 = zeros(size(n,1));Boxp2 = zeros(size(n,1));cou=1;  
for i=n1:n2  
Boxp1(cou) = sum(Varpast(:,:,i),"all");  
Boxp2(cou) = sum(Varproj(:,:,i),"all");  
cou=cou+1;  
end
```


Daily and Cumulative plots

```
figure
```

```

plot(Tpast(n1:n2),Boxp1,'r-',Tpast(n1:n2),Boxp2,'k-',LineWidth=1);datetick('x',6);yl
abel('daily sum');xlabel('date');
hold on
yyaxis
right;plot(Tpast(n1:n2),cumsum(Boxp1),'r--',Tpast(n1:n2),cumsum(Boxp2),'k--');dateti
ck('x',6);ylabel('cumulative');xlabel('date');
legend(num2str(year(Tpast(1))),num2str(year(Tproj(1))),num2str(year(Tpast(1))),num2s
tr(year(Tproj(1))))

```




Box charts

```

figure
group = categorical([repmat("past", size(Boxp1));repmat("projection",
size(Boxp2))]);
boxchart([Boxp1',Boxp2']);xticklabels({'past','projection'});
title(["Daily Sum Average","past/projection ratio = "+num2str(sum(Boxp1)/
sum(Boxp2),'%7.1f')])

```



Perform Statistical Tests

```
[p.Wilcoxon,h.Wilcoxon] = ranksum(Boxp1,Boxp2); % null hypothesis, h0, equal medians
[h.ttest2,p.ttest2] = ttest2(Boxp1,Boxp2); % null hypothesis, h0, equal means
[h.kstest2,p.kstest2] = kstest2(Boxp1,Boxp2); % null hypothesis, h0, samples come
from the same distribution
h.ttest2 = logical(h.ttest2);
```

```
if h.Wilcoxon==0;disp(['>> The <strong>Wilcoxon</strong> rank sum test FAILS to
reject the hypothesis of equal medians at ',num2str(p.Wilcoxon) ' significance'])
else; disp(['>> The <strong>Wilcoxon</strong> rank sum test REJECTS the hypothesis
of equal medians at ',num2str(p.Wilcoxon), ' significance'])
end
```

>> The **Wilcoxon** rank sum test REJECTS the hypothesis of equal medians at 0.0017175 significance

```
if h.ttest2==0;disp(['>> The 2-sample <strong>t-test</strong> FAILS to reject the
hypothesis of equal means at ', num2str(p.ttest2) ' significance'])
else; disp(['>> The 2-sample <strong>t-test</strong> REJECTS the hypothesis of
equal means at ',num2str(p.ttest2), ' significance'])
end
```

>> The 2-sample **t-test** REJECTS the hypothesis of equal means at 0.011602 significance

```

if h.kstest2==0;disp(['>> The 2-sample <strong>Kolmogorov-Smirnov</strong> test
FAILS to reject the hypothesis that the two samples come from the same distribution
at ',num2str(p.kstest2) ' significance'])
else; disp(['>> The 2-sample <strong>Kolmogorov-Smirnov</strong> test REJECTS
the hypothesis that the two samples come from the same distribution at
',num2str(p.kstest2), ' significance'])
end


```

>> The 2-sample **Kolmogorov-Smirnov** test REJECTS the hypothesis that the two samples come from the same distribution

Publish reusable MATLAB code for reproducible results


To enable collaboration partners, reviewers and the community reuse your MATLAB code and reproduce your results.

- Publish your MATLAB code (eg: on GitHub) and generate a [DOI](#) (digital object identifier) by [linking it to a DOI generating portal](#) (e.g. [Figshare®](#), [Zenodo®](#)). Make your research output findable by including as much information as needed in the metadata. Document your code well explaining steps required to reproduce clearly and explicitly.
- Make sure you include a license for your code that specifies reuse and re-distribution rights for the code. Various open source licenses are [available](#). BSD, MIT and Apache licenses are commonly used for open research software.
- [Link your GitHub repository to File Exchange](#) to make your MATLAB code available to MATLAB users via the Add-Ons button.




- Make your MATLAB code **interoperable**. MATLAB is [interoperable](#) with several other languages including C, Fortran and Python. MATLAB can be directly called from Python using the [MATLAB Engine for Python®](#) which is available as a PyPI® package and can be installed using the command `pip install matlab.engine` from Python. MATLAB code can also be [packaged as a Python library](#) and called from Python. Deep Learning models from other frameworks are [interoperable with MATLAB](#) either using the [ONNX™ interface](#) or via direct interfaces that exist, for example, for Pytorch® and Tensorflow™ models.

Functions that Import Deep Learning Networks



- MATLAB is interoperable with cloud architectures such as [JupyterHub®](#) and MATLAB code can also be used within Jupyter Notebooks. Here is a link to a Jupyter® notebook of the same example used here. There is an official MATLAB kernel for Jupyter Notebooks - read about it [here](#). **To easily convert a Live Script into a Jupyter® notebook use the [export](#) function.**



- Run your [MATLAB code on the browser directly from GitHub](#). Copy and paste the GitHub repo address into [this app](#). That will generate a command, which when pasted into your README, will create a "Open in MATLAB Online™" button on your GitHub repository. By clicking on this button, users will be able to run your code in the browser on MATLAB Online.

Open a GitHub repository in MATLAB Online

Author and repository name or URL*

author-name/repo-name or URL

File path (optional)

Path to a file

Line number (optional)

#

Project path (optional)

Path to a project (.prj file)

Copy the URL below to share a link that opens this repository in MATLAB Online

Fill in the fields above to create link



Copy the markdown below and paste it into your README to display this button: [Open in MATLAB Online](#)

Fill in the fields above to create markdown



- Make your MATLAB code reproducible by using a reproducibility portals that supports MATLAB. One example is Code Ocean®. On Code Ocean, you can [upload your MATLAB code](#) including dependencies. Once uploaded, your code is tested and published as a Code Ocean "capsule" which can be run online or downloaded and run locally by users. Code Ocean also generates a DOI for your code capsule. For Live Scripts, convert the .mlx file into e.g., a .m file, a .html or a.ipynb file using the [export](#) function for best results. Here is the DOI for the Code Ocean capsule of the this code. Read more about MATLAB on Code Ocean [here](#).

Code Reproducibility and Reuse Sites that Host MATLAB

A number of sites that focus on code reproducibility and reuse host MATLAB on the cloud. Researchers can view and download the MATLAB code for their own use, while publishers, such as Nature, use the sites to conduct peer reviews including running and verifying the code for computation-based papers.

For example, we shared [information about Code Ocean](#) a few years ago, showcasing how researchers can upload their code and run code posted by others on [Code Ocean](#) without needing to access their own MATLAB license.

The screenshot displays the Code Ocean web interface. On the left is a file browser showing the contents of a capsule, including folders like 'environment', 'code', and 'data', and files like 'LICENSE', 'run_example_1.m', and 'run_example_2.m'. The central area shows the 'Metadata' for a capsule named 'g_rect: a MATLAB package for georectifying oblique images'. It includes a description of the toolbox, the authors (Daniel Bourgault, Rich Pawlowicz, Clark Richards), and a DOI: 10.24438/CO.4927878.v1. The right sidebar shows a 'Reproducible Run' section with a timeline of events, including 'Published Version 1.0' and 'Created capsule'.

- **Warning:** Before making your code available on the cloud, make sure all dependencies including any data that is needed for your code to run is uploaded along with the code. Also make sure any path and/or filenames that refer to local directories are appropriately renamed.

- FAIR standards: FAIR is an acronym that stands for **F**indable, **A**ccessible, **I**nteroperable and **R**eproducible. It is an [accepted standard](#) for research output (code, data) and is often required for your research results to be in [compliance with "Open Science" standards](#). Adhering to the above pointers helps in making your MATLAB code FAIR

Copyright 2024 - 2024 The MathWorks, Inc.