

# Ondansetron in alcohol use disorder: a systematic review

Matias Ceau \*

## Abstract

**Context** Alcohol use disorder (AUD) is a frequent disorder. Few treatments have shown a great efficacy in clinical studies. It is hypothesized that this is due to the fact that AUD suffering people constitute a heterogeneous group composed of various endophenotypes. Ondansetron, a selective 5-HT<sub>3</sub> receptor antagonist, has been evaluated in AUD and particularly in certain subgroups of patients. This review presents clinical studies evaluating its impact.

**Methods** A systematic review was conducted on Pubmed and Wiley Online Library. All 19 clinical trials involving the use of ondansetron in AUD were included.

**Results** Ondansetron was found to be effective mostly in certain subgroups of AUD-suffering patients. The identified subgroups were based on clinical evaluation (age of onset, personality type) and genotype. A number of limitations remains and further studies are needed.

**Keywords** *alcohol use disorder, alcohol dependence, ondansetron, 5-HT<sub>3</sub> receptors, serotonin, craving, pharmacogenetics, personalized medicine*

## Introduction

Alcohol use disorder (AUD) is a heterogeneous and chronic relapsing disorder resulting in a complex interaction between neurobiological, genetic and environmental factors. Despite the demonstrated efficacy of some approved medications (acamprosate, naltrexone, disulfiram), a key barrier is the fact that these medications are not effective in every patient pointing out the need for more personalized therapy approaches to overcome this heterogeneity. In this perspective, major advances in pharmacogenetics identified distinct clinical subgroups of AUD according to genetic profiles, that could be associated with differential treatment responses. Thus, the identification of patient subgroups that are most likely to respond favorably to different medications is crucial with a need for a better targeting of medication to specific patients.

Ondansetron (IUPAC name: (RS)-9-Methyl-3-[(2-methyl-1H-imidazol-1-yl)methyl]-2,3-dihydro-1H-carbazol-4(9H)-one), is a selective antagonist of the 5-HT<sub>3</sub> receptor. It is approved by ANSM in France and FDA in the USA as an antiemetic for cancer treatment-induced and anesthesia-related nausea and vomiting.

Serotonin function is an important determinant of alcohol consumption through its relationship with the mesolimbic dopaminergic reward system. In the late 1980s, Hagan et al. [1] showed that the injection of ondansetron in the ventral

tegmental area of the rat brain lessened induced hyperactivation in the nucleus accumbens.

The rewarding effect of alcohol being mediated by dopaminergic activity in the nucleus accumbens, researchers have hypothesized that ondansetron, by modulating the reinforcing effect of alcohol, could also reduce alcohol consumption in AUD suffering patients.

Some phase 1 clinical studies [2], [3] have shown promising results on the ability of ondansetron to attenuate the reinforcing properties of alcohol and of the desire to use by 5-HT<sub>3</sub> receptor blockade, in healthy male volunteers.

Later phase 2 clinical studies ([4], [5]) have shown differential effects among alcohol-use disorder suffering patients. Different typologies have been previously proposed, based on personality such as type I/II (Cloninger et al. 1987 [6]) or type A/B personalities (Babor et al. 1992 [7], [8]) or simply defined by the age of onset of alcoholism (Early-onset alcoholism/Late-onset alcoholism (EOA/LOA), Varma et al. 1994 [9]) ) characterized by differential course, prognosis and treatment responses. Results from these studies indicated distinct effect of ondansetron among EOA compared to LOA, which is hypothesized to be linked to genetic variations among individuals.

As the functional state of the serotonin transporter protein (5-HTT) is an important factor of the serotonergic function control, more recent pharmacogenetic studies have investigated the potential role of 5-HTT genotype on drinking behaviors and alcohol craving ([10]). 5-HTT gene polymorphisms, involving two variants, a short form (S) and a

\*Collège Sciences médicales, Université de Bordeaux. Email: matias.ceau@u-bordeaux.fr

long form (L), have been shown to be associated with differential 5-HT neurotransmission, which could moderate the rewarding effects and the craving for alcohol, and thereby ondansetron treatment response among AUD suffering patients.

Direct inhibition of 5-HT<sub>3</sub> receptor is thus hypothesized to alleviate alcohol craving, that is currently considered as a key determinant of relapse vulnerability as well as a major treatment target. A better knowledge of the potential impact of 5-HT antagonist medication on alcohol consumption and craving in AUD, as well as individual characteristics and genotype associated with treatment response is therefore a critical issue to improve treatment approaches and develop personalized medicine in the pharmacotherapy of alcohol use disorder.

We conducted a systematic review of the literature in order to summarize scientific evidence of the efficacy of ondansetron on alcohol consumption and craving as well as the genetic moderation of treatment response

## Methods

This systematic review was registered in PROSPERO (Registration Number:).

**Databases and search strategy** A PubMed search was conducted in the MEDLINE database and the Wiley Online Library.

For Medline search, the relevant articles were identified by combining the terms: ("ondansetron"[MeSH Terms] OR "ondansetron"[All Fields]) AND ("alcoholism"[MeSH Terms] OR "alcoholism"[All Fields] OR "alcohol use disorder"[All Fields] OR "alcohol abuse"[All Fields] OR "AUD"[All Fields])

For the Wiley Online Library search we used the keywords "ondansetron" and "alcoholism".

Two authors independently examined titles and abstracts. Relevant articles were obtained in full-text and assessed for inclusion criteria blindly by the two reviewers. Disagreement was resolved via discussion to reach consensus. Data from the eligible articles were independently extracted by two reviewers using a standardized data extraction form. Extracted data included participant characteristics, study design, treatment outcomes and results, how confounders were controlled for, and limitations.

Included studies were all clinical trials involving the use of ondansetron as a treatment of AUD.

Exclusion criteria were studies in healthy volunteers, pre-clinical studies, reviews, opinion papers, protocols, case reports, and studies not published in English.




FIGURE 1: Flow chart

## Results

**Search results** The Prisma flowchart presenting the studies selection is shown in Figure 1. A total of 90 results were found in the MEDLINE database and 8 additional studies were found on Wiley Online Library.

In total, 92 unique records were screened. Of the screened studies, 19 were finally retained meeting the inclusion criteria, published between 1994 and 2015.

**Study characteristics** Of the 19 clinical trials, 17 were randomized controlled trials (RCT) or analysis of previous RCT, and 2 were prospective open-label studies [5] [18]. The study duration ranged from 2 to 12 weeks.

The 19 clinical trials, published between 1994 and 2015, analysed 10 distinct study populations described in Table 1.

In total, 1053 subjects were enrolled, of which 1036 met criteria for alcohol use disorder. Patients were mostly males (75.7 %), and had a mean age of 40.7.

Patients enrolled were diagnosed as alcohol dependent according to the DSM-III-TR ([11], [4]), DSM-IV ([19], [12], [13], [5], [18], [21]), DSM-IV-TR ([20], [26]) or ICD-10 ([25]). Some studies required additional criteria, such as more than 35 standard drinks per week for men or 28 for women ([20], [26]), more than 30 drinks per week for men or 21 for women ([23]), more than 3 standard drinks per day and a Michigan Alcohol Screening Test greater than 5 ([12], [4], [13]) or an AUDIT score greater than 8 ([21]). Whereas most trials ([11], [12], [4], [13], [5], [18]) concerned treatment-seeking patients, some ([19], [20], [26]) did not. Most of the studies excluded patients using any other drug than alcohol.

In most studies, using drugs was considered as an exclusion criteria either directly ([5], [19], [20], [21], [26]) and/or after positive drug screening ([11], [12], [4], [13], [19], [25]). Nicotine use and nicotine use disorder were often not considered as an exclusion criteria ([4], [5]) and cannabis use was tolerated in a few trials ([18], [19]).

Psychiatric disorders were exclusion criteria in most studies ([4], [12], [13] [5], [20], [21], [27]). Two studies specified

TABLE 1: Population description

Original publication; subsequent analysis	Location	Sample size	Proportion of male	Mean age	Dropout rate	Treatment seeking	Psychiatric diagnosis	Other SUD‡
A Sellers 1994 [11]	North America	71	100 %	43.55	21 %	yes	yes*	no
B Johnson 2000a [12]; [13][14]	San Antonio, USA	20	75 %	38.0	30 %	yes	no	no
C Johnson 2000b [4]; [15][16][17]	Houston, USA	321	70 %	40.6	41 %	yes	no	no
D Kranzler 2003 [5]	North America	40	67.5 %	43.7	30 %	yes	no	no
E Dawes 2005 [18]	San Antonio, USA	12	58.3 %	18	25 %	yes	yes*	yes†
F Myrick 2008 [19]	Northridge, USA	107	72.9 %	25.69	N.A.	no	no	yes†
G Kenna 2009 [20]	Providence, USA	20	80 %	44.1	25 %	no	no	no
H Johnson 2011 [21]; [22][23][24]	North America	283	73.1 %	44.7	33 %	yes	no	no
I Corrêa Filho 2013 [25]	São Paulo, Brazil	102	100 %	42.91	50 %	Un.	yes*	no
J Kenna 2014 [26]; [27]	Providence, USA	77	65 %	43.4	29 %	no	no	no
Total		1053	75.7 %	40.7	35.4 %	78.5 %	17.6 %	11.3 %

Description of the 10 distinct study population used by the 19 clinical trials. Letters A to J are attributed to each study population in chronological order of first publication for later referral.

‡: *except nicotine*; †: *cannabis use allowed*; \*: *except clinically significant disorders*; c: *except cannabis-use disorder*; n: *except nicotine-use disorder*; N.A.: *non applicable*; SD: *standard deviation*; SUD: *substance-use disorder*; Un.: *unknown*

that only major diagnosis ([19]) or clinically significant disorders ([25] [18]) were exclusion criteria major. In the study of Edward et al. ([11]) a Montgomery/asberg Depression scale score below 15 and a Spielberger State-Trait anxiety inventory score below 55 were required.

Many studies also required good health at enrollment and notably excluded frequent AUD comorbidities such as elevated bilirubin ([4], [20]), liver enzymes ([18], [20]), liver cirrhosis ([25]) or severe alcohol withdrawal ([12], [4], [18]).

Different subgroups were identified. Early onset alcoholics (EOA) and late onset alcoholics (LOA) were defined by the age of onset of their substance use disorder, respectively before or after the age of 25. This paradigm was used in 8 studies ([4], [12], [13], [14], [15], [16], [5], [17]). Six studies ([21], [23], [22], [27], [26], [24]) delineated subgroups based on their genotype. SLC6A4, the gene coding for the serotonin transporter contains a polymorphism in the promoter region, the 5-HTT-linked polymorphic region, with a "short" (S) and "long" (L). Individuals possessing two long alleles (L/L) have been found to respond differently to serotonergic treatments than those having either one (L/S) or two (S/S) short alleles.

**Treatment protocols** Ondansetron dosage was between 1 µg/kg bid (twice a day) and 16 mg per day with the most frequent dosage being 4 µg/kg bid. Four studies (127 patients) used ondansetron in combination with naltrexone at a dosage of 50 mg per day ([12], [13], [14], [19]). Three stud-

ies (97 patients) evaluated ondansetron against sertraline, at a dosage of 200 mg per day ([20], [27], [26]).

The treatment duration varied between 8 days to 11 weeks.

Treatment compliance was evaluated through pill count or riboflavin dosage.

**Treatment outcomes** Efficacy was mostly assessed by evaluating the number of standard drinks and derived variables such as defined by the Alcohol Timeline Followback (TLFB) method [28]. Drinking outcomes were drinks per day (DD), drinks per drinking day (DDD), percentage of day abstinent (PDA), heavy drinking days (days with more than 5 drinks per day) percentage of heavy drinking day (PHDD). Standard drink definition varied between different studies, it was defined as 12 g ([4][21]), 13 g ([11]) or 14 g[25]) of pure ethanol.

Some studies ([13], [15][19]) evaluated alcohol craving, either with a visual analogical scale or with the obsessive compulsive drinking scale (OCDS [29]).

One study ([19]) used functional magnetic resonance imaging to determine ventral striatum activation. Another study used the Profile of Mood States [30] to evaluate attenuation of mood disturbances.

**Study results** The 19 clinical trials were subdivided in two main subgroups according to their primary outcomes. The first set of studies (Tables 2, 3 and 4) evaluated the efficacy of ondansetron, by evaluating its impact on alcohol

TABLE 2: Treatment efficacy on alcohol use reduction

	Sample characteristics	Design	Study Timeframe	Treatment protocol		Primary outcome measures	Results
	Size (†)		(weeks)	Dosage	Duration		
Sellers 1994 [11]	74 (A)	RCT	9	OND: 0.25 mg 2mg	6 w	DDD (standard drink 13g)	Principal analysis: trend but N.S.; <i>Post hoc</i> analysis: effective when heavy drinkers (>10DD) excluded *
Johnson 2000a [12]	20 (B)	RCT	8	OND 4 µg/kg bid +NAL 25 mg bid	8 w	DD, DDD and PDA	Reduced DD*, DDD* and PDA (N.S) compared to placebo
Johnson 2000b [4]	321 (C)	RCT	12	OND: 1 µg/kg bid 4 µg/kg bid 16 µg/kg bid	11 w	DD, PDA and plasma CDT (standard drink 12 g)	In EOA, OND was superior* to placebo on DD, DDD for all dosage and on CDT for 1 and 4 µg/kg bid. OND at 4µg/kg bid was superior to placebo on PDA and DAW
Ait-Daoud 2001b [14]	20 (B)	RCT	8	OND 4 µg/kg bid +NAL 50 mg	8 w	log serum CDT	EOA treated with OND+NAL had lower CDT levels compared to placebo*
Kranzler 2003 [5]	40 (D)	PT	8	OND: 4 µg/kg bid	8 w	DD, DrInC score, log CDT ratio	DD, DDD, EOA had greater* decrease in DD, DDD and alcohol related problems than LOA
Kenna 2009 [20]	20 (G)	RCT	8	SER 200 mg OND 0.5 mg	2/3 w	ASAE volume (mL), DDD	In L/L, OND reduced* ASAE alcohol volume and DDD compared to SER
Johnson 2011 [21]	283 (H)	RCT	12	OND: 4 µg/kg bid	11 w	DDD, PDA	In L/L, less DDD and higher PDA in OND vs. placebo*; In OND, less DDD and higher PDA in L/L vs L-S/S
Corrêa Filho 2013 [25]	102 (I)	RCT	12	OND: 16 mg	12 w	PDA, PHDD (standard drink 14g)	OND was superior than placebo to reduce %HDD *
Kenna 2014a [26]	77 (J)	RCT	9	SER 200 mg or OND 0.5 mg	3 w	ASAE volume (mL), DDD	OND resulted in reduction in DDD compared to SER*
Kenna 2014b [27]	77 (J)	RCT	9	SER 200 mg or OND 0.5 mg	3 w	ASAE volume (mL), DDD	Among women only, L/L + OND and L-S/S + SER had fewer DDD* and reduced ASAE* L/L + SER and L-S/L + OND

\*: significant result; †: population (A) to (J) as defined in Table 1 ASAE: alcohol self-administration experiment; bid: twice a day; CDT: carbohydrate deficient transferrin; d: day; DD: drinks per day; DDD: drinks per drinking day; L/L: L/L genotype; L-S/S: L/S or S/S genotype; NAL: naltrexone; N.S.: non significant; OND: ondansetron; PDA: proportion of days abstinent; PHDD: percentage of heavy drinking days; SD: standard drinks; SER: sertaline; w: week;

TABLE 3: Treatment efficacy on craving and cue-induced craving

	Sample characteristics	Design	Study Timeframe	Treatment protocol		Primary outcome measures	Results
	Size (†)		(weeks)	Dosage	Duration		
Ait-Daoud 2001a [13]	20 (B)	RCT	8	OND 4 µg/kg bid + NAL 50 mg	8 w	Craving (measured with OCDS)	OND+NAL was effective at reducing craving*; reduction in craving was correlated by reduction of drinking in medication group
Johnson 2002 [15]	253 (C)	RCT	12	OND: 1 µg/kg bid 4 µg/kg bid 16 µg/kg bid	11 w	Craving (Visual Analog Scale)	OND at 4µg/kg was associated in craving reduction in EOA but not LOA compared to placebo*
Myrick 2008 [19]	107 (F)	RCT	2	OND: 0.5 mg + NAL: 50 mg or OND: 0.5 mg or NAL: 50 mg	8 d	Craving for Alcohol Score, Alcohol Cue-Induced Ventral Striatum Activation Score	Diminished craving and cue-induced activation of the striatum with NAL*, NAL+OND* and OND (N.S)

\*: significant result; †: population (A) to (J) as defined in Table 1 ASAE: alcohol self-administration experiment; bid: twice a day; CDT: carbohydrate deficient transferrin; d: day; DD: drinks per day; DDD: drinks per drinking day; L/L: L/L genotype; L-S/S: L/S or S/S genotype; NAL: naltrexone; N.S.: non significant; OND: ondansetron; PDA: proportion of days abstinent; PHDD: percentage of heavy drinking days; SD: standard drinks; SER: sertaline; w: week;

TABLE 4: Treatment efficacy on mood disturbances

	Sample characteristics	Design	Study Timeframe	Treatment protocol		Primary outcome measures	Results
	Size (†)		(weeks)	Dosage	Duration		
Johnson 2003 [16]	253 (C)	RCT	12	OND: 1 µg/kg bid 4 µg/kg bid 16 µg/kg bid	11 w	Mood (Profile of Mood States)	OND 16 µg/kg bid was effective* on reducing some mood disturbances in EOA

\*: significant result; †: population (A) to (J) as defined in Table 1 ASAE: alcohol self-administration experiment; bid: twice a day; CDT: carbohydrate deficient transferrin; d: day; DD: drinks per day; DDD: drinks per drinking day; L/L: L/L genotype; L-S/S: L/S or S/S genotype; NAL: naltrexone; N.S.: non significant; OND: ondansetron; PDA: proportion of days abstinent; PHDD: percentage of heavy drinking days; SD: standard drinks; SER: sertaline; w: week;

TABLE 5: Moderators of treatment outcomes

	Sample characteristics	Design	Study Timeframe	Treatment protocol		Primary outcome measures	Results
	Size (†)		(weeks)	Dosage	Duration		
Dawes 2005 [18]	12 (E)	PT	8	OND: 4 µg/kg bid	8 w	Safety, tolerability, DD, DDD and PDA	No discontinuation due to adverse effect; within-group decrease in DD* and DDD*
Roache 2008 [17]	271 (C)	RCT	11	OND: 1 µg/kg bid 4 µg/kg bid 16 µg/kg bid	11 w	DD, PDA	DDD, Babor type A/B better discriminated subtypes based on baseline severity; EOA/LEO better predicted response to OND
Seneviratne 2012 [23]	41 (H)	RCT	11	OND: 4 µg/kg bid	11 w	DDD	In OND group, DDD was associated positively with 5'-HTTLPR mRNA levels in L/L
Johnson 2013 [22]	283 (H)	RCT	11	OND: 4 µg/kg bid	11 w	DDD, PHDD, PDA	5 different genotypes predicted efficacy of OND on alcohol consumption*
Hou 2015 [24]	251 (H)	RCT	11	OND: 4 µg/kg bid	11 w	reduction of PHDD from baseline	Data mining approaches (such as interaction trees and virtual twins) successfully identified subgroups benefiting from OND

\*: significant result; †: population (A) to (J) as defined in Table 1 ASAE: alcohol self-administration experiment; bid: twice a day; CDT: carbohydrate deficient transferrin; d: day; DD: drinks per day; DDD: drinks per drinking day; L/L: L/L genotype; L-S/S: L/S or S/S genotype; NAL: naltrexone; N.S.: non significant; OND: ondansetron; PDA: proportion of days abstinent; PHDD: percentage of heavy drinking days; SD: standard drinks; SER: sertaline; w: week;

use, craving and mood disturbances. The second set (Table 5) evaluated the moderators of treatment outcomes.

**Alcohol use reduction** Ten studies, summarized in Table 2 evaluated the impact of ondansetron, alone or in combination with naltrexone on alcohol use. The main outcomes were mostly self-reported changes in alcohol consumption in standard drinks (using the previously defined TLFB method). A few studies used objective measures: plasma CDT and the volume of alcohol consumed in an alcohol self administration premenstrual.

In a sample of 71 males suffering from alcohol dependence (DSM-III-TR), Sellers et al. [11] showed a trend ( $p = 0,06$ ), and *post hoc* analysis indicated a significant impact on AUD patients drinking less than 10 drinks a day. The effect of ondansetron was shown to be non-linear, as 0,25 mg was more effective than 2 mg.

In a small scale randomized control trial [12] ( $n = 20$ , exclusively EOA), a combination of ondansetron and naltrexone showed significant effect on reduction of drinks per day ( $0.99 \pm 0.60$  vs  $3.68 \pm 0.63$ , effect size = 1.42) and drinks per drinking day ( $3.14 \pm 0.87$  vs  $6.76 \pm 0.71$ , effect size = 1.71), as well as a trend in reducing the percentage of abstinent days, compared to placebo.

A subsequent analysis of the sample by Ait-Daoud et al. showed that the combination of ondansetron and naltrexone was associated with significantly lower CDT levels [14].

In a later trial (Johnson et al. 2000b [4]), 271 patients of the 321 enrolled were given ondansetron at various dosages (1, 4 and 16  $\mu\text{g/kg}$  of body weight, twice a day). In this study, ondansetron was found to be significantly more effective than placebo in reducing alcohol consumption among EOA but not LOA. Ondansetron at 4  $\mu\text{g/kg}$  *b.i.d.*, which was (non significantly) superior to the other dosages, was more effective than placebo on drinks per day (1.56 vs 3.30,  $p = 0.01$ ), drinks per drinking day (4.28 vs 6.90,  $p = 0.004$ ), percentage of day abstinent (70.10 vs 50.20,  $p = 0.02$ ) and mean log CDT ratio (-0.19 vs 0.12,  $p = 0.01$ ). Among EOA, all other dosages were superior to placebo on the two first criteria.

These results were subsequently replicated by Kranzler et al. in 2003 [5], who showed a significant reduction (compared to baseline) in most alcohol-related measures (drinks per day, drinks per drinking days, DrinC total score) among EOA and LOA who received ondansetron (4  $\mu\text{g/kg}$  *b.i.d.*). A significant difference was also found between EOA and LOA receiving ondansetron, benefiting the former on drinks per day, drinks per drinking day and DrinC total score.

A larger-scale clinical trial, conducted by Johnson et al. [21], enrolling 283 patients showed that L/L-subjects receiving ondansetron significantly reduced their alcohol consumption, measured by drinks per drinking day and percentage of days abstinent as compared to placebo (respectively -1.62,  $p = 0.007$  and 11.27%,  $p = 0.023$ ).

In a small-scale study with 15 non-treatment seeking individuals, Kenna et al. [20] showed that patient with L/L genotype on the 5-HTTLPR promoter region of SLC6A4 (further referred as L/L-subjects) that were administered ondansetron (4  $\mu\text{g/kg}$  *b.i.d.*) for 3 weeks drank significantly less alcohol at an alcohol self-administration, compared to similar patients administered sertraline (200 mg per day).

Another clinical trial involving 77 patients, showed limited support that ondansetron may reduce drinking in non-treatment seeking L/L-subjects and was inconclusive in evaluating the effectiveness of sertraline in S/L or S/S-subjects ([26]). Further analysis ([27]) pinpointed gender differences as L/L women treated with ondansetron and S/L or S/S women treated with sertraline had significantly less drinks per drinking days and drank less at alcohol self-administration evaluations.

In the only trial taking place outside of North America, Corrêa Filho et al. [25] showed a significant reduction of heavy drinking days (7,8 % vs 11,7%,  $p=0.02$ ) but not of other measured outcomes.

**Craving and cue-induced craving** Three studies (Table 3) evaluated the impact of ondansetron on craving.

An analysis of the study population of Johnson et al. 2000a ([12]) found that the combination of ondansetron and naltrexone was significantly better than placebo at reducing craving among EOA [13].

Johnson et al. 2002 [15] showed that ondansetron at 4  $\mu\text{g/kg}$  *b.i.d.* was associated with a significant reduction in craving (measured by visual analog scale) compared to the placebo group, but only in EOA. In contrast, craving was significantly increased by ondansetron at 1  $\mu\text{g/kg}$  *b.i.d.* among LOA.

In a BOLD-MRI laboratory study, Myrick et al. [19] evaluated ventral striatum activation of AUD suffering people, treated for 7 days by either naltrexone, ondansetron, a combination of both or placebo, and "social-drinkers" (control group), when shown alcohol cues or neutral beverage cues. Ventral striatum activation was significantly reduced in the naltrexone, combination and social-drinkers groups. This was correlated with reduced craving scores in these groups. Ondansetron alone wasn't significantly effective in the reduction of ventral striatum activation nor craving. There was however a trend in the reduction of both.

**Mood disturbances** One study, presented in Table 4, showed that among EAO only, ondansetron significantly reduced mood disturbances as measured by the POMS scale ([30]).

**Moderators of treatment outcomes** Finally, five studies, presented in Table 5 didn't directly evaluate the efficacy of

ondansetron but provided useful information on prediction factors or safety and tolerability.

As ondansetron had previously been found useful in early onset alcoholism, Dawes et al. [18] evaluated ondansetron among adolescent in a prospective, open-label trial, which showed that ondansetron was safe and well tolerated in adolescents with alcohol dependence.

Roache et al. 2008 [17] compared the prediction capabilities of the EOA/LOA typology to the type A/type B typology precedently described by Babor et al 1992 [7][8](derived from Type I/II description by Cloninger et al. [6]). The A/B typology better described baseline severity of alcohol dependance but treatment response to ondansetron was significantly better predicted by the EOA/LOA typology.

Seneviratne et al. [23] produced some evidence that 5'-HTTLPR mRNA levels could be used as biomarker to evaluate treatment effectiveness in L/L-subjects treated with ondansetron.

Two studies analysed the population of Johnson et al. 2011 [21] and identified genotypes predicting treatment success. Johnson et al. 2013 [22] found 5 genotypes which presence predicted efficacy of ondansetron and which were present in a third of the population. Finally, Hou et al. [24] further worked on identifying ways of predicting effectiveness of ondansetron and found that data mining approaches, such as interaction trees and virtual twins could simplify subgroup identification while limiting statistical errors.

## Discussion

Overall, these studies show promising result for the use of ondansetron in AUD-suffering people. In particular subgroups, such as early-onset alcoholics, LL-genotypes for the SLA6A4 gene, or AUD-suffering people having one of the five genotypes identified in [22], ondansetron was shown to have a therapeutic effect which might be comparable or better than currently used medication. This paves the way to new discoveries and application in personalized medicine.

However, most of the studies have stringent inclusion criteria which greatly limits their external validity. Particularly, patients suffering from dual diagnosis or addicted to several substance (with the exception of nicotine) were excluded more often than not.

Moreover, ethical issues related to mass genotyping may prevent these type of information to be used for treatment prediction.

**Conclusion** Whereas growing evidence tends to suggest efficacy of ondansetron as a treatment of alcohol use disorder in particular genetic subgroups, new studies will be needed to fully conclude. In particular, there is a need for bigger studies evaluating long term changes in alcohol con-

sumption. These studies should also have less exclusion criteria to maximize their external validity. :

## References

- [1] Russell M. Hagan, Arthur Butler, Julia M. Hill, Christopher C. Jordan, Simon J. Ireland, and Michael B. Tyers. Effect of the 5-HT<sub>3</sub> receptor antagonist, GR38032f, on responses to injection of a neurokinin agonist into the ventral tegmental area of the rat brain. *European Journal of Pharmacology*, 138(2):303–305, 1987.
- [2] Kathleen A. Grant and James E. Barrett. Blockade of the discriminative stimulus effects of ethanol with 5-HT<sub>3</sub> receptor antagonists. *Psychopharmacology*, 104(4):451–456, 1991.
- [3] B. A. Johnson, G. M. Campling, P. Griffiths, and P. J. Cowen. Attenuation of some alcohol-induced mood changes and the desire to drink by 5-HT<sub>3</sub> receptor blockade: a preliminary study in healthy male volunteers. *Psychopharmacology*, 112(1):142–144, 1993.
- [4] B. A. Johnson, J. D. Roache, M. A. Javors, C. C. Di-Clemente, C. R. Cloninger, T. J. Prihoda, P. S. Bordnick, N. Ait-Daoud, and J. Hensler. Ondansetron for reduction of drinking among biologically predisposed alcoholic patients: A randomized controlled trial. *JAMA*, 284(8):963–971, 2000.
- [5] Henry R. Kranzler, Amira Pierucci-Lagha, Richard Feinn, and Carlos Hernandez-Avila. Effects of ondansetron in early- versus late-onset alcoholics: A prospective, open-label study. *Alcoholism: Clinical & Experimental Research*, 27(7):1150–1155, 2003.
- [6] C Robert Cloninger. A systematic method for clinical description and classification of personality variants: A proposal. *Archives of general psychiatry*, 44(6):573–588, 1987.
- [7] Thomas F. Babor, Michael Hofmann, Frances K. DelBoca, Victor Hesselbrock, Roger E. Meyer, Zelig S. Dolinsky, and Bruce Rounsaville. Types of Alcoholics, I: Evidence for an Empirically Derived Typology Based on Indicators of Vulnerability and Severity. *Archives of General Psychiatry*, 49(8):599–608, 08 1992.
- [8] Mark D. Litt, Thomas F. Babor, Frances K. DelBoca, Ronald M. Kadden, and Ned L. Cooney. Types of Alcoholics, II: Application of an Empirically Derived Typology to Treatment Matching. *Archives of General Psychiatry*, 49(8):609–614, 08 1992.



- [9] Vijoy K Varma, Debasish Basu, Anil Malhotra, Avneet Sharma, and Surendra K Mattoo. Correlates of early- and late-onset alcohol dependence. *Addictive behaviors*, 19(6):609–619, 1994.
- [10] Nassima Ait-Daoud, John D Roache, Michael A Dawes, Lei Liu, Xin-Qun Wang, Martin A Javors, Chamindi Seneviratne, and Bankole A Johnson. Can serotonin transporter genotype predict craving in alcoholism? *Alcoholism: Clinical and Experimental Research*, 33(8):1329–1335, 2009.
- [11] Edward M. Sellers, Tony Toneatto, Myroslava K. Romach, Gail R. Somer, Linda C. Sobell, and Mark B. Sobell. Clinical efficacy of the 5-HT<sub>3</sub> antagonist ondansetron in alcohol abuse and dependence. *Alcoholism: Clinical and Experimental Research*, 18(4):879–885, 1994.
- [12] Bankole A. Johnson, Nassima Ait-Daoud, and Thomas J. Prihoda. Combining ondansetron and naltrexone effectively treats biologically predisposed alcoholics: From hypotheses to preliminary clinical evidence. *Alcoholism: Clinical and Experimental Research*, 24(5):737–742, 2000.
- [13] N. Ait-Daoud, B. A. Johnson, T. J. Prihoda, and I. D. Hargita. Combining ondansetron and naltrexone reduces craving among biologically predisposed alcoholics: preliminary clinical evidence. *Psychopharmacology*, 154(1):23–27, 2001.
- [14] Nassima Ait-Daoud, Bankole A. Johnson, Martin Javors, John D. Roache, and Nursen A. Zanca. Combining ondansetron and naltrexone treats biological alcoholics: Corroboration of self-reported drinking by serum carbohydrate deficient transferrin, a biomarker. *Alcoholism: Clinical and Experimental Research*, 25(6):847–849, 2001.
- [15] Bankole A. Johnson, John D. Roache, Nassima Ait-Daoud, Nursen A. Zanca, and Madeline Velazquez. Ondansetron reduces the craving of biologically predisposed alcoholics. *Psychopharmacology*, 160(4):408–413, 2002.
- [16] Bankole A. Johnson, Nassima Ait-Daoud, Jennie Z. Ma, and Yanmei Wang. Ondansetron reduces mood disturbance among biologically predisposed, alcohol-dependent individuals. *Alcoholism: Clinical & Experimental Research*, 27(11):1773–1779, 2003.
- [17] John D. Roache, Yanmei Wang, Nassima Ait-Daoud, and Bankole A. Johnson. Prediction of serotonergic treatment efficacy using age of onset and type a/b typologies of alcoholism. *Alcoholism, Clinical and Experimental Research*, 32(8):1502–1512, 2008.
- [18] Michael A. Dawes, Bankole A. Johnson, Nassima Ait-Daoud, Jennie Z. Ma, and Jack R. Cornelius. A prospective, open-label trial of ondansetron in adolescents with alcohol dependence. *Addictive Behaviors*, 30(6):1077–1085, 2005.
- [19] Hugh Myrick, Raymond F Anton, Xingbao Li, Scott Henderson, Patrick K Randall, and Konstantin Voronin. Effect of naltrexone and ondansetron on alcohol cue-induced activation of the ventral striatum in alcohol-dependent people. *Archives of general psychiatry*, 65(4):466–475, 2008.
- [20] George A. Kenna, William H. Zywiak, John E. McGeary, Lorenzo Leggio, Chinatsu McGeary, Shirley Wang, Andrea Grew, and Robert M. Swift. A within-group design of nontreatment seeking 5-HTTLPR genotyped alcohol-dependent subjects receiving ondansetron and sertraline. *Alcoholism, Clinical and Experimental Research*, 33(2):315–323, 2009.
- [21] Bankole A. Johnson, Nassima Ait-Daoud, Chamindi Seneviratne, John D. Roache, Martin A. Javors, Xin-Qun Wang, Lei Liu, J. Kim Penberthy, Carlo C. DiClemente, and Ming D. Li. Pharmacogenetic approach at the serotonin transporter gene as a method of reducing the severity of alcohol drinking. *The American Journal of Psychiatry*, 168(3):265–275, 2011.
- [22] Bankole A. Johnson, Chamindi Seneviratne, Xin-Qun Wang, Nassima Ait-Daoud, and Ming D. Li. Determination of genotype combinations that can predict the outcome of the treatment of alcohol dependence using the 5-HT<sub>3</sub> antagonist ondansetron. *The American Journal of Psychiatry*, 170(9):1020–1031, 2013.
- [23] Chamindi Seneviratne and Bankole A. Johnson. Serotonin transporter genomic biomarker for quantitative assessment of ondansetron treatment response in alcoholics. *Frontiers in Psychiatry*, 3:23, 2012.
- [24] Jue Hou, Chamindi Seneviratne, Xiaogang Su, Jeremy Taylor, Bankole Johnson, Xin-Qun Wang, Heping Zhang, Henry R. Kranzler, Joseph Kang, and Lei Liu. Subgroup identification in personalized treatment of alcohol dependence. *Alcoholism, Clinical and Experimental Research*, 39(7):1253–1259, 2015.
- [25] João Maria Corrêa Filho and Danilo Antonio Baltieri. A pilot study of full-dose ondansetron to treat heavy-drinking men withdrawing from alcohol in Brazil. *Addictive Behaviors*, 38(4):2044–2051, 2013.
- [26] George A. Kenna, William H. Zywiak, Robert M. Swift, John E. McGeary, James S. Clifford, Jessica R. Shoaff, Cynthia Vuittonet, Samuel Fricchione, Michael Brickley, Kayla Beaucage, Carolina L. Haass-Koffler,

- and Lorenzo Leggio. Ondansetron reduces naturalistic drinking in nontreatment-seeking alcohol-dependent individuals with the LL 5'-HTTLPR genotype: a laboratory study. *Alcoholism, Clinical and Experimental Research*, 38(6):1567–1574, 2014.
- [27] George A. Kenna, William H. Zywiak, Robert M. Swift, John E. McGeary, James S. Clifford, Jessica R. Shoaff, Samuel Fricchione, Michael Brickley, Kayla Beaucage, Carolina L. Haass-Koffler, and Lorenzo Leggio. Ondansetron and sertraline may interact with 5-HTTLPR and DRD4 polymorphisms to reduce drinking in non-treatment seeking alcohol-dependent women: Exploratory findings. *Alcohol*, 48(6):515–522, 2014.
- [28] Linda C Sobell and Mark B Sobell. *Timeline follow-back: user's guide*. Addiction Research Foundation, 1996.
- [29] Raymond F Anton, Darlene H Moak, and Patricia K Latham. The obsessive compulsive drinking scale: A new method of assessing outcome in alcoholism treatment studies. *Archives of general psychiatry*, 53(3):225–231, 1996.
- [30] Douglas McNair, Maurice Lorr, and Leo Droppleman. Profile of mood states (poms). 1989.