


Accepted Manuscript

Title: Executive function performance in obesity and overweight individuals: A meta-analysis and review

Authors: Yingkai Yang, Grant S. Shields, Cheng Guo, Yanling Liu



PII: S0149-7634(17)30479-7
DOI: <https://doi.org/10.1016/j.neubiorev.2017.11.020>
Reference: NBR 3011

To appear in:

Received date: 5-7-2017
Revised date: 8-9-2017
Accepted date: 30-11-2017

Please cite this article as: Yang, Yingkai, Shields, Grant S., Guo, Cheng, Liu, Yanling, Executive function performance in obesity and overweight individuals: A meta-analysis and review. *Neuroscience and Biobehavioral Reviews* <https://doi.org/10.1016/j.neubiorev.2017.11.020>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Executive function performance in obesity and overweight individuals:

A meta-analysis and review

Yingkai Yang^{1*}, Grant S. Shields², Cheng Guo^{1*}, and Yanling Liu¹

¹The Lab of Mental Health and Social Adaptation, Faculty of Psychology, Research Center of Mental Health Education, Southwest University, Chongqing, China;

²Department of Psychology, University of California, Davis, CA, USA.

*Corresponding Author:

Cheng Guo, Faculty of Psychology, Southwest University, No.2 Tiansheng Street, Beibei District, 400715 Chongqing, China. E-mail: guochen@swu.edu.cn;

Yingkai Yang, Faculty of Psychology, Southwest University, No.2 Tiansheng Street, Beibei District, 400715 Chongqing, China. E-mail: yykjiyisui pian@gmail.com;

Abstract Word Count: 170

Body Word Count: 12733

Number of Figures: 11

Number of Tables: 5

Supplementary Materials: 1 table, 2 page of text

Highlights

- We conducted a meta-analysis of associations between obesity/overweight and various executive functions.
- Obese participants showed broad impairments on executive function.
- Overweight participants showed significant deficits in inhibition and working memory performances.
- Measures of executive function moderated effects of obesity on working memory and decision-making.

Abstract

Prior research has suggested that obesity/overweight may be associated with deficits in executive function. If true, this has important clinical implications. In this review, we synthesize the current literature by conducting a meta-analysis of studies comparing executive functions in overweight/obese individuals to normal weight controls. We identified 72 studies—with 4,904 overweight/obese participants—that met our inclusion criteria. Effect sizes were analyzed using the robust variance estimation random effects meta-regression technique. It was found that obese participants showed broad impairments on executive function, including on tasks primarily utilizing inhibition, cognitive flexibility, working memory, decision-making, verbal fluency, and planning; overweight participants only showed significant deficits in inhibition and working memory. The only moderator of effects of obesity to emerge significant was

the task used to assess the respective executive function, which moderated effects of obesity on working memory and decision-making. There were not enough studies of overweight individuals to make strong claims about moderating effects in those studies.

In sum, current evidence supports the existence of broad executive function deficits in obese individuals, and inhibition and working memory deficits in overweight individuals.

Keywords: obesity; overweight; executive functions; meta-analysis

1. Introduction

The prevalence of obesity and overweight is rising problematically in developed and developing nations worldwide (Ng et al., 2014). The World Health Organization reported that in 2014, more than 600 million adults were obese and over 1.9 billion were overweight (World Health Organization, 2017). This fact has far-reaching and costly implications, because excessive weight contributes to the development of numerous diseases, including cardiovascular disease (Lavie, Milani, & Ventura, 2009), diabetes (Mokdad et al., 2003), and some cancers (Kyrgiou et al., 2017), and it is a risk

factor for psychiatric disorders such as depression and anxiety (Luppino, 2010). Not surprisingly, excess weight has become a cause of growing health care costs (Withrow, & Alter, 2011) and accounts for over 2.8 million deaths per year (World Health Organization, 2017).

While it is clear that obesity/overweight correlate with poor mental and physical health, recent studies have linked overweight and obesity to poorer cognitive functioning. In particular, research has suggested that individual cognitive performance declines with increases in body mass index (BMI) (Bocarsly et al., 2015; Smith, Hay, Campbell, & Trollor, 2011; Vainik, U., Dagher, Dubé, & Fellows, 2013). These deficits can be observed throughout life, from childhood to late adulthood (Liang, Matheson, Kaye, & Boutelle, 2014; Smith et al., 2011). More recently, it has been proposed that the cognitive processes collectively known as executive function may be particularly vulnerable to weight-related impairments (Appelhans, 2009; Bartholdy, Dalton, O'Daly, Campbell, & Schmidt, 2016; Jansen, Houben, & Roefs, 2015; Fitzpatrick, Gilbert, & Serpell, 2013; Nederkoorn, Braet, Van Eijs, Tanghe, & Jansen, 2006).

1.1 Executive function

Although executive function has been defined in different ways, these definitions all share the idea that executive functions are the higher cognitive processes that enable forethought and goal-directed action (Banich, 2009; Diamond, 2013).

According to an influential model of executive function (Miyake, Friedman, Emerson, Witzki, Howerter, & Wager, 2000; Miyake, & Friedman, 2012), there are three key aspects of EF: (1) inhibition, which refers to the ability to suppress impulsive or automatic (prepotent) responses, (2) cognitive flexibility, which refers to the ability to shift attention as well as mental sets or rules when situationally appropriate, and (3) working memory, which refers to the ability to monitor the relevance of incoming stimuli and update information in memory as required.

Although inhibition, cognitive flexibility, and working memory are important aspects of EF, they may not be the only components (Snyder, 2013). Indeed, several other EF domains have been well defined in the literature (Collins, & Koechlin, 2012; Lezak, 2012; Suchy, 2009; Testa, Bennett, & Ponsford, 2012), including (a) decision making (e.g., Rangel, Camerer, & Montague, 2008), defined as the cognitive process that occurs whenever an individual has to make a choice from several alternative

possibilities, (b) verbal fluency (e.g., Troyer, Moscovitch, & Winocur, 1997), defined as the ability to generate as many words as possible from a semantic category (or that start with certain letters) in a given time, and (c) planning (e.g., Lezak, 2012), defined as formulating, evaluating and selecting a sequence of thoughts and actions to achieve a goal.

1.2 Theories of obesity/overweight with executive function

Theories of the association between obesity/overweight and executive function agree that excessive weight should be associated with impairments in executive function (Appelhans, 2009; Castanon, Lasselin, & Capuron, 2014; Guillemot-Legris, & Muccioli, 2017; Miller, & Spencer, 2014; Nederkoorn et al., 2006; Spyridaki, Avgoustinaki, & Margioris, 2016; Stoeckel et al., 2017), although the exact reason for this association is still unclear.

Excessive weight may exert detrimental effects on cognition, including executive function. What's more, it is gradually becoming clearer that these deleterious effects of adiposity on cognition are not solely mediated by the commonly accepted clinical consequences (e.g., type 2 diabetes, hypoglycemic episodes, hyperlipidemia)

of adiposity (Smith et al., 2011; Spyridaki, Avgoustinaki, & Margioris, 2016). Recent research suggests that body mass and executive function deficits can be indirectly associated via obesity-induced activation of innate immunity which directly caused low-grade inflammation in obesity (Bourassa, & Sbarra, 2017; D O'Brien, Hinder, Callaghan, & Feldman, 2017; Lasselin et al., 2016). This association is furthered by a newly proposed model, the immunologic model of self-regulatory failure, which suggested that immune system activity—especially components of the immune system involved in inflammation—impair executive function (Shields, Moons, & Slavich, 2017).

As fundamental and important aspects for lifestyle habits, it is also possible that differences on executive function could predispose individuals to excessive weight. The dual process model argues that much of our behavior is determined by an interaction between the impulsive system and the executive control system (Hofmann, Friese, & Strack, 2009; Hofmann, Schmeichel, & Baddeley, 2012; Strack & Deutsch, 2004). In the case of excessive weight, coupled with a strong automatic approach response to high-calorie food and food cues, people who show low levels of executive control are

particularly susceptible to obesity-related behaviors and outcomes (e.g., increased intake of fatty foods, weight gain), whereas those with effective cognitive control may be protected (Appelhans, 2009).

More recently, using non-invasive brain stimulation procedures, studies have found that the dorsolateral prefrontal cortex (dlPFC)—a major neural area implicated in executive function (Nee, Wager, & Jonides, 2007)—plays a causal role in the inhibition of high-calorie food craving and consumption (Hall, & Vincent, 2017). For instance, in a study examining the effects of multi-session of transcranial direct current stimulation (tDCS) targeting the right dlPFC in 30 normal and overweight young adults, significant food craving reductions in the active stimulation condition (but not sham) were observed over the 5-day treatment interval and the 30-day follow up assessment (Ljubisavljevic, Maxood, Bjekic, Oommen, & Nagelkerke, 2016). These data further suggested that executive function—or at least neural regions supporting it—may play a causal role in the development of abnormal eating behavior that increases excessive weight risk (Hall, 2016).

1.3. Evidence for impaired executive function in obesity/overweight

Despite theories converged in positing that obesity/overweight should be associated with greater impairments in executive functioning, results have been inconsistent in studies examining these associations. Many studies have indeed found significant deficits in executive function when comparing obese individuals to normal weight participants (e.g., Nederkoorn et al., 2006; Kamijo et al., 2014); however, other studies have reported no significant differences in executive function between those groups (e.g., Ariza et al., 2012; Delgado-Rico, Soriano-Mas, Verdejo-Román, S Río-Valle, & Verdejo-García, 2013). Researchers have consequently reached a wide range of conclusions about the relationship between excess weight and executive function, ranging from pronounced and broad impairments on neuropsychological measures of executive functioning (e.g., Cohen, Yates, Duong, & Convit, 2011) to no apparent impairments on some aspects of executive function (e.g., Lawyer, Boomhower, & Rasmussen, 2015). In sum, although weight-related impairments in inhibition, cognitive flexibility, working memory, decision-making, verbal fluency, planning are often reported, a systematic meta-analysis is needed to assess the consistency and magnitude of such deficits.

A number of informative narrative and systematic reviews have already been conducted to summarize empirical findings regarding the effects of obesity/overweight on executive function performance. However, these reviews either (a) did not apply meta-analytical techniques to quantitatively synthesize the data (e.g., Fitzpatrick, Gilbert, & Serpell, 2013; Prickett, Brennan, & Stolwyk, 2015; Stojek, & MacKillop, 2017; Smith et al., 2011; Vainik et al., 2013), (b) solely focused on one outcome measure (e.g., the stop signal task) or type (e.g., cognitive flexibility) of executive function (e.g., Amlung, Petker, Jackson, Balodis, & MacKillop, 2016; Bartholdy et al., 2016; Lavagnino, Arnone, Cao, Soares, & Selvaraj, 2016; McClelland, Dalton, Kekic, Bartholdy, Campbell, & Schmidt, 2016; Rotge, Poitou, Fossati, Aron-Wisnewsky, & Oppert, 2017; Wu, Brockmeyer, Hartmann, Skunde, Herzog, & Friederich, 2014, 2016), (c) only considered studies with children and adolescents (e.g., Liang et al., 2014), and/or (d) only focused on weight disorder without distinguishing between obesity and overweight (e.g., Wu et al., 2014). In addition, the number of neuropsychological studies on executive function in excess weight is growing rapidly; as such, prior meta-analytic results likely need updated to reflect the current literature, especially given

increased power to detect moderating effects.

1.4 The present meta-analysis

In the current study, we conducted a quantitative meta-analysis of existing studies examining executive functions in obesity/overweight individuals as compared with normal weight comparison groups. Such an analysis is important since it allowed for a combination of effect sizes across studies and provided a more powerful estimate of true population differences.

The mixed results across studies may be due to the diversity of participant or task-related characteristics—that is, potential moderators. Therefore, a secondary aim of this meta-analysis was to identify moderators (whenever possible) of differences in executive functions between obese/overweight people and normal weight controls. The moderators we examined include age, percent of participants in the sample that were female, BMI (in adult samples), and measures of executive functions.

We considered the aforementioned factors as potential moderators mainly because previous evidence suggested that effect sizes diverged when those study characteristics differed. More specifically, previous meta-analysis on altered decision-

making with excess weight revealed nonsignificant effects of excess weight in adolescent samples (Wu et al., 2016). In addition, some researchers have obtained nonsignificant effect sizes when using some measures (e.g., stop signal task) of executive functions (e.g., Lawyer et al., 2015; Schiff et al., 2016) or examining groups that only included males (e.g., Weller, Cook, Avsar, & Cox, 2008). Finally, there is also some evidence that executive function impairments are larger in obese individuals with higher BMIs (e.g., Perpiñá, Segura, & Sánchez-Reales, 2016; Fagundo et al. 2016). Therefore, given the above, we examined age, percentage of female participants, BMI, and measures of executive function as potential moderators of the differences in executive functions between obese/overweight people and normal weight controls.

2. Method

2.1. Study Selection and Inclusion Criteria

2.1.1. Literature review. To obtain studies for use in the meta-analysis, a topic search in the databases PubMed, ISI Web of Knowledge, and PsycINFO was conducted for all papers published until February 20, 2017 (see supplementary material for a list of keywords). In this search, PubMed returned 9120 results, ISI Web of Knowledge

returned 7196 results, and PsycINFO returned 907 results. Abstracts of articles were reviewed and the full text of an article was read whenever a paper's title or abstract indicated that the study might be relevant to analyses. In addition, to help ensure that all studies on this topic were included, references from relevant articles were reviewed, and studies that were potentially relevant were examined from those references. Finally, ProQuest Dissertations and Theses was also searched to identify unpublished dissertations. Additionally, a Google search was conducted to further identify unpublished studies. Figure 1 outlines the detailed study selection procedure.

2.1.2. Inclusion criteria. Studies were incorporated into this meta-analysis if they (1) examined human participants (2) compared at least one group of obese or overweight individuals (minimum age 6 years) to a normal weight control group; (3) used at least one task known or shown to depend upon or assess executive function; and (4) provided data or statistical information that allowed for effect size calculation. In samples of adult participants, an obese group was defined as an average BMI of 30 kg/m² or above, and an overweight group as an average BMI between 25–30 kg/m². In samples of children/adolescents, obesity was defined as a BMI percentile of 95th or

above, and overweight as a BMI between the 85th and 95th percentile; if BMI percentile was not reported, international cutoff points of BMI were used for defining obesity and overweight (Cole, Bellizzi, Flegal, & Dietz, 2000). In addition, we excluded studies which explicitly indicated that all their overweight/obese participants had a known psychological/psychiatric disorder (e.g., depression, schizophrenia, binge eating disorder).

2.1.3. Selected studies. Our search and study inclusion criteria led to the incorporation of 72 studies, 71 of which were published in peer-reviewed papers, and 1 of which was an unpublished thesis. Of these 72 studies, 44 assessed differences between obese/overweight individuals and normal weight on inhibition, 25 assessed differences in cognitive flexibility, 22 assessed differences in working memory, 29 assessed differences in decision-making, 9 assessed differences in verbal fluency, and 8 assessed differences in planning.

2.2. Coding of Variables

Executive function tasks were coded as assessing inhibition, cognitive flexibility, working memory, decision-making, verbal fluency, or planning based upon

previous empirical or theoretical evidence suggesting that a given task primarily utilized the particular coded executive function. See the supplementary material table S1 for a complete description of task coding. This list is not meant to be exhaustive of all executive functions, but rather of the range of neuropsychological measures of executive function included in the obesity/overweight literature.

Within the obese or overweight groups, mean age, percentage female participants, and BMI were coded as continuous variables. Measures of executive function were contrast coded and submitted to F tests (as described below) to examine differences.

The methodological quality of included studies was assessed through a linear combination of factors that may influence obtained differences in executive function between obese/overweight individuals and normal weight individuals such that a higher score indicated better precision in assessing potential executive function differences.

Where exclusion of age, gender, IQ/socioeconomic status (SES) and education differences, illnesses, medication use, are all dummy-coded as 1 for excluded and 0 for not excluded, the equation is as follows: Study quality = (Age Difference Excluded [0–

1]) + (Gender Difference Excluded [0–1]) + (Education Difference Excluded [0–1]) + (IQ/ SES Difference Excluded [0–1]) + (Illnesses Excluded [0–1]) + (Medication Excluded [0–1]) (Differences between the groups were analyzed using *t*-test or χ^2 tests, $p > 0.05$ means difference excluded).

2.3. Statistical Analyses

The effect size measure of interest was the standardized mean difference between obese/overweight and normal weight groups. Hedges' *g*, rather than Cohen's *d*, was used as the effect size for analysis, given that the former is a relatively unbiased estimate of the population standardized mean difference effect size while the latter is a biased estimate. Whenever possible, we firstly calculated Cohen's *d* from the means, standard deviations, and sample sizes presented in the article. If means and standard deviations were not reported and the design was between-studies, we used *t* or one-way *F* statistics—or *p* values resulting from tests of those two statistics—to calculate the effect size. If none of these statistics were reported, that study was excluded from analysis. Then, all Cohen's *d* effect sizes were transformed into Hedges' *g* effect sizes and used for analysis (Borenstein, Hedges, Higgins, Rothstein, 2009).

Given the multifaceted nature of executive function, most studies often report more than one outcome (e.g., errors of commission and errors of omission on the go/no-go task). Multiple outcomes are a problem for conventional meta-analytic methods, as averaging effect sizes within studies without accounting for their correlations can alter or obscure true effect size estimates (Scammacca, Roberts, & Stuebing, 2014). Thus, we employed the meta-analytic technique of robust variance estimation, a random-effects meta-regression that can account for dependence between effect size estimates (Hedges, Tipton, & Johnson, 2010; Tanner-Smith & Tipton, 2014). This technique robustly estimates effect size weights and standard errors for the given effects, allowing for multiple outcomes within studies (Hedges, Tipton, & Johnson, 2010). We employed the `robu()` function of the `robumeta` package, version 2.0, in **R**, version 3.4.0, to conduct these analyses using the correlated weights given by Hedges, Tipton and Johnson (2010) with our primary analyses using the small sample corrections suggested by Tipton (2015). To account for dependency, ρ was set to the recommended 0.80 (Tanner-Smith & Tipton, 2014). To conduct F tests on models with contrast-coded coefficients, we used the `Wald_test()` function in the `clubSandwich` package,

version 0.2.2, in **R**, with the bias-reduced linearization adjustment for clustered standard errors and degrees of freedom estimated with Hotelling's T_Z^2 method.

Degrees of freedom for all primary analyses were estimated using the Satterwaite approximation, where $df = 2/cv^2$ and cv represents the coefficient of variation, as simulation studies have indicated that this method of estimating degrees of freedom is most analytically valid with study set sizes under 40 using the RVE meta analytic technique (Tipton, 2015).

Finally, we used the procedures described by Viechtbauer and Cheung (2010) to derive extreme outliers (identified by inspecting z-score of the standardised residuals) and influential studies (identified by inspecting Cook's distance plots). If the z-score of the standardized residuals exceeded 1.96, the study was deemed to be an outlier. And if Cook's distance plots showed the outlier exert significant influence on the results, the outlier was excluded and only results from the meta-analysis without the outlier were reported in full.

For all of the following analyses, a negative effect size means that obese or overweight group performed worse compared to the normal weight control group,

whereas a positive effect size indicates that the obese or overweight group performed better than normal weight individuals. In addition, because the outcome in these analyses is the standardized mean difference between groups (the effect size), a significant moderator means that the effect size estimate depends upon levels of that variable. We only conducted meta-regression for outcomes in which there are at least 10 samples to 1 covariate (Borenstein et al., 2009)

3. Results

3.1. Preliminary Analyses

3.1.1. Study characteristics. The final sample consisted of 72 studies (i.e., total $m = 72$). Of those 72 studies, 65 examined differences in executive function between obese and normal weight individuals, whereas 24 studies examined differences in executive function between overweight and normal weight individuals. There were 183 effect sizes (i.e., total $k = 183$) between obese participants and normal weight participants, and 69 effect sizes (i.e., total $k = 69$) between overweight participants and normal weight participants. The number of effect sizes per study that we obtained is relatively common in social science research (e.g., Shields, Sazma, & Yonelinas, 2016;

Yang, Cao, Shields, Teng, & Liu, 2017).

Differences in inhibition were examined in 39 studies ($k = 50$) with 1,782 obese participants and in 10 studies ($k = 20$) with 641 overweight participants. Differences in cognitive flexibility were examined in 24 studies ($k = 33$) with 1,145 obese participants, and in 5 studies ($k = 6$) with 382 overweight participants. Differences in working memory were examined in 20 studies ($k = 35$) with 785 obese participants and in 8 studies ($k = 17$) with 406 overweight participants. Differences decision-making were examined in 27 studies ($k = 46$) with 1,205 obese participants and in 6 studies ($k = 18$) with 210 overweight participants. Differences in verbal fluency were examined in 9 studies ($k = 12$) with 927 obese participants and in 3 studies ($k = 4$) with 897 overweight participants. Effect sizes on planning were examined in 6 studies ($k = 7$) with 359 obese participants, and in 3 studies ($k = 4$) with 348 overweight participants.

Table 1 presents each study and its characteristics.

3.1.2. Assessment of publication bias. Egger's test (Egger, Smith, Schneider, & Minder, 1997) for funnel plot asymmetry was used to conduct the publication bias test on each executive function individually. In assessment of obesity (Figure 2), the test for

publication bias returned nonsignificant for inhibition, $t(38) = -0.49$, $p = 0.626$; cognitive flexibility, $t(22) = -0.01$, $p = 0.995$; working memory, $t(18) = -0.32$, $p = 0.752$; decision-making, $t(27) = -0.78$, $p = 0.440$; verbal fluency $t(7) = -0.20$, $p = 0.849$; and planning, $t(4) = -0.41$, $p = 0.701$. In assessment of overweight, the test for publication bias returned nonsignificant for inhibition (Figure 3), $t(10) = -0.50$, $p = 0.630$, cognitive flexibility, $t(4) = 0.43$, $p = 0.690$, decision making, $t(4) = -1.27$, $p = 0.274$; verbal fluency, $t(1) = 0.57$, $p = 0.671$; and planning, $t(2) = -1.63$, $p = 0.245$. The test indicated significant publication bias in studies examining working memory (Figure 3), $t(7) = 3.43$, $p = .011$; notably, however, a trim and fill analysis indicated that the two missing effect sizes were associated with larger effects of overweight on working memory than the average published study. Including these estimated missing working memory studies increased the impairing effect of overweight on working memory by $B = -0.0138$. Importantly, these publication bias results did not differ if the removed, outlying effect sizes were included in analyses, indicating that removing these effect sizes did not hide any potential publication bias. Therefore, these results indicate that any effects observed in this meta-analysis are unlikely to be due to publication bias.

3.2. Primary Analyses of Obesity Effects

3.2.1 Testing overall differences in executive functions. We first tested whether potential differences in executive functions between obese and normal weight individuals were broad and nonspecific or whether any potential differences varied by executive function domain. Results indicated that differences in executive functions between obese and normal weight individuals did not differ across the domains of inhibition, cognitive flexibility, working memory, decision-making, verbal fluency, and planning, $F(5, 16.2) = 0.49, p = .78$. However, a lack of overall differences may mask varying effects of moderators on individual executive functions. Thus, we present analyses of each executive function below.

3.2.2. Inhibition. The analysis of studies examining differences between obese and normal weight participants on measures of inhibition ($m = 40, k = 50$) revealed a significant overall effect, $g^+ = -0.363, t(37.2) = -6.62, p < .001, 95\% \text{ CI } [-0.473, -0.252]$ (Figure 4), such that obese participants performed worse on inhibition tasks. There was low heterogeneity across these studies' effects, $\tau^2 = 0.069$, indicating that the impaired performance on inhibition tasks was relatively consistent across various

conditions. However, in the interest of examining potential factors that may contribute to some inconsistencies in studies examining effects of obesity on inhibition, we conducted moderator analyses of these effects, described below.

Moderator analyses showed that age, percentage of female participants, and BMI did not significantly moderate effects of obesity on inhibition (see Table 2). Similarly, the task used to assess inhibition did not moderate effects of obesity on inhibition, $F(3, 16.9) = 0.25, p = .862$ (see Table 2). Nine different tasks were used to assess inhibition in the studies included in our analyses of obesity effects. The most frequently used tasks were the Stroop task, the Stop Signal Task, and the Go/No-Go. The effect sizes comparing obese people to normal weight people across these tasks were comparable (see Table 2). Thus, we did not find evidence for significant moderation of effects of obesity on inhibition by age, sex, BMI, or task used to assess inhibition.

3.2.3. Cognitive flexibility. The analysis of studies examining differences between obese and normal weight participants on measures of cognitive flexibility ($m = 24, k = 33$) revealed a significant overall effect, $g^+ = -0.369, t(21) = -5.32, p < .001$,

95% CI [-0.513, -0.224] (Figure 5), such that obese participants performed worse than normal weight participants on tasks assessing cognitive flexibility. There was low heterogeneity across these studies' effects, $\tau^2 = 0.059$, indicating that the obesity-related performance impairment on tasks requiring cognitive flexibility was relatively consistent across various conditions.

Moderator analyses showed that age, percentage of female participants, and BMI did not significantly moderate effects of obesity on cognitive flexibility (Table 3). In addition, measures of cognitive flexibility did not significantly influence the effect sizes, $F(2, 12.6) = 0.92$, $p = .425$ (see Table 3). Nine different tasks were used to assess overweight/obesity in the studies examining effects of obesity included in our analyses. The most frequently used tasks were the Trail Making Test (TMT), and the Wisconsin Card Sorting Test (WCST). Effect sizes comparing obese to normal weight individuals were similar across these tasks (Table 2). Thus, we did not find evidence for significant moderation of effects of obesity on cognitive flexibility by age, sex, BMI, or task used to assess cognitive flexibility.

3.2.4. Working memory. The analysis of studies examining differences

between obese and normal weight participants on measures of working memory ($m = 20, k = 35$) revealed a significant overall effect, $g^+ = -0.332, t(17.9) = -4.48, p < .001$, 95% CI $[-0.488, -0.177]$ (Figure 6), such that obese participants performed worse than healthy weight participants on tasks utilizing working memory. There was low heterogeneity across these studies' effects, $\tau^2 = 0.071$, indicating that the obesity-related performance impairment of tasks requiring working memory was relatively consistent across various conditions.

Moderator analyses showed that age, percentage of female participants, and BMI did not significantly moderate effects of obesity on working memory (Table 4). However, the task used to assess working memory significantly moderated effects of obesity on working memory, $F(2, 6.8) = 6.98, p = .023$ (Table 4). Ten different tasks were used to assess working memory in studies comparing the working memory of obese to normal weight individuals. The most frequently used tasks were the Digit span task, and the Letter–number sequencing task. Six studies included the Digit span task, and performance on this task was significantly worse in obese than healthy weight individuals, $g^+ = -0.346, 95\% \text{ CI } [-0.542, -0.149]$. Four studies included the Letter–

number sequencing task, but performance on this task did not differ between obese and normal weight individuals, $g^+ = 0.026$, 95% CI $[-0.166, 0.218]$. Other working memory tasks, used in a total of 12 studies, showed an obesity-related performance impairment similar to the digit span, $g^+ = -0.478$, 95% CI $[-0.663, -0.292]$. As might be clear, the moderating effect of the measure used to assess working memory was driven by differences in the effect of obesity on the letter–number sequencing task compared to effects of obesity on other tasks, $B = 0.469$, $t(4.4) = 3.99$, $p = .013$. Thus, although relatively consistent across conditions, the effect of obesity on working memory is moderated by the task used to assess working memory, with less of an effect on the Letter–number sequencing task.

3.2.5. Decision-making. The analysis of studies examining differences between obese and healthy weight individuals in measures of decision-making ($m = 29$, $k = 46$) revealed a significant overall effect, $g^+ = -0.441$, $t(25.7) = -6.82$, $p < .001$, 95% CI $[-0.574, -0.308]$ (Figure 7), such that obese participants performed worse than healthy weight participants on measures of decision-making. There was low heterogeneity across these studies' effects, $\tau^2 = 0.060$, indicating that the altered performance on

decision-making is relatively consistent across various conditions.

Moderator analyses showed that age, percentage of female participants, and BMI did not significantly influence the pooled effect sizes (Table 5). However, the task used to assess decision-making significantly moderated effects of obesity on decision-making, $F(2, 13.6) = 4.51, p = .032$ (Table 5). Seven different tasks were used to assess decision-making in studies examining differences in decision-making between obese and healthy weight individuals. The most frequently used tasks were the Delay discounting task, and the Iowa gambling task. Eleven studies included the Delay discounting task, and obese participants performed worse on this task than healthy weight participants, $g^+ = -0.416$, 95% CI $[-0.602, -0.231]$. Twelve studies included Iowa gambling task, and obese participants performed worse on this task than healthy weight participants, $g^+ = -0.580$, 95% CI $[-0.815, -0.344]$. Seven studies included other measures of decision-making. Although obese participants performed worse on these other decision-making tasks than healthy weight participants, $g^+ = -0.200, t(4.2) = -2.87, p = .044$, the difference between obese and healthy weight participants in these other decision-making tasks was not as large as the difference in the Delay discounting

and Iowa gambling tasks, $B = 0.352$, $t(7.8) = 2.97$, $p = .019$.

3.2.6. Verbal fluency. The analysis of studies examining differences between obese and healthy weight participants in verbal fluency ($m = 9$, $k = 12$) revealed a significant overall effect, $g^+ = -0.308$, $t(5.7) = -3.90$, $p = .009$, 95% CI $[-0.504, -0.112]$ (Figure 8), such that obese participants performed worse on measures of verbal fluency than healthy weight participants. There was low heterogeneity across these effects, $\tau^2 = 0.023$, indicating that the obesity-related performance impairment in verbal fluency was relatively consistent across various conditions. There were not enough studies or task variability within studies examining of effects of obesity on verbal fluency to conduct reliable moderator analyses.

3.2.7. Planning. The analysis of studies examining differences between obese and healthy weight participants in planning ($m = 6$, $k = 7$) revealed a significant overall effect, $g^+ = -0.346$, $t(4.5) = -3.08$, $p = .031$, 95% CI $[-0.643, -0.048]$ (Figure 9), such that obese participants performed worse than healthy weight participants on tasks assessing planning. There was low heterogeneity across these studies' effects, $\tau^2 = 0.036$, indicating that the impaired performance on planning is relatively consistent

across various conditions. There were not enough studies or task variability within studies examining of effects of obesity on planning to conduct reliable moderator analyses.

3.3. Primary Analyses of Overweight Effects

We first tested whether potential differences in executive functions between overweight and normal weight individuals were broad and nonspecific or whether any potential differences varied by executive function domain. Results indicated that differences in executive functions between overweight and normal weight individuals did not differ across the domains of inhibition, cognitive flexibility, working memory, decision-making, verbal fluency, and planning, $F(5, 4.0) = 0.70, p = .655$. Nonetheless, we present analyses of each executive function below.

The analysis of studies examining effects of overweight on inhibition ($m = 10, k = 20$) revealed a significant overall effect, $g^+ = -0.234, t(9.8) = -2.56, p = .029, 95\% \text{ CI } [-0.438, -0.030]$ (Figure 10), such that overweight participants performed worse on measures of inhibition than healthy weight participants. Similarly, the analysis of studies examining effects of overweight on working memory ($m = 8, k = 17$) also

revealed a significant overall effect, $g^+ = -0.132$, $t(3.8) = -5.51$, $p = .006$, 95% CI $[-0.199, -0.064]$ (Figure 11), such that overweight participants performed worse on measures of working memory than healthy weight participants. There were not enough studies comparing executive functions of overweight to healthy weight participants to conduct moderator analyses.

Nonsignificant overall effects were founded when analyzing effect sizes on cognitive flexibility ($m = 6$, $k = 6$), $g^+ = -0.096$, $t(3.0) = -1.52$, $p = .227$, 95% CI $[-0.299, 0.107]$, decision-making ($m = 6$, $k = 18$), $g^+ = -0.030$, $t(4.4) = -0.271$, $p = .838$, 95% CI $[-0.397, 0.337]$, verbal fluency ($m = 3$, $k = 4$), $g^+ = -0.007$, $t(1.3) = -0.07$, $p = .952$, 95% CI $[-0.739, 0.725]$, and planning ($m = 3$, $k = 4$), $g^+ = -0.323$, $t(2.9) = -1.81$, $p = .171$, 95% CI $[-0.904, 0.258]$. However, because of the small study set sizes, it is difficult to interpret these nonsignificant effects, especially in light of the nonsignificant F test examining differences in effects of overweight on executive functions.

4. Discussion

From resisting temptation to keeping long-term goals in mind, executive functions play a critical role in everyday life. However, despite the obvious relevance

of executive functions to overweight and obesity, much is still unknown about executive function abilities in individuals with excess weight. We addressed this gap in the literature by conducting a meta-analysis of effects of obesity and overweight on inhibition, cognitive flexibility, working memory, decision-making, verbal fluency, and planning. Our meta-analysis indicated that obese individuals showed poorer executive functioning across all domains than healthy weight individuals. Unexpectedly, however, overweight participants only showed significant deficits in inhibition and working memory; no differences between overweight and healthy weight individuals emerged on tasks assessing cognitive flexibility, decision-making, verbal fluency, and planning. However, there were too few studies comparing overweight and healthy weight individuals on tasks assessing cognitive flexibility, decision-making, verbal fluency, and planning to make strong claims about the lack of significant differences between groups. Also unexpectedly, age, gender, BMI did not moderate effects of obesity on executive functions. Interestingly, though, the measures used to assess working memory and decision-making moderated effects of obesity on those executive functions. There were not enough studies of overweight individuals to examine moderators of

those effects.

The observed magnitude of effect sizes testing differences in executive functions between obese/overweight individuals and normal weight controls resembles magnitudes found in previous reviews (Lavagnino et al., 2016a; Wu et al., 2016).

However, our meta-analysis extends the findings from previous reviews in several ways.

First, we included a larger total sample of obese/overweight individuals. Second, extending previous meta-analyses that focused on one specific executive function (Wu et al., 2016) or one selected executive function task (Lavagnino et al., 2016a), the present meta-analysis took a wider range of executive function measures into account to cover a broad spectrum of executive function in obese/overweight individuals. In addition, our review was the first to systematically summarize findings of altered inhibition, working memory, verbal fluency and planning in obese/overweight by means of meta-analysis. Finally, the current meta-analysis on executive functions differentiated between obese and overweight individuals. Of note, we found sufficient evidence for a broad impairment of executive functions in obese individuals, whereas there were not enough studies to make definitive inferences about the breadth of

executive function impairment in overweight.

Taken together, the findings of the present meta-analysis support the notion that obesity is associated with poor executive function. Past researchers have assumed or argued that clinical consequences of adiposity, such as hypertension, have an impact on cognition, which may explain this association. However, this hypothesis was not strongly supported by our results, as this meta-analysis included large young samples who were not likely to have cardiovascular disease. As such, it is possible that obesity itself may impair executive functioning. Specifically, some evidence indicates that obesity-related inflammatory activity may impair executive function in obese individuals (e.g., Bourassa, & Sbarra, 2017; Lasselin et al., 2016). As might be inferred from the above, obesity is associated with sustained inflammatory activity, which originates from several factors (Capuron, Lasselin, & Castanon, 2017; Guillemot-Legris, & Muccioli, 2017; Tilg, & Moschen, 2006). In addition, data from several sources, including correlational (e.g., Bourassa, & Sbarra, 2017), longitudinal (e.g., Trompet et al., 2008), experimental (e.g., Grigoleit et al., 2011), and genetic studies (e.g., Mooijaar et al., 2013), converge to suggest that heightened inflammatory activity

contributes to poorer executive function (D O'Brien, et al., 2017; Shields, Moons, & Slavich, 2017). In sum, obesity-related inflammation may be one biological pathway involved in the link between obesity and poor executive function.

It is also possible that poor executive function is a risk factor for an increased BMI, indicating a bidirectional relationship between excess weight and executive function (Martin, & Davidson, 2014; Smith et al., 2011; Stoeckel et al., 2017). The dual-process model argues that our behavior is determined by an interaction between the impulsive system and the executive control system (Strack & Deutsch, 2004). In the case of obesity, coupled with strong impulsive desires regarding high-calorie food and food cues, individuals who show a lack of control over these desires are particularly susceptible to obesity-related behaviors and outcomes (Ziauddeen, Alonso-Alonso, Hill, Kelley, & Khan, 2015). Indeed, both cross-sectional and longitudinal studies showed that lower executive function is associated with greater intake of fatty foods (e.g., Hall, 2012; Allom, & Mullan, 2014; Powell, McMinn, & Allan, 2017), less intake of fruits and vegetables (e.g., Zhou, Gan, Miao, Hamilton, Knoll, & Schwarzer, 2015; Wyckoff, Evans, Manasse, Butryn, & Forman, 2017), poorer adherence to dietary intentions (e.g.,

Hall, Fong, Epp, & Elias, 2008; Spitznagel, Galioto, Limbach, Gunstad, & Heinberg, 2013), less physical activity (e.g., Riggs, Chou, Spruijt-Metz, & Pentz, 2010), higher BMI (e.g., Emery, & Levine, 2017), weight gain (e.g., Nederkoorn, Houben, Hofmann, Roefs, & Jansen, 2010; Nelson, James, Hankey, Nelson, Lundahl, & Espy, 2016), and poor treatment outcomes during weight loss intervention (e.g., Witbracht, Laugero, Van Loan, Adams, & Keim, 2012). In sum, although obesity may itself impair executive function, much evidence indicates that poor executive function also contributes to excessive weight.

The weight-related impairments we observed in executive function are consistent with prior neuroimaging research, which has found altered connectivity among brain regions, structural abnormalities, and task related prefrontal cortical dysfunction (e.g., fronto-striato-parietal circuits) in obese individuals (e.g., García-García et al., 2013; Gearhardt, Yokum, Stice, Harris, & Brownell, 2014). From a functional point of view, several studies have observed obesity-related hypoactivation in prefrontal cortical areas (e.g., inferior frontal gyrus) during performance of executive function tasks (e.g., go/no-go) (Batterink, Yokum, & Stice, 2010; Filbey, & Yezhuvath,

2017; Hendrick, Luo, Zhang, & Li, 2012). Similarly, obese individuals showed less activation of prefrontal regions (e.g., dlPFC, ventral lateral prefrontal cortex) than healthy weight individuals when trying to inhibit responses to high-calorie food stimuli (Batterink, Yokum, & Stice, 2010; Gearhardt et al., 2014; Silvers et al., 2014). In addition, excessive weight is related to lower functional connectivity in frontal-striatal networks during processing of food-related stimuli (García-García et al., 2013; Verdejo-Román, Fornito, Soriano-Mas, Vilar-López, & Verdejo-García, 2017). These results are also consistent with studies focusing on the structural differences between normal and obese individuals. That is, prior studies have found that obesity is related to lower volumes and cortical thickness in several subdivisions of the frontal cortex, including the superior frontal gyrus, middle frontal gyrus, inferior frontal gyrus, and orbitofrontal cortex (Brooks et al., 2013; Kurth et al., 2013; Marqués-Iturria et al., 2013; Pannacciulli et al., 2006; Taki et al., 2008; Walther, Birdsill, Glisky, Ryan, 2010; Wang, Wen, Cheng, & Li, 2017). Importantly, reduced gray matter volume in the bilateral superior frontal gyrus was related to increased BMI at a 1-year follow-up (Yokum, Ng, & Stice, 2012) and the negative relationship between cortical thickness in the right

superior frontal gyrus and BMI was mediated by Stroop task performance (Lavagnino, et al., 2016b). It is important to note that we could not test whether all of the weight-related impairments in executive function we observed were due to these known weight-related alterations in neural structure and function, but examining the neural correlates of the effects we observed is an exciting opportunity for future research.

4.1. Discussion of Moderators

To further illuminate contextual factors that may influence the associations between excess weight and executive function, this meta-analysis examined the effects of potential moderators of these associations, including age, sex, BMI, and measures of executive function. Unexpectedly, however, age, gender, BMI did not moderate any effects of obesity we were able to test. This suggests that the associations between obesity and executive functions may be independent of these demographic factors. The lack of age effect on effects of weight on decision-making is different from what may have been expected based upon the result of the previous meta-analysis, which found non-significant overall decision-making effect size in adolescence (Wu et al., 2016), given that we found an effect on decision-making. However, this difference may be due

to differences in analytic strategy, as Wu and colleagues did not separate obese and overweight samples in their analyses.

We found that the measures used to assess respective executive functions moderated effects of obesity on working memory and decision-making. Studies using the letter-number sequencing task did not show significant effects of obesity on working memory, whereas studies using other working memory tasks, such as the digit span, showed significant deficits in obesity. Similarly, compared to the difference between obese and healthy weight participants in other decision-making tasks, larger effect sizes were observed in decision-making when either the delay discounting task or the Iowa gambling task were used. A possible explanation of this moderating effect might be to do with the sensitivity of the tasks used (Fitzpatrick, Gilbert, & Serpell, 2013). For example, the letter-number sequencing task may be insensitive to detecting subtle differences in working memory performance. However, these results must be interpreted with caution, as the mechanism(s) behind these task-specific effects in working memory and decision-making are currently unknown.

As mentioned earlier, there is some inconsistency in the literature describing

associations between excessive weight and executive function. However, almost none of the potential moderating effects we examined in this meta-analysis emerged as significant. Inconsistencies across studies, such as heterogeneity in executive function tasks, may have obscured the emergence of moderating effects. For example, executive function tasks are likely to vary in sensitivity (Fitzpatrick, Gilbert, & Serpell, 2013). That is, some neuropsychological tasks are designed for use with brain-injured populations and thus are less sensitive to relatively subtle impairments in obese individuals than those designed to assess individual differences in the normal range. In addition, even in analyses of single tasks, task versions and dependent measures may vary, and in turn cause different sensitivities. Thus, we hesitate to draw conclusions about nonsignificant moderators given these and other between-study inconsistencies.

4.2. Limitations and Future Directions

Despite its strengths, this meta-analysis has limitations. First, the small number of studies examining the association between obesity and verbal fluency or planning limited our ability to make inferences about effects of obesity on these executive functions, especially with regard to moderators. As such, when more studies examine

verbal fluency or planning ability in obese individuals, an additional meta-analysis of these associations will be warranted. In addition, the effects on cognitive flexibility, decision making, verbal fluency and planning in overweight also need be interpreted with due caution due to the limited number of original studies. Second, there may be moderators of the associations between obesity and executive functions that are unaccounted for in our analyses. Some studies failed to report important information, such as clinical status of participants, education, economic status, whether participants took any obesity-treating medication, or if the participants had a psychiatric disorder, precluding moderator analyses on these variables. As a result, we could not explain the high heterogeneity in subgroup analysis of studies using Wisconsin Card Sorting Test and moderate heterogeneity in subgroup analysis of studies using Stop Signal Task. In addition, limited number of studies investigated obese individuals with a BMI over 40, which may underestimate the moderating effect of BMI. As such, we do not claim to present a complete picture of moderators of the associations between obesity and executive functions. Indeed, future large-scale studies assessing executive functions in obese/overweight individuals should therefore provide a full characterization of the

participants and explore whether additional factors moderate effects of excessive weight on executive function. Third, although four studies in the obese sample also included some overweight individuals, we classified these samples as obese due to the group's average BMI. However, it is possible that including these studies in analyses may have underestimated true effect sizes. Although analyses suggest against this possibility, we note here that our results should be considered to be a conservative estimate of the association between obesity and executive function. Additionally, we recommend future studies clearly select participants and avoid including overweight and obese individuals in the same group. Fourth, obesity was operationalized via BMI in this meta-analysis, which is a relatively coarse measure of body density and may overlook relevant physical characteristics, such as body fat and anthropometric features (World Health Organization, 2011). Fifth, we were not able to examine the potential role of food stimuli in moderating effects of the association between obesity and executive function. Although we initially coded this variable in our dataset, only four studies used an executive function task that included food content—across all types of executive function tasks. As such, we were unable to analyze the contribution of this

variable. Thus, future research could examine whether the association between obesity and executive function differs by varying stimuli used in the executive function tasks.

What's more, although we have uncovered associations between obesity/overweight and poor executive functions, the studies included were cross-sectional. There is a relative lack of longitudinal studies to investigate whether altered EF is a risk factor or a consequence of obesity/overweight. Thus, more research is needed to investigate causal links between excessive weight and executive function. Finally, although a thorough search was conducted for unpublished studies, because we did not send unsolicited requests to researchers who had published other studies in the field that asked for them to provide data from unpublished studies, it is possible that some unpublished data were not included missed.

5. Conclusion

In conclusion, we found evidence to support the presence of broad executive function deficits in obese participants compared to healthy weight controls. In addition, overweight participants showed significant deficits on inhibition and working memory and nonsignificant deficits on cognitive flexibility, decision-making, verbal fluency,

and planning performance compared to healthy weight controls. Age, gender, BMI did not moderate effects of obesity on executive function. However, the measures used to assess respective executive functions emerged as significant moderators of effects of obesity on working memory and decision-making. Research aimed at determining the causal mechanisms underlying associations between excessive weight and executive functions, as well as effects of executive function interventions on weight status in obese/overweight individuals, deserve further attention because of their important public health implications.

Author Contributions: Yingkai Yang, Cheng Guo and Yanling Liu developed the concept for this article and wrote the manuscript, Grant S. Shields provided critical revisions to the paper, and all authors read and approved the final version.

Declaration of Conflicting Interests: The authors declare that they had no conflicts of interest with respect to their authorship or the publication of this article.

Funding: None

References

- Alarcón, G., Ray, S., & Nagel, B. J. (2016). Lower working memory performance in overweight and obese adolescents is mediated by white matter microstructure. *Journal of the International Neuropsychological Society*, 22(03), 281-292.
- Allom, V., & Mullan, B. (2014). Individual differences in executive function predict distinct eating behaviours. *Appetite*, 80, 123-130.
- Amlung, M., Petker, T., Jackson, J., Balodis, I., & MacKillop, J. (2016). Steep discounting of delayed monetary and food rewards in obesity: a meta-analysis. *Psychological Medicine*, 46(11), 2423-2434.
- Appelhans, B. M. (2009). Neurobehavioral inhibition of reward-driven feeding: implications for dieting and obesity. *Obesity*, 17(4), 640-647.
- Ariza, M., Garolera, M., Jurado, M. A., Garcia-Garcia, I., Hernan, I., Sanchez-Garre, C., ... & Segura, B. (2012). Dopamine genes (DRD2/ANKK1-TaqA1 and DRD4-7R) and executive function: their interaction with obesity. *Plos One*, 7(7), e41482.
- Banich, M. T. (2009). Executive function: The search for an integrated account. *Current Directions in Psychological Science*, 18(2), 89-94.
- Bartholdy, S., Dalton, B., O'Daly, O. G., Campbell, I. C., & Schmidt, U. (2016). A systematic review of the relationship between eating, weight and inhibitory control using the stop signal task. *Neuroscience & Biobehavioral Reviews*, 64, 35-62.
- Batterink, L., Yokum, S., & Stice, E. (2010). Body mass correlates inversely with

inhibitory control in response to food among adolescent girls: an fMRI study. *Neuroimage*, 52(4), 1696-1703.

Benito-León, J., Mitchell, A. J., Hernández-Gallego, J., & Bermejo-Pareja, F. (2013).

Obesity and impaired cognitive functioning in the elderly: a population-based cross-sectional study (NEDICES). *European Journal of Neurology*, 20(6), 899-e77.

Blanco-Gómez, A., Ferré, N., Luque, V., Cardona, M., Gispert-Llauradó, M., Escribano,

J., ... & Canals-Sans, J. (2015). Being overweight or obese is associated with inhibition control in children from six to ten years of age. *Acta Paediatrica*, 104(6), 619-625.

Bocarsly, M. E., Fasolino, M., Kane, G. A., LaMarca, E. A., Kirschen, G. W.,

Karatsoreos, I. N., ... & Gould, E. (2015). Obesity diminishes synaptic markers, alters microglial morphology, and impairs cognitive function. *Proceedings of the National Academy of Sciences*, 112(51), 15731-15736.

Bongers, P., van de Giessen, E., Roefs, A., Nederkoorn, C., Boonij, J., van den Brink,

W., & Jansen, A. (2015). Being impulsive and obese increases susceptibility to speeded detection of high-calorie foods. *Health Psychology*, 34(6), 677-685.

Borenstein, M., Hedges, L.V., Higgins, J.P., Rothstein, H.R. (2009). *Introduction to*

Meta-analysis. Wiley, Chichester.

Bourassa, K., & Sbarra, D. A. (2017). Body mass and cognitive decline are indirectly

associated via inflammation among aging adults. *Brain, Behavior, and Immunity*, 60, 63-70.

- Brogan, A., Hevey, D., & Pignatti, R. (2010). Anorexia, bulimia, and obesity: shared decision making deficits on the Iowa Gambling Task (IGT). *Journal of the International Neuropsychological Society*, 16(04), 711-715.
- Brogan, A., Hevey, D., O'Callaghan, G., Yoder, R., & O'Shea, D. (2011). Impaired decision making among morbidly obese adults. *Journal of Psychosomatic Research*, 70(2), 189-196.
- Brooks, S. J., Benedict, C., Burgos, J., Kempton, M. J., Kullberg, J., Nordenskjöld, R., ... & Ahlström, H. (2013). Late-life obesity is associated with smaller global and regional gray matter volumes: a voxel-based morphometric study. *International journal of obesity*, 37(2), 230-236.
- Calvo, D., Galioto, R., Gunstad, J., & Spitznagel, M. B. (2014). Uncontrolled eating is associated with reduced executive functioning. *Clinical Obesity*, 4(3), 172-179.
- Capuron L, Lasselin J, Castanon N (2017). Role of adiposity-driven inflammation in depressive morbidity. *Neuropsychopharmacology*, 42, 115–128.
- Castanon, N., Lasselin, J., & Capuron, L. (2014). Neuropsychiatric comorbidity in obesity: role of inflammatory processes. *Frontiers in Endocrinology*, 5, 74.
- Chamberlain, S. R., Derbyshire, K. L., Leppink, E., & Grant, J. E. (2015). Obesity and dissociable forms of impulsivity in young adults. *CNS spectrums*, 20(05), 500-507.
- Cohen, J. I., Yates, K. F., Duong, M., & Convit, A. (2011). Obesity, orbitofrontal structure and function are associated with food choice: a cross-sectional study. *BMJ open*, 1(2), e000175.
- Cole, T. J., Bellizzi, M. C., Flegal, K. M., & Dietz, W. H. (2000). Establishing a standard

definition for child overweight and obesity worldwide: international survey. *British Medical Journal*, 320(7244), 1240.

Collins, A., & Koechlin, E. (2012). Reasoning, learning, and creativity: frontal lobe function and human decision-making. *Plos Biology*, 10(3), e1001293.

Coppin, G., Nolan-Poupart, S., Jones-Gotman, M., & Small, D. M. (2014). Working memory and reward association learning impairments in obesity. *Neuropsychologia*, 65, 146-155.

Cserjési, R., Molnár, D., Luminet, O., & Lénárd, L. (2007). Is there any relationship between obesity and mental flexibility in children?. *Appetite*, 49(3), 675-678.

Danner, U. N., Ouwehand, C., Haastert, N. L., Hornsveld, H., & Ridder, D. T. (2012). Decision-making impairments in women with binge eating disorder in comparison with obese and normal weight women. *European Eating Disorders Review*, 20(1), e56-e62.

Delgado-Rico, E., Río-Valle, J. S., González-Jiménez, E., Campoy, C., & Verdejo-García, A. (2012). BMI predicts emotion-driven impulsivity and cognitive inflexibility in adolescents with excess weight. *Obesity*, 20(8), 1604-1610.

Delgado-Rico, E., Soriano-Mas, C., Verdejo-Román, J., S Río-Valle, J., & Verdejo-García, A. (2013). Decreased insular and increased midbrain activations during decision-making under risk in adolescents with excess weight. *Obesity*, 21(8), 1662-1668.

Deux, N., Schlarb, A. A., Martin, F., Holtmann, M., Hebebrand, J., & Legenbauer, T. (2017). Overweight in adolescent, psychiatric inpatients: A problem of general or

food-specific impulsivity?. *Appetite*, 112, 157-166.

Diamond, A. (2013). Executive functions. *Annual Review of Psychology*, 64, 135-168.

D O'Brien, P., Hinder, L. M., Callaghan, B. C., & Feldman, E. L. (2017). Neurological consequences of obesity. *The Lancet Neurology*, 16(6), 465-477.

Egger, M., Smith, G. D., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected by a simple, graphical test. *British Medical Journal*, 315(7109), 629-634.

Eisenstein, S. A., Gredysa, D. M., Antenor-Dorsey, J. A., Green, L., Arbelaez, A. M., Koller, J. M., ... & Hershey, T. (2015). Insulin, central dopamine D2 receptors, and monetary reward discounting in obesity. *Plos one*, 10(7), e0133621.

Emery, R. L., & Levine, M. D. (2017). Questionnaire and behavioral task measures of impulsivity are differentially associated with body mass index: a comprehensive meta-analysis. *Psychological Bulletin*, in press.

Fagundo, A. B., De la Torre, R., Jiménez-Murcia, S., Agüera, Z., Granero, R., Tárrega, S., ... & Forcano, L. (2012). Executive functions profile in extreme eating/weight conditions: from anorexia nervosa to obesity. *Plos one*, 7(8), e43382.

Fagundo, A. B., Jiménez-Murcia, S., Giner-Bartolomé, C., Agüera, Z., Sauchelli, S., Pardo, M., ... & de la Torre, R. (2016). Modulation of Irisin and Physical Activity on Executive Functions in Obesity and Morbid obesity. *Scientific Reports*, 6.

Fields, S. A., Sabet, M., & Reynolds, B. (2013). Dimensions of impulsive behavior in obese, overweight, and healthy-weight adolescents. *Appetite*, 70, 60-66.

Fields, S., Sabet, M., Peal, A., & Reynolds, B. (2011). Relationship between weight status and delay discounting in a sample of adolescent cigarette

- smokers. *Behavioural Pharmacology*, 22(3), 266-268.
- Filbey, F. M., & Yezhuvath, U. S. (2017). A multimodal study of impulsivity and body weight: Integrating behavioral, cognitive, and neuroimaging approaches. *Obesity*, 25(1), 147-154.
- Fitzpatrick, S., Gilbert, S., & Serpell, L. (2013). Systematic review: are overweight and obese individuals impaired on behavioural tasks of executive functioning?. *Neuropsychology Review*, 23(2), 138-156.
- Galioto Wiedemann, R., Calvo, D., Meister, J., & Spitznagel, M. B. (2014). Self-reported physical activity is associated with cognitive function in lean, but not obese individuals. *Clinical Obesity*, 4(6), 309-315.
- Gameiro, F., Perea, M. V., Ladera, V., Rosa, B., & García, R. (2017). Executive functioning in obese individuals waiting for clinical treatment. *Psicothema*, 29(1), 61-66.
- García-García, I., Jurado, M.A., Garolera, M., Segura, B., Marqués-Iturria, I., Pueyo, R., Vernet-Vernet, M., Sender-Palacios, M.J., Sala-Llonch, R., Ariza, M., Narberhaus, A., Junqué, C., 2013. Functional connectivity in obesity during reward processing. *NeuroImage*, 66, 232–239.
- Gearhardt, A. N., Yokum, S., Stice, E., Harris, J. L., & Brownell, K. D. (2014). Relation of obesity to neural activation in response to food commercials. *Social Cognitive and Affective Neuroscience*, 9(7), 932-938.
- Goldschmidt, A. B., O'Brien, S., Lavender, J. M., Pearson, C. M., Le Grange, D., & Hunter, S. J. (2017). Executive functioning in a racially diverse sample of children

who are overweight and at risk for eating disorders. *Appetite*, *in press*.

Gonzales, M. M., Tarumi, T., Miles, S. C., Tanaka, H., Shah, F., & Haley, A. P. (2010).

Insulin sensitivity as a mediator of the relationship between BMI and working memory-related brain activation. *Obesity*, *18*(11), 2131-2137.

Grigoleit, J. S., Kullmann, J. S., Wolf, O. T., Hammes, F., Wegner, A., Jablonowski,

S., ... & Schedlowski, M. (2011). Dose-dependent effects of endotoxin on neurobehavioral functions in humans. *Plos one*, *6*(12), e28330.

Guerrieri, R., Nederkoorn, C., & Jansen, A. (2008). The interaction between

impulsivity and a varied food environment: its influence on food intake and overweight. *International Journal of Obesity*, *32*(4), 708-714.

Guillemot-Legris, O., & Muccioli, G. G. (2017). Obesity-Induced Neuroinflammation:

Beyond the Hypothalamus. *Trends in Neurosciences*, *in press*.

Gunstad, J., Paul, R. H., Cohen, R. A., Tate, D. F., Spitznagel, M. B., & Gordon, E.

(2007). Elevated body mass index is associated with executive dysfunction in otherwise healthy adults. *Comprehensive Psychiatry*, *48*(1), 57-61.

Gunstad, J., Spitznagel, M. B., Paul, R. H., Cohen, R. A., Kohn, M., Luyster, F. S., ...

& Gordon, E. (2008). Body mass index and neuropsychological function in healthy children and adolescents. *Appetite*, *50*(2), 246-251.

Hall, P. A. (2012). Executive control resources and frequency of fatty food consumption:

findings from an age-stratified community sample. *Health Psychology*, *31*(2), 235.

Hall, P.A. (2016). Executive-control processes in high-calorie food

consumption. *Current Directions in Psychological Science*, *25*(2), 91-98.

- Hall, P. A., Fong, G. T., Epp, L. J., & Elias, L. J. (2008). Executive function moderates the intention-behavior link for physical activity and dietary behavior. *Psychology and Health, 23*(3), 309-326.
- Hall, P. A., & Vincent, C. (2017). Non-invasive brain stimulation for food cravings, consumption, and disorders of eating: A review of methods, findings and controversies. *Appetite*, in press.
- Hedges, L. V., Tipton, E., & Johnson, M. C. (2010). Robust variance estimation in meta-regression with dependent effect size estimates. *Research Synthesis Methods, 1*(1), 39-65.
- Hendrick, O. M., Luo, X., Zhang, S., & Li, C. S. R. (2012). Saliency processing and obesity: a preliminary imaging study of the stop signal task. *Obesity, 20*(9), 1796-1802.
- Hofmann, W., Friese, M., & Strack, F. (2009). Impulse and self-control from a dual-systems perspective. *Perspectives on Psychological Science, 4*(2), 162-176.
- Hofmann, W., Schmeichel, B. J., & Baddeley, A. D. (2012). Executive functions and self-regulation. *Trends in Cognitive Sciences, 16*(3), 174-180.
- Hong, C. (2013). *Deficiencies in inhibitory mechanism in obesity* (Master dissertation, University of Delaware, USA).
- Hsu, C. L., Voss, M. W., Best, J. R., Handy, T. C., Madden, K., Bolandzadeh, N., & Liu-Ambrose, T. (2015). Elevated body mass index and maintenance of cognitive function in late life: exploring underlying neural mechanisms. *Frontiers in Aging Neuroscience, 7*, 155.

- Jansen, A., Houben, K., & Roefs, A. (2015). A cognitive profile of obesity and its translation into new interventions. *Frontiers in Psychology*, 6, 1807.
- Kamijo, K., Khan, N. A., Pontifex, M. B., Scudder, M. R., Drollette, E. S., Raine, L. B., ... & Hillman, C. H. (2012a). The relation of adiposity to cognitive control and scholastic achievement in preadolescent children. *Obesity*, 20(12), 2406-2411.
- Kamijo, K., Pontifex, M. B., Khan, N. A., Raine, L. B., Scudder, M. R., Drollette, E. S., ... & Hillman, C. H. (2014). The negative association of childhood obesity to cognitive control of action monitoring. *Cerebral Cortex*, 24(3), 654-662.
- Kamijo, K., Pontifex, M. B., Khan, N. A., Raine, L. B., Scudder, M. R., Drollette, E. S., ... & Hillman, C. H. (2012b). The association of childhood obesity to neuroelectric indices of inhibition. *Psychophysiology*, 49(10), 1361-1371.
- Kittel, R., Schmidt, R., & Hilbert, A. (2017). Executive functions in adolescents with binge-eating disorder and obesity. *International Journal of Eating Disorders*, in press.
- Kulendran, M., Patel, K., Darzi, A., & Vlaev, I. (2016). Diagnostic validity of behavioural and psychometric impulsivity measures: An assessment in adolescent and adult populations. *Personality and Individual Differences*, 90, 347-352.
- Kulendran, M., Vlaev, I., Sugden, C., King, D., Ashrafian, H., Gately, P., & Darzi, A. (2014). Neuropsychological assessment as a predictor of weight loss in obese adolescents. *International Journal of Obesity*, 38(4), 507-512.
- Kurth, F., Levitt, J.G., Phillips, O.R., Luders, E., Woods, R.P., Mazziotta, J.C., Toga, A. W., Narr, K.L., 2013. Relationships between gray matter, body mass index, and

- waist circumference in healthy adults. *Human Brain Mapping*, 34 (7), 1737–1746.
- Kyrgiou, M., Kalliala, I., Markozannes, G., Gunter, M. J., Paraskevaïdis, E., Gabra, H., ... & Tsilidis, K. K. (2017). Adiposity and cancer at major anatomical sites: umbrella review of the literature. *British Medical Journal*, 356, j477.
- Lasselin, J., Magne, E., Beau, C., Aubert, A., Dexpert, S., Carrez, J., ... & Capuron, L. (2016). Low-grade inflammation is a major contributor of impaired attentional set shifting in obese subjects. *Brain, Behavior, and Immunity*, 58, 63-68.
- Lavagnino, L., Arnone, D., Cao, B., Soares, J. C., & Selvaraj, S. (2016a). Inhibitory control in obesity and binge eating disorder: A systematic review and meta-analysis of neurocognitive and neuroimaging studies. *Neuroscience & Biobehavioral Reviews*, 68, 714-726.
- Lavagnino, L., Mwangi, B., Bauer, I. E., Cao, B., Selvaraj, S., Prossin, A., & Soares, J. C. (2016b). Reduced inhibitory control mediates the relationship between cortical thickness in the right superior frontal gyrus and body mass index. *Neuropsychopharmacology*, 41 (9), 2275–2282.
- Lavie, C. J., Milani, R. V., & Ventura, H. O. (2009). Obesity and cardiovascular disease. *Journal of the American College of Cardiology*, 53(21), 1925-1932.
- Lawyer, S. R., Boomhower, S. R., & Rasmussen, E. B. (2015). Differential associations between obesity and behavioral measures of impulsivity. *Appetite*, 95, 375-382.
- Lezak, M. D. (2012). *Neuropsychological assessment*. Oxford University Press, USA.
- Ljubisavljevic, M., Maxood, K., Bjekic, J., Oommen, J., & Nagelkerke, N. (2016). Long-term effects of repeated prefrontal cortex transcranial direct current

- stimulation (tDCS) on food craving in normal and overweight young adults. *Brain stimulation*, 9(6), 826-833.
- Liang, J., Matheson, B. E., Kaye, W. H., & Boutelle, K. N. (2014). Neurocognitive correlates of obesity and obesity-related behaviors in children and adolescents. *International Journal of Obesity*, 38(4), 494-506.
- Loeber, S., Grosshans, M., Korucuoglu, O., Vollmert, C., Vollstädt-Klein, S., Schneider, S., ... & Kiefer, F. (2012). Impairment of inhibitory control in response to food-associated cues and attentional bias of obese participants and normal-weight controls. *International Journal of Obesity*, 36(10), 1334-1339.
- Luppino, F. S., de Wit, L. M., Bouvy, P. F., Stijnen, T., Cuijpers, P., Penninx, B. W., & Zitman, F. G. (2010). Overweight, obesity, and depression: a systematic review and meta-analysis of longitudinal studies. *Archives of General Psychiatry*, 67(3), 220-229.
- Maayan, L., Hoogendoorn, C., Sweat, V., & Convit, A. (2011). Disinhibited eating in obese adolescents is associated with orbitofrontal volume reductions and executive dysfunction. *Obesity*, 19(7), 1382-1387.
- Mallorquí-Bagué, N., Fagundo, A. B., Jimenez-Murcia, S., de la Torre, R., Baños, R. M., Botella, C., ... & Frühbeck, G. (2016). Decision Making Impairment: A Shared Vulnerability in Obesity, Gambling Disorder and Substance Use Disorders?. *Plos One*, 11(9), e0163901.
- Marqués-Iturria, I., Pueyo, R., Garolera, M., Segura, B., Junqué, C., García-García, I., ... & Jurado, M. Á. (2013). Frontal cortical thinning and subcortical volume

- reductions in early adulthood obesity. *Psychiatry Research: Neuroimaging*, 214(2), 109-115.
- Martin, A. A., & Davidson, T. L. (2014). Human cognitive function and the obesogenic environment. *Physiology & Behavior*, 136, 185-193.
- Mata, F., Verdejo-Roman, J., Soriano-Mas, C., & Verdejo-Garcia, A. (2015). Insula tuning towards external eating versus interoceptive input in adolescents with overweight and obesity. *Appetite*, 93, 24-30.
- McClelland, J., Dalton, B., Kekic, M., Bartholdy, S., Campbell, I. C., & Schmidt, U. (2016). A systematic review of temporal discounting in eating disorders and obesity: Behavioural and neuroimaging findings. *Neuroscience & Biobehavioral Reviews*, 71, 506-528.
- Miller, A. A., & Spencer, S. J. (2014). Obesity and neuroinflammation: a pathway to cognitive impairment. *Brain, Behavior, and Immunity*, 42, 10-21.
- Miyake, A., & Friedman, N. P. (2012). The nature and organization of individual differences in executive functions four general conclusions. *Current Directions in Psychological Science*, 21(1), 8-14.
- Miyake, A., Friedman, N. P., Emerson, M. J., Witzki, A. H., Howerter, A., & Wager, T. D. (2000). The unity and diversity of executive functions and their contributions to complex “frontal lobe” tasks: A latent variable analysis. *Cognitive Psychology*, 41(1), 49-100.
- Mokdad, A. H., Ford, E. S., Bowman, B. A., Dietz, W. H., Vinicor, F., Bales, V. S., & Marks, J. S. (2003). Prevalence of obesity, diabetes, and obesity-related health risk

- factors, 2001. *Jama*, 289(1), 76-79.
- Mole, T. B., Irvine, M. A., Worbe, Y., Collins, P., Mitchell, S. P., Bolton, S., ... & Voon, V. (2015). Impulsivity in disorders of food and drug misuse. *Psychological Medicine*, 45(04), 771-782.
- Moreno-López, L., Soriano-Mas, C., Delgado-Rico, E., Rio-Valle, J. S., & Verdejo-García, A. (2012). Brain structural correlates of reward sensitivity and impulsivity in adolescents with normal and excess weight. *Plos one*, 7(11), e49185.
- Navas, J. F., Vilar-López, R., Perales, J. C., Steward, T., Fernández-Aranda, F., & Verdejo-García, A. (2016). Altered decision-making under risk in obesity. *Plos one*, 11(6), e0155600.
- Nederkoorn, C. (2014). Effects of sales promotions, weight status, and impulsivity on purchases in a supermarket. *Obesity*, 22(5), E2-E5.
- Nederkoorn, C., Braet, C., Van Eijs, Y., Tanghe, A., & Jansen, A. (2006a). Why obese children cannot resist food: the role of impulsivity. *Eating behaviors*, 7(4), 315-322.
- Nederkoorn, C., Coelho, J. S., Guerrieri, R., Houben, K., & Jansen, A. (2012). Specificity of the failure to inhibit responses in overweight children. *Appetite*, 59(2), 409-413.
- Nederkoorn, C., Houben, K., Hofmann, W., Roefs, A., & Jansen, A. (2010). Control yourself or just eat what you like? Weight gain over a year is predicted by an interactive effect of response inhibition and implicit preference for snack foods. *Health Psychology*, 29(4), 389-393.

Nederkoorn, C., Smulders, F. T., Havermans, R. C., Roefs, A., & Jansen, A. (2006b).

Impulsivity in obese women. *Appetite*, 47(2), 253-256.

Nee, D. E., Wager, T. D., & Jonides, J. (2007). Interference resolution: insights from a meta-analysis of neuroimaging tasks. *Cognitive, Affective, & Behavioral Neuroscience*, 7(1), 1-17.

Nelson, T. D., James, T. D., Hankey, M., Nelson, J. M., Lundahl, A., & Espy, K. A. (2016). Early executive control and risk for overweight and obesity in elementary school. *Child Neuropsychology*, 1-9.

Ng, M., Fleming, T., Robinson, M., Thomson, B., Graetz, N., Margono, C., ... & Abraham, J. P. (2014). Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. *The Lancet*, 384(9945), 766-781.

Pannacciulli, N., Del Parigi, A., Chen, K., Le, D.S., Reiman, E.M., Tataranni, P.A., 2006. Brain abnormalities in human obesity: a voxel-based morphometric study. *NeuroImage*, 31 (4), 1419–1425.

Perpiñá, C., Segura, M., & Sánchez-Reales, S. (2016). Cognitive flexibility and decision-making in eating disorders and obesity. *Eating and Weight Disorders-Studies on Anorexia, Bulimia and Obesity*, 1-10.

Pignatti, R., Bertella, L., Albani, G., Mauro, A., Molinari, E., & Semenza, C. (2006). Decision-making in obesity: a study using the Gambling Task. *Eating and Weight Disorders-Studies on Anorexia, Bulimia and Obesity*, 11(3), 126-132.

Powell, D. J., McMinn, D., & Allan, J. L. (2017). Does Real Time Variability in

Inhibitory Control Drive Snacking Behavior? An Intensive Longitudinal Study.

Health Psychology, in press.

Prickett, C., Brennan, L., & Stolwyk, R. (2015). Examining the relationship between obesity and cognitive function: a systematic literature review. *Obesity Research & Clinical Practice, 9*(2), 93-113.

Qavam, S. E., Anisan, A., Fathi, M., & Pourabbasi, A. (2015). Study of relationship between obesity and executive functions among high school students in Bushehr, Iran. *Journal of Diabetes & Metabolic Disorders, 14*(1), 79.

Rangel, A., Camerer, C., & Montague, P. R. (2008). A framework for studying the neurobiology of value-based decision making. *Nature Reviews Neuroscience, 9*(7), 545-556.

Restivo, M. R., McKinnon, M. C., Frey, B. N., Hall, G. B., Syed, W., & Taylor, V. H. (2017). The impact of obesity on neuropsychological functioning in adults with and without major depressive disorder. *Plos One, 12*(5).

Reyes, S., Peirano, P., Peigneux, P., Lozoff, B., & Algarin, C. (2015). Inhibitory control in otherwise healthy overweight 10-year-old children. *International Journal of Obesity, 39*(8), 1230-1235.

Riggs, N., Chou, C. P., Spruijt-Metz, D., & Pentz, M. A. (2010). Executive cognitive function as a correlate and predictor of child food intake and physical activity. *Child Neuropsychology, 16*(3), 279-292.

Ross, N., Yau, P. L., & Convit, A. (2015). Obesity, fitness, and brain integrity in adolescence. *Appetite, 93*, 44-50.

Rotge, J. Y., Poitou, C., Fossati, P., Aron-Wisnewsky, J., & Oppert, J. M. (2017).

Decision-making in obesity without eating disorders: a systematic review and meta-analysis of Iowa gambling task performances. *Obesity Reviews*, *in press*.

Scammacca, N., Roberts, G., & Stuebing, K. K. (2014). Meta-analysis with complex research designs dealing with dependence from multiple measures and multiple group comparisons. *Review of Educational Research*, *84*(3), 328-364.

Schiff, S., Amodio, P., Testa, G., Nardi, M., Montagnese, S., Caregaro, L., ... & Sellitto, M. (2016). Impulsivity toward food reward is related to BMI: evidence from intertemporal choice in obese and normal-weight individuals. *Brain and Cognition*, *110*, 112-119.

Schiff, S., Amodio, P., Testa, G., Nardi, M., Montagnese, S., Caregaro, L., ... & Sellitto, M. (2016). Impulsivity toward food reward is related to BMI: evidence from intertemporal choice in obese and normal-weight individuals. *Brain and Cognition*, *110*, 112-119.

Sellaro, R., & Colzato, L. S. (2017). High body mass index is associated with impaired cognitive control. *Appetite*, *113*, 301-309.

Shields, G. S., Moons, W. G., & Slavich, G. M. (2017). Inflammation, Self-Regulation, and Health: An Immunologic Model of Self-Regulatory Failure. *Perspectives on Psychological Science*, *12*(4), 588-612.

Shields, G. S., Sazma, M. A., & Yonelinas, A. P. (2016). The effects of acute stress on core executive functions: A meta-analysis and comparison with cortisol. *Neuroscience & Biobehavioral Reviews*, *68*, 651-668.

Silveira, L. E., Kozicky, J. M., Muralidharan, K., Bücker, J., Torres, I. J., Bond, D. J., ...

& Yatham, L. N. (2014). Neurocognitive functioning in overweight and obese patients with bipolar disorder: data from the systematic treatment optimization program for early mania (STOP-EM). *The Canadian Journal of Psychiatry*, 59(12), 639-648.

Silvers, J. A., Insel, C., Powers, A., Franz, P., Weber, J., Mischel, W., ... & Ochsner, K.

N. (2014). Curbing craving behavioral and brain evidence that children regulate craving when instructed to do so but have higher baseline craving than adults. *Psychological Science*, 25, 1932–1942.

Simmank, J., Murawski, C., Bode, S., & Horstmann, A. (2015). Incidental rewarding

cues influence economic decisions in people with obesity. *Frontiers in Behavioral Neuroscience*, 9.

Skoranski, A. M., Most, S. B., Lutz-Stehl, M., Hoffman, J. E., Hassink, S. G., & Simons,

R. F. (2013). Response monitoring and cognitive control in childhood obesity. *Biological Psychology*, 92(2), 199-204.

Smith, E., Hay, P., Campbell, L., & Trollor, J. N. (2011). A review of the association

between obesity and cognitive function across the lifespan: implications for novel approaches to prevention and treatment. *Obesity Reviews*, 12(9), 740-755.

Snyder, H. R. (2013). Major depressive disorder is associated with broad impairments

on neuropsychological measures of executive function: A meta-analysis and review. *Psychological Bulletin*, 139 (1) ,81-132.

Spitoni, G. F., Ottaviani, C., Petta, A. M., Zingaretti, P., Aragona, M., Sarnicola, A., &

- Antonucci, G. (2017). Obesity is associated with lack of inhibitory control and impaired heart rate variability reactivity and recovery in response to food stimuli. *International Journal of Psychophysiology*, in press.
- Spitznagel, M. B., Galioto, R., Limbach, K., Gunstad, P. D. J., & Heinberg, P. D. L. (2013). Cognitive function is linked to adherence to bariatric postoperative guidelines. *Surgery for Obesity and Related Diseases*, 9(4), 580-585.
- Spyridaki, E. C., Avgoustinaki, P. D., & Margioris, A. N. (2016). Obesity, inflammation and cognition. *Current Opinion in Behavioral Sciences*, 9, 169-175.
- Stingl, K. T., Kullmann, S., Ketterer, C., Heni, M., Häring, H. U., Fritsche, A., & Preissl, H. (2012). Neuronal correlates of reduced memory performance in overweight subjects. *NeuroImage*, 60(1), 362-369.
- Stoeckel, L. E., Birch, L. L., Heatherton, T., Mann, T., Hunter, C., Czajkowski, S., ... & Savage, C. R. (2017). Psychological and neural contributions to appetite self-regulation. *Obesity*, 25(S1), 17-25.
- Stojek, M. M., & MacKillop, J. (2017). Relative reinforcing value of food and delayed reward discounting in obesity and disordered eating: A systematic review. *Clinical Psychology Review*, in press.
- Strack, F., & Deutsch, R. (2004). Reflective and impulsive determinants of social behavior. *Personality and Social Psychology Review*, 8(3), 220-247.
- Suchy, Y. (2009). Executive functioning: Overview, assessment, and research issues for non-neuropsychologists. *Annals of behavioral medicine*, 37(2), 106-116.
- Sweat, V., Yates, K. F., Migliaccio, R., & Convit, A. (2017). Obese Adolescents Show

Reduced Cognitive Processing Speed Compared with Healthy Weight Peers. *Childhood Obesity*, in press.

Taki, Y., Kinomura, S., Sato, K., Inoue, K., Goto, R., Okada, K., Uchida, S., Kawashima, R., Fukuda, H., 2008. Relationship between body mass index and gray matter volume in 1428 healthy individuals. *Obesity*, 16 (1), 119–124.

Tanner-Smith, E. E., & Tipton, E. (2014). Robust variance estimation with dependent effect sizes: practical considerations including a software tutorial in Stata and SPSS. *Research Synthesis Methods*, 5(1), 13-30.

Testa, R., Bennett, P., & Ponsford, J. (2012). Factor analysis of nineteen executive function tests in a healthy adult population. *Archives of Clinical Neuropsychology*, 27, 213-224.

Tilg, H., & Moschen, A. R. (2006). Adipocytokines: Mediators linking adipose tissue, inflammation and immunity. *Nature Reviews Immunology*, 6, 772–783.

Tipton, E. (2015). Small sample adjustments for robust variance estimation with meta-regression. *Psychological Methods*, 20(3), 375-393.

Trompet, S., De Craen, A. J. M., Slagboom, P., Shepherd, J., Blauw, G. J., Murphy, M. B., ... & Macfarlane, P. W. (2008). Genetic variation in the interleukin-1 β -converting enzyme associates with cognitive function. The PROSPER study. *Brain*, 131(4), 1069-1077.

Troyer, A. K., Moscovitch, M., & Winocur, G. (1997). Clustering and switching as two components of verbal fluency: evidence from younger and older healthy adults. *Neuropsychology*, 11(1), 138-146.

- Tsai, C. L., Chen, F. C., Pan, C. Y., & Tseng, Y. T. (2016). The neurocognitive performance of visuospatial attention in children with obesity. *Frontiers in Psychology*, 7.
- Vainik, U., Dagher, A., Dubé, L., & Fellows, L. K. (2013). Neurobehavioural correlates of body mass index and eating behaviours in adults: a systematic review. *Neuroscience & Biobehavioral Reviews*, 37(3), 279-299.
- Verdejo-García, A., Pérez-Expósito, M., Schmidt-Río-Valle, J., Fernández-Serrano, M. J., Cruz, F., Pérez-García, M., ... & Campoy, C. (2010). Selective alterations within executive functions in adolescents with excess weight. *Obesity*, 18(8), 1572-1578.
- Verdejo-Román, J., Fornito, A., Soriano-Mas, C., Vilar-López, R., & Verdejo-García, A. (2017). Independent functional connectivity networks underpin food and monetary reward sensitivity in excess weight. *NeuroImage*, 146, 293-300.
- Viechtbauer, W., & Cheung, M. W. L. (2010). Outlier and influence diagnostics for meta-analysis. *Research Synthesis Methods*, 1(2), 112-125.
- Voon, V., Irvine, M. A., Derbyshire, K., Worbe, Y., Lange, I., Abbott, S., ... & Wood, J. (2014). Measuring “waiting” impulsivity in substance addictions and binge eating disorder in a novel analogue of rodent serial reaction time task. *Biological Psychiatry*, 75(2), 148-155.
- Walther, K., Birdsill, A.C., Glisky, E.L., Ryan, L., 2010. Structural brain differences and cognitive functioning related to body mass index in older females. *Human Brain Mapping*, 31, 1052–1064.
- Wang, H., Wen, B., Cheng, J., & Li, H. (2017). Brain Structural Differences between

- Normal and Obese Adults and their Links with Lack of Perseverance, Negative Urgency, and Sensation Seeking. *Scientific Reports*, 7, 40595.
- Weller, R. E., Cook, E. W., Avsar, K. B., & Cox, J. E. (2008). Obese women show greater delay discounting than healthy-weight women. *Appetite*, 51(3), 563-569.
- Weller, R. E., Cook, E. W., Avsar, K. B., & Cox, J. E. (2008). Obese women show greater delay discounting than healthy-weight women. *Appetite*, 51(3), 563-569.
- Witbracht, M. G., Laugero, K. D., Van Loan, M. D., Adams, S. H., & Keim, N. L. (2012). Performance on the Iowa Gambling Task is related to magnitude of weight loss and salivary cortisol in a diet-induced weight loss intervention in overweight women. *Physiology & Behavior*, 106(2), 291-297.
- Withrow, D., & Alter, D. A. (2011). The economic burden of obesity worldwide: a systematic review of the direct costs of obesity. *Obesity Reviews*, 12(2), 131-141.
- World Health Organization (2011). *Waist Circumference and Waist-Hip Ratio: Report of a WHO Expert Consultation*. Geneva: World Health Organization.
- World Health Organization. (2017). *Obesity and Overweight*. Geneva: World Health Organization.
- Wu, M., Brockmeyer, T., Hartmann, M., Skunde, M., Herzog, W., & Friederich, H. C. (2016). Reward-related decision making in eating and weight disorders: A systematic review and meta-analysis of the evidence from neuropsychological studies. *Neuroscience & Biobehavioral Reviews*, 61, 177-196.
- Wu, M., Brockmeyer, T., Hartmann, M., Skunde, M., Herzog, W., & Friederich, H. C. (2014). Set-shifting ability across the spectrum of eating disorders and in

- overweight and obesity: a systematic review and meta-analysis. *Psychological Medicine*, 44(16), 3365-3385.
- Wu, N., Chen, Y., Yang, J., & Li, F. (2017). Childhood Obesity and Academic Performance: The Role of Working Memory. *Frontiers in Psychology*, 8.
- Wu, X., Nussbaum, M. A., & Madigan, M. L. (2016). Executive function and measures of fall risk among people with obesity. *Perceptual and Motor Skills*, 122(3), 825-839.
- Wyckoff, E. P., Evans, B. C., Manasse, S. M., Butryn, M. L., & Forman, E. M. (2017). Executive functioning and dietary intake: Neurocognitive correlates of fruit, vegetable, and saturated fat intake in adults with obesity. *Appetite*, 111, 77-85.
- Yang, Y., Cao, S., Shields, G. S., Teng, Z., & Liu, Y. (2017). The relationships between rumination and core executive functions: A meta-analysis. *Depression and Anxiety*, 34(1), 37-50.
- Yau, P. L., Kang, E. H., Javier, D. C., & Convit, A. (2014). Preliminary evidence of cognitive and brain abnormalities in uncomplicated adolescent obesity. *Obesity*, 22(8), 1865-1871.
- Yeomans, M. R., Leitch, M., & Mobini, S. (2008). Impulsivity is associated with the disinhibition but not restraint factor from the Three Factor Eating Questionnaire. *Appetite*, 50(2), 469-476.
- Yokum, S., Ng, J., & Stice, E. (2012). Relation of regional gray and white matter volumes to current BMI and future increases in BMI: A prospective MRI study. *International Journal of Obesity*, 36, 656-664.

Zhou, G., Gan, Y., Miao, M., Hamilton, K., Knoll, N., & Schwarzer, R. (2015). The role of action control and action planning on fruit and vegetable consumption. *Appetite*, 91, 64-68.

Ziauddeen, H., Alonso-Alonso, M., Hill, J. O., Kelley, M., & Khan, N. A. (2015). Obesity and the neurocognitive basis of food reward and the control of intake. *Advances in Nutrition: An International Review Journal*, 6(4), 474-486.

Figure 1 Flow diagram illustrating the process of our review, screening, and article selections

Figure 2 Funnel plots to ascertain evidence for publication bias in obesity

Figure 3 Funnel plots to ascertain evidence for publication bias of inhibition and working memory in overweight

Figure 4 Inhibition deficit in obesity individuals

Figure 5 Cognitive flexibility deficit in obesity individuals

Figure 6 Working memory deficit in obesity individuals

Figure 7 Decision making deficit in obesity individuals

Figure 8 Verbal fluency deficit in obesity individuals

Figure 9 Planning deficit in obesity individuals

Figure 10 Inhibition deficit in overweight individuals

Figure 11 Working memory deficit in overweight individuals




Figure 1 Flow diagram illustrating the process of our review, screening, and article

selections




Figure 2 Funnel plots to ascertain evidence for publication bias in obesity




Figure 3 Funnel plots to ascertain evidence for publication bias of inhibition and working memory in overweight




Figure 4 Inhibition deficit in obesity individuals




Figure 5 Cognitive flexibility deficit in obesity individuals




Figure 6 Working memory deficit in obesity individuals




Figure 7 Decision making deficit in obesity individuals




Figure 8 Verbal fluency deficit in obesity individuals




Figure 9 Planning deficit in obesity individuals




Figure 10 Inhibition deficit in overweight individuals




Figure 11 Working memory deficit in overweight individuals

Table 1 studies included in the meta-analysis

Study	Subject (sample size)	Female (%)	Mean age (years)	Mean BMI (kg/m ²)	Executive functions studied	Task(s) Used	Quality score (x/6)
Alarcón et al., 2016	OB (18)	33.3%	14.4	30.9	Working memory	Spatial 2-back	5
	OW (46)	45.7%	13.8	24.2		Verbal 2-back	
	NW (88)	45.5%	14.2	20.3			
Ariza et al., 2012	OB (42)	66.7%	31.8	38.3	Inhibition	Stroop	3
	NW (42)	69.1%	29.7	22.1	Cognitive flexibility	Trail Making Test B/B-A	
					Working memory	WCST	
					Verbal fluency	Letter-Number Sequencing Phonemic Verbal Fluency	
Benito-Leon et al., 2013	OB (592)	62.8%	74.9	≥30	Verbal fluency	Semantic Verbal Fluency	1
	OW (850)	51.2%	75.7	25 - 29.9			
	NW (507)	58.4%	76.9	< 25			
Blanco-Gomez et al., 2015	OB (130)	46.2%	8.5	N.R.	Inhibition	Five Digit Test	4
	NW (90)	40%	8.5	N.R.	Cognitive flexibility		
Bongers et al., 2014	OB (185)	70.8%	35.2	38.2	Inhibition	Stop Signal Task.	3
	NW (134)	73.9%	33.0	22.4	Decision making	Delay Discounting Task	

Brogan et al., 2010	OB (18)	100%	52.1	36.2	Decision making	The Iowa Gambling Task	4
	NW (20)	100%	27.8	21.6			
Brogan et al., 2011	OB (42)	71.4%	52.2	41.5	Decision making	The Iowa Gambling Task	3
	NW (50)	66.0%	47.3	24.4			
Brooks et al., 2013	OB (112)	100%	72.5	≥ 30	Cognitive flexibility	Trail Making Test B-A	2
	NW (180)	100%	72.5	< 25			
Calvo et al., 2014	OB (30)	60%	21.2	36.4	Inhibition	Go/No-Go	5
	NW (32)	53.1%	21.1	21.7	Working memory	RMCP	
Chamberlain et al., 2015	OB (55)	60%	24.2	35.6	Inhibition	Stop Signal Task.	1
	OW (110)	29.1%	23	26.9	Cognitive flexibility	IDET	
	NW (346)	34.1%	21.3	21.6	Decision making	Cambridge Gamble Task	
Cohen et al., 2011					Planning	OTSCT	3
	OB (42)	47.7%	58.9	31.8	Inhibition	Stroop	
	NW (107)	52.3%	61.2	24.1	Cognitive flexibility	Trail Making Test B-A	
					Working memory	WCST	
						Digit Span	
Coppin et al., 2014						Visual Working Memory Task	5
	OB (17)	52.9%	25.2	36	Working memory	CCPT	
	OW (16)	43.8%	24.9	27.6			

	NW (14)	64.3%	24.3	27.6			
Cserje' si et al., 2007	OB (12)	0%	12.1	27.2	Cognitive flexibility	WCST	4
	NW (12)	0%	12.4	16.9			
Danner et al., 2012	OB (18)	100%	44.7	30.8	Decision making	The Iowa Gambling Task	2
	NW (30)	100%	36.1	22.3			
Delgado-Rico et al., 2012	OB (42)	66.7%	14.2	29.2	Inhibition	Stroop	5
	NW (21)	47.6%	14.1	19.8	Cognitive flexibility		
Delgado-Rico et al., 2013	OB (14)	64.3%	14.2	31.1	Decision making	Risk Gain Task	4
	OW (13)	69.2%	14.1	24.7			
	NW (13)	61.5%	13.7	20.4			
Deux et al., 2017	OB (20 & 23)	55%& 64.4 %	14.4	N.R.	Inhibition	Go/no-go	4
	NW (59 & 57)	56.5%&64.9 %	15.3	N.R.			
Eisenstein et al., 2015	OB (27)	85.2%	31.5	39.9	Decision making	Delay Discounting Task	4
	NW (20)	75%	28.6	22.4		Probability Discounting Task	
Fagundo et al., 2012	OB (52)	100%	40.5	39.8	Inhibition	Stroop	3
	NW (137)	100%	24.8	21.5	Decision making	The Iowa Gambling Task	
Fagundo et al., 2016	OB (44 & 21)	100%	42.3 & 49.2	46.3 & 35.5	Inhibition	Stroop	3
	NW (49)	100%	29	21.61	Cognitive flexibility	WCST	
					Decision making	The Iowa Gambling Task	

Fields et al., 2011	OB (16)	68.8%	17.2	35.4	Decision making	Delay Discounting Task	3
	NW (20)	20%	17.4	21.3			
Fields et al., 2013	OB (21)	52.4%	14.9	34.4	Decision making	Delay Discounting Task	3
	OW (20)	55%	15.2	26.1			
	NW (20)	60%	15	20.4			
Galioto Wiedemann et al., 2014	OB (36)	61.1%	21.2	36.4	Inhibition	Go/No-Go	4
	NW (36)	50%	20.7	22	Working memory	RMCPT	
Gameiro et al., 2017	OB (76)	68.4%	43.2	> 30	Inhibition	Stroop	4
	NW (38)	71.1%	40.5	18-24.9	Shifting	WCST	
						Color trail test	
Goldschmidt et al., 2017	OB (34 & 26)	55.9%&61.5%	10.8 &10.2	N.R.	Inhibition	Flanker Task	5
	NW (15)	60%	10.4	N.R.	Cognitive flexibility	DCCTT	
					Working memory	List Sorting Task	
					Decision making	The Iowa Gambling Task	
					Planning	Tower of London Task	
Gonzales et al., 2010	OB (12)	50%	48.5	34.3	Cognitive flexibility	Trail Making Test B	4
	OW (11)	45.5%	52	27.4	Working memory	Digit Span	
	NW (9)	77.9%	51.8	22.4	Verbal fluency	Phonemic Verbal Fluency	
Guerrieri et al., 2008	OW (15)	N.R.	9	21.8	Inhibition	Stop Signal Task	2

	NW (63)	N.R.	9	16.3	Decision making	Door Open Task	
Gunstad et al., 2007(1)	OW (140)	46.4%	32.4	28.4	Inhibition	Stroop	5
	NW (178)	55.1%	31.4	22.1	Cognitive flexibility	Trail Making Test B	
					Working memory	Digit Span	
					Planning	Visual Working Memory Task	
						Maze task	
Gunstad et al., 2007(2)	OW (58)	55.1%	60.4	29.2	Inhibition	Stroop	5
	NW (32)	53.1%	58.3	23.1	Cognitive flexibility	Trail Making Test B	
					Working memory	Digit Span	
					Planning	Visual Working Memory Task	
						Maze Task	
Gunstad et al., 2008	OB (45)	38%	10.8	26.2	Cognitive flexibility	Trail Making Test B	2 & 4
	OW (76)	47%	12.6	23.1	Working memory	Digit Span	
	NW (330)	51%	12.6	18.5	Verbal fluency	Semantic Verbal Fluency	
Hendrick et al., 2012	OB (13)	100%	34.8	33.2	Inhibition	Stop Signal Task	2
	OW (8)	100%	33.2	25.6			
	NW (18)	100%	26.2	20.0			
Hong, 2013	OB (27)	N.R.	N.R.	> 30	Inhibition	Stop Signal Task	2
	NW (29)	N.R.	N.R.	18-24.9			

Hsu et al., 2015	OB (15)	86.7%	75	32.7	Inhibition	Stroop	4 & 3
	OW (27)	55.6%	74	27.6	Cognitive flexibility	Trail Making Test B-A	
	NW (24)	87.5%	74	22.3	Working memory	Digit Span	
Kamijo et al., 2012a	OB (30)	56.7%	9	26.1	Inhibition	Go/No-Go	5
	OW (26)	46.2%	8.7	19.5			
	NW (70)	48.6%	8.9	16.5			
Kamijo et al., 2012b	OB (37)	51.4%	7.9	25.3	Inhibition	Go/No-Go	5
	NW (37)	51.4%	9	16.8			
Kamijo et al., 2014	OB (37)	54.1%	8.9	25.2	Inhibition	Flanker task	5
	NW (37)	54.1%	8.8	16.7			
Kittel et al., 2017	OB (22)	81.8%	14.8	N.R.	Inhibition	Stroop	5
	NW (22)	81.8%	15.2	N.R.	Cognitive flexibility	Comprehensive Trail Making	
					Decision making	Test	
						Iowa Gambling Task	
Kulendran et al., 2014	OB (53)	60.1%	14.28	33.75	Inhibition	Stop Signal Task	4
	NW (50)	N.R.	13.81	23.83	Decision making	Delay Discounting Task	
Kulendran et al., 2016(1)	OB (20 & 45)	N.R.	36.2 & 44.3	39.7 & 43.4	Inhibition	Stop Signal Task	1
	NW (20)	N.R.	22.1	23.8	Decision making	Temporal discounting task	
Kulendran et al., 2016(2)	OB (47)	N.R.	14.3	33.2	Inhibition	Stop Signal Task	4

	NW (50)	N.R.	13.3	20.6	Decision making	Temporal discounting task	
Lawyer et al., 2015	OB (56)	65.5%	22.6	> 30	Inhibition	Stop Signal Task	2
	NW (235)	53.6%	21.5	18-24.9	Decision making	Delay Discounting task	
						Probability Discounting Task	
Loeber et al., 2012	OB (20)	65%	47.9	38.8	Inhibition	Go/No-Go	4
	NW (20)	65%	44.9	22.6	Working memory	Auditive verbal learning task	
Maayan et al., 2012	OB (54)	63%	17.5	39.9	Inhibition	Stroop	5
	NW (37)	56.8%	17.3	21.7	Cognitive flexibility	Trail Making Test B	
					Working memory	WRALM	
					Verbal fluency	Phonemic Verbal Fluency	
Mallorquí-Bague et al., 2016	OB (113)	77%	43.4	N.R.	Decision making	The Iowa Gambling Task	3
	NW (194)	79%	25	N.R.			
Mata et al., 2015	OB (32)	65.6%	15.1	29.4	Decision making	Risky-Gains task	4
	NW (22)	54.5%	15.5	21.2			
Mole et al., 2015	OB (30)	36.7%	44.1	32.7	Inhibition	Stop Signal Task	5
	NW (30)	36.7%	43.6	24.1	Decision making	Delay Discounting Task	
Moreno-Lo'pez et al., 2012	OB (36)	72.2%	14.2	28.5	Inhibition	Stroop	5
	NW (30)	56.3%	14.1	20.3			
Navas et al., 2016	OB (20)	55%	32.2	33.5	Decision making	The Iowa Gambling Task	5

	OW (21)	52.4%	35	27		Wheel of Fortune Task	
	NW (38)	57.9%	33.2	22.2			
Nederkoorn et al., 2006a	OB (15)	60%	13.9	25.8	Inhibition	Stop Signal Task	3
	NW (31)	61.3%	13.7	19.3	Decision making	Door Open Task	
Nederkoorn et al., 2006b	OB (31)	100%	40.9	39	Inhibition	Stop Signal Task	2
	NW (28)	100%	41.8	22.5	Decision making	Delay Discounting Task	
Nederkoorn et al., 2012	OW (14)	78.6%	8.1	21.1	Inhibition	Stop Signal Task	3
	NW (75)	50.7%	8.1	16.1			
Nederkoorn, 2014	OW (45)	84.4%	31	29.1	Inhibition	Stop Signal Task	4
	NW (73)	87.7%	29.2	21.7			
Perpiñá et al., 2016	OW (27)	85.2%	47.8	43.9	Cognitive flexibility	WCST	1
	NW (39)	76.9%	31.9	23.2	Decision making	The Iowa Gambling Task	
Pignatti et al., 2006	OB (20)	70%	43.4	42.2	Decision making	The Iowa Gambling Task	4
	NW (20)	50%	46.7	22.2			
Qavam et al., 2015	OW (40)	0%	16.5	N.R.	Planning	Tower of London Task	3
	NW (40)	0%	16.5	N.R.			
Restivo et al., 2017	OB (25)	92%	43.9	44.7	Inhibition	Stroop	2
	NW (20)	90%	43.8	22.4	Cognitive flexibility	WCST	
					Verbal fluency	Phonemic Verbal Fluency	

						Semantic Verbal Fluency	
Reyes et al., 2015	OW (69)	69 & 93	N.R.& 44%	10.2	Inhibition	Stroop	3 & 4
	NW (63)	63 & 92	N.R.&45.6%	10.3		Go/No-Go	
Ross et al., 2015	OB (79)	64.6%	19.6	35.6	Cognitive flexibility	Trail Making Test B-A	5
	NW (51)	49%	19.4	22.9	Working memory	Letter-Number Sequencing	
					Planning	VWMI	
						Tower of London Task	
Schiff et al., 2016	OB (23)	78.3%	36.2	36.2	Inhibition	Simon Task	4
	NW (23)	78.3%	22.8	22.4	Cognitive flexibility	Trail Making Test B	
					Working memory	Sternberg Task	
Sellaro and Colzato. 2017(1)	OW (17)	64.1%	23.4	27.7	Inhibition	Stop Signal Task	4
	NW (22)	77.3%	21.2	21.9			
Sellaro and Colzato. 2017(2)	OW (19)	57.9%	22.9	28.7	Inhibition	Simon Task	4
	NW (24)	79.2%	20.5	21.7			
Silveira et al., 2014	OW (9)	55.6%	25.4	28.3	Working memory	Letter-Number Sequencing	4
	NW (28)	57.1%	23.1	22.2			
Simmank et al., 2015	OW (26)	50%	27.1	34.7	Decision making	Delay Discounting Task	4
	NW (26)	50%	26	22.6			
Skoranski et al., 2013	OB (24)	N.R.	12.8	N.R.	Inhibition	Simon Task	3

	NW (27)	N.R.	12.8	N.R.			
Spitoni et al., 2017	OB (24)	79.2%	49.8	41.1	Inhibition	THSCT	1
	NW (37)	64.9%	35.7	22.5	Cognitive flexibility	The Rule Shift Cards	
Sting et al., 2012	OB (34)	70.6%	36.5	30.4	Working memory	Visual 1-back	5
	NW (34)	70.6%	38.4	22			
Sweat et al., 2017	OB (108)	63%	19.6	35.6	Inhibition	Stroop	5
	NW (54)	54%	19.4	21.5	Cognitive flexibility	Trail making Test B-A	
					Working memory	Letter-Number Sequencing	
					Verbal fluency	Semantic Verbal Fluency	
					Planning	Tower of London Task	
Tsai et al. 2016	OB (26)	30.8%	9.5	27.4	Inhibition	Serial Reaction Time Task	5
	NW (26)	30.8%	9.5	18.5			
Verdejo-García et al., 2010	OB (27)	45.8%	14.3	31.6	Inhibition	Five Digit test	5
	NW (34)	38.2%	15.3	21	Cognitive flexibility	Stroop task	
					Working memory	Trail Making Test B-A	
					Decision making	Letter-number Sequencing	
					Planning	The Iowa Gambling Task	
						Zoo Map	
Voon et al., 2014	OB (30)	45.8%	44.1	32.7	Inhibition	Premature Responding Task	5

	NW (30)	43.6%	43.6	24.1			
Weller et al., 2008(1)	OB (19)	0%	19.2	35.4	Decision making	Delay Discounting Task	5
	NW (21)	0%	19.4	22.3			
Weller et al., 2008(2)	OB (29)	100%	19.6	38.4	Decision making	Delay Discounting Task	5
	NW (26)	100%	20	21.9			
Wu et al., 2016	OB (19 & 18)	73.7%	N.R.	33	Inhibition	Stroop	3
	NW (20 & 18)	70%	N.R.	22.2	Cognitive flexibility	Trail making Test B-A	
					Working memory	Digit Span	
					Verbal fluency	Phonemic Verbal Fluency	
						Semantic Verbal Fluency	
Wu et al., 2017	OB (44)	31.8%	12.4	N.R.	Working memory	Digit Span	2
	OW (23)	26.1%	11.8	N.R.			
	NW (92)	56.5%	11.9	N.R.			
Yau et al., 2014	OB (30)	56.7%	17.6	35.5	Inhibition	Stroop	6
	NW (30)	63.3%	17.2	21.1	Cognitive flexibility	Trail making Test B	
					Working memory	WCST	
					Verbal fluency	Phonemic Verbal Fluency	
					Planning	Tower of London Task	
Yeomans et al., 2008	OW (31)	100%	N.R.	N.R.	Decision making	Delay Discounting Task	1

NW (116)	100%	N.R.	N.R.
----------	------	------	------

Note: N.R.= not reported; OB=obesity group; OW= overweight group; NW= normal weight group; RMCPT =Running Memory Continuous Performance Task; OTSCT= One-Touch Stockings of Cambridge Task; IDEDT= Intradimensional/ Extradimensional Task; CCPT= Conditioned cue preference test; DCCTT = Dimensional Change Card Sort Task; WRALM = Wide Range Assessment of Learning and Memory; VWMI=Visual Working Memory Index; THSCT= The Hayling Sentence Completion Test.

Table 1. Moderator analyses for studies of inhibition in obesity.

Variable	β	t	F	m	g	df	p
Participant Age	.032	0.65				14.3	.529
Percent Female	.064	1.31				11.4	.217
Participants							
BMI (adult sample)	.050	0.88				8.1	.404
Measures of inhibition			0.25			16.9	.862
Stop Signal task				10	-.402	9.7	.008
Go/No-go task				6	-.411	4.8	< .001
Stroop task				15	-.311	13.5	.008
Other				9	-.407	7.5	.005

Note: Significant ($p < .05$) moderating effects are listed in boldface font.

Table 2. Moderator analyses for studies of cognitive flexibility in obesity.

Variable	β	t	F	m	g	df	p
Participant Age	-.053	-0.92				7.4	.389
Percent Female Participants	.017	0.33				6.8	.754
BMI (adult sample)	-.127	-1.23				5.1	.272
Measures of cognitive flexibility			0.92			12.6	.425
Wisconsin Card Sorting Test				8	-.532	6.9	.036
Trail Making Test B/B-A				13	-.312	10.1	.002
Other				8	-.395	6.5	.014

Note: Significant ($p < .05$) moderating effects are listed in boldface font.

Table 3. Moderator analyses for studies of working memory in obesity.

Variable	β	t	F	m	g	df	p
Participant Age	.020	0.30				4.3	.778
Percent Female Participants	-.016	-0.29				7.0	.784
BMI (adult sample)	.106	1.00				4.3	.372
Measures of working memory			6.98			6.8	.023
Digit Span				6	-.346	3.2	.011
Letter–number sequencing				4	.026	2.7	.748
Other					-.478	10.5	< .001
				12			

Note: Significant ($p < .05$) moderating effects are listed in boldface font.

Table 4. Moderator analyses for studies of decision-making in obesity.

Variable	β	t	F	m	g	df	p
Participant Age	-.107	-1.80				18.4	.088
Percent Female Participants	-.100	-1.69				6.7	.138
BMI (adult sample)	-.047	-0.59				7.8	.573
Measure of decision-making			4.51			13.6	.032
Delay Discounting Task				11	-.416	10.6	< .001
The Iowa Gambling Task				12	-.580	10.2	< .001
Other				7	-.200	4.2	.044

Note: Significant ($p < .05$) moderating effects are listed in boldface font.