

Bioverse: Potentially Observable Exoplanet Biosignature Patterns Under the UV Threshold Hypothesis for the Origin of Life

MARTIN SCHLECKER ¹, DÁNIEL APAI ^{1, 2}, ANTONIN AFFHOLDER ³, SUKRIT RANJAN ^{2, 4}, RÉGIS FERRIÈRE ^{3, 5, 6},
KEVIN K. HARDEGREE-ULLMAN ¹, TIM LICHTENBERG ⁷ AND STÉPHANE MAZEVET ⁸

¹Steward Observatory, The University of Arizona, Tucson, AZ 85721, USA; schlecker@arizona.edu

²Lunar and Planetary Laboratory, The University of Arizona, Tucson, AZ 85721, USA

³Department of Ecology and Evolutionary Biology, University of Arizona, Tucson AZ, USA

⁴Blue Marble Space Institute of Science, Seattle, 98104, USA

⁵Institut de Biologie de l'École Normale Supérieure, ENS, PSL, Paris, France

⁶International Research Laboratory for Interdisciplinary Global Environmental Studies (iGLOBES), CNRS, ENS, PSL, University of Arizona, Tucson AZ, USA

⁷Kapteyn Astronomical Institute, University of Groningen, PO Box 800, 9700 AV Groningen, The Netherlands

⁸Observatoire de la Côte d'Azur, Université Côte d'Azur, Nice, France

ABSTRACT

A wide variety of scenarios for the origin of life have been proposed, with many influencing the prevalence and distribution of biosignatures across exoplanet populations. This relationship suggests these scenarios can be tested by predicting biosignature distributions and comparing them with empirical data. Here, we demonstrate this approach by focusing on the cyanosulfidic origins-of-life scenario and investigating the hypothesis that a minimum near-ultraviolet (NUV) flux is necessary for abiogenesis. Using Bayesian modeling and the Bioverse survey simulator, we constrain the probability of obtaining strong evidence for or against this “UV Threshold Hypothesis” with future biosignature surveys. Our results indicate that a correlation between past NUV flux and current biosignature occurrence is testable for sample sizes of $\gtrsim 50$ planets. The diagnostic power of such tests is critically sensitive to the intrinsic abiogenesis rate and host star properties, particularly maximum past NUV fluxes. Surveys targeting a wide range of fluxes, and planets orbiting M dwarfs enhance the chances of conclusive results, with sample sizes $\gtrsim 100$ providing $\gtrsim 80\%$ likelihood of strong evidence if abiogenesis rates are high and the required NUV fluxes are moderate. For required fluxes exceeding a few hundred erg/s/cm², both the fraction of inhabited planets and the diagnostic power sharply decrease. Our findings demonstrate the potential of exoplanet surveys to test origins-of-life hypotheses. Beyond specific scenarios, this work underscores the broader value of realistic survey simulations for future observatories (e.g., HWO, LIFE, ELTs, Nautilus) in identifying testable science questions, optimizing mission strategies, and advancing theoretical and experimental studies of abiogenesis.

1. INTRODUCTION

A wide variety of scenarios for the origin of life have been proposed (e.g., Baross & Hoffman 1985; Brasier et al. 2011; Mulkidjanian et al. 2012; Fox & Strasdeit 2013; Deamer & Georgiou 2015; Westall et al. 2018). While we may still be far from conclusively testing these scenarios, new prospects in the search for conditions favorable to life have opened up by thinking of the origin of life as a planetary phenomenon and identifying global-scale environmental properties that might support pathways to life (Sasselov et al. 2020). In particular, specific planetary conditions are needed to create stockpiles of initial compounds for prebiotic chemistry; and planetary processes are required to trigger the prebiotic synthesis.

Such planetary conditions can be hypothesized for exoplanets located in the habitable zone (HZ) of their host star, with persistent liquid water on their surface. For example, if deep-sea or sedimentary hydrothermalism is required for abiogenesis, then the insulation of an ocean from the planetary crust minerals (e.g., due to high-pressure ices) may reduce or eliminate the chances of life emerging (e.g., Baross & Hoffman 1985). The alternate scenario of a surface locally subject to wet-dry cycles requires a planetary exposure to mid-range Ultraviolet (UV) irradiation, as a source of energy and an agent of selection in chemical evolution (e.g., Deamer et al. 2019). This “UV Threshold Hypothesis” states that UV light in a specific wavelength range played a constructive role in

getting life started on Earth (Ranjan & Sasselov 2016; Ranjan et al. 2017a; Rimmer et al. 2018; Rapf & Vaida 2016), and it could provide a probabilistic approach to the interpretation of possible future biosignature detections (e.g., Catling et al. 2018a; Walker et al. 2018).

The association of chemical pathways to life and planetary environmental conditions offers a new opportunity to test alternate scenarios for life emergence based on planetary-level data collected from the upcoming observations of populations of exoplanets. Deep-sea hydrothermal scenarios require planetary conditions that may not be met on ocean worlds with large amounts of water, where the water pressure on the ocean floor is high enough to form high-pressure ices (Noack et al. 2016; Kite & Ford 2018). In this case, a testable prediction would be that planets with high-pressure ices do not show biosignatures. Likewise, if UV light is required to get life started, then there is a minimum planetary UV flux requirement to have an inhabited world. This requirement is set by competing thermal processes; if the photoreaction does not move forward at a rate faster than the competitor thermal process(es), then the abiogenesis scenario cannot function. On the other hand, abundant UV light vastly in excess of this threshold does not increase the probability of abiogenesis, since once the UV photochemistry is no longer limiting, some other thermal process in the reaction network will be the rate-limiting process instead. Therefore, a putative dependence of life on UV light is best described as a step function (e.g., Ranjan et al. 2017a; Rimmer et al. 2018, 2021a).

The goal of this work is to evaluate the potential of future exoplanet surveys to test the hypothesis that a minimum past NUV flux is required for abiogenesis. We focus on one version of the UV Threshold Hypothesis, the so-called cyanosulfidic scenario, which has been refined to the point where the required threshold flux has been measured to be $F_{\text{NUV},\text{min}} = (6.8 \pm 3.6) \times 10^{10} \text{ photons cm}^{-2} \text{ s}^{-1} \text{ nm}^{-1}$ integrated from 200–280 nm at the surface (Rimmer et al. 2018, 2021b; Rimmer 2023; Ranjan et al. 2023).

We follow a semi-analytical Bayesian analysis to estimate probabilities of obtaining strong evidence for or against this hypothesis. Under the UV Threshold Hypothesis (H_1), the probability of an exoplanet having detectable biosignatures is zero if the near-ultraviolet (near-Ultraviolet (NUV)) irradiation is less than the threshold, and it is equal to the (unknown) probability of **life emerging and persisting**, f_{life} if NUV exceeds the threshold for a sufficiently long period of time. Under the null hypothesis (H_{null}), that probability simply is f_{life} , that is, it does not correlate with the UV flux.

Figure 1 shows these hypotheses as derived from the predictions of the cyanosulfidic scenario. Given a sample of planets, where for some of them we have convincing biosignature detections but remain agnostic on f_{life} , we ask what evidence for H_1 and H_{null} we can expect to obtain.

A real exoplanet survey will be subject to observational biases and sample selection effects, and will be constrained by the underlying demographics of the planet sample. To assess the information gain of a realistic exoplanet survey, we employed Bioverse (Bixel & Apai 2021; Hardegree-Ullman et al. 2023; Schlecker et al. 2024; Hardegree-Ullman et al. 2024), a framework that integrates multiple components including statistically realistic simulations of exoplanet populations, a survey simulation module, and a hypothesis testing module to evaluate the statistical power of different observational strategies.

This paper is organized as follows: In Section 2, we introduce both our semi-analytical approach and Bioverse simulations for testing the UV Threshold Hypothesis. Section 3 presents the results of these experiments for a generic survey as well as for a realistic transit survey. In Section 4, we discuss our findings before concluding with a summary in Section 5.

2. METHODS

2.1. Fraction of inhabited planets with detectable biosignatures

Here, we conduct a theoretical experiment on the UV Threshold Hypothesis by relating the occurrence of life on an exo-earth candidate with a minimum past quiescent stellar UV flux, focusing on the prebiotically interesting NUV range from 200–280 nm (Ranjan & Sasselov 2016). Our core hypothesis shall be that life only occurs on planets that at some point in their history have received such radiation at a flux exceeding a threshold $F_{\text{NUV},\text{min}}$.

2.2. Semi-analytical approach

We first assessed the expected probabilities of obtaining true negative or true positive evidence for the UV Threshold Hypothesis (H_1) above, as well as the probability for misleading or inconclusive evidence, under idealized conditions. This serves as a first-order estimate of the information content of a survey, before we take into account the effects of exoplanet demographics, sample selection, and survey strategy.

Presumably, not all habitable worlds are inhabited and not all inhabited worlds develop detectable biosignatures. The fraction of exo-Earth candidates (EEC) that are both inhabited and exhibit detectable biosig-

Hypothesis: Life only originates on planets with particular UV irradiance




Figure 1. UV Threshold Hypothesis and null hypothesis derived from the cyanosulfidic scenario.

natures at the time of observation is unknown and is represented by the term f_{life} . This encompasses the probability of life both emerging and persisting to produce detectable biosignatures. Let us consider the probability to detect a biosignature $P(L)$, and let our observable be the inferred past NUV flux of the planet F_{NUV} . Under Hypothesis H_1 , there exists a special unknown value of F_{NUV} , noted $F_{\text{NUV},\text{min}}$ such that

$$P(L|F_{\text{NUV}}, H_1) = f_{\text{life}} \quad \text{if } F_{\text{NUV}} > F_{\text{NUV},\text{min}} \quad (1)$$

$$P(L|F_{\text{NUV}}, H_1) = 0 \quad \text{otherwise} \quad (2)$$

where f_{life} is the unknown probability of abiogenesis. The corresponding null hypothesis H_{null} is that there exists no such special value of F_{NUV} and that

$$P(L|F_{\text{NUV}}, H_{\text{null}}) = f_{\text{life}}. \quad (3)$$

In other words, H_{null} states that $P(L)$ is independent of F_{NUV} .

Defining a sample of size n as $X = \{F_{\text{NUV},i}, L_i\}_{i \in [1,n]}$ where L_i is equal to 1 if life is detected and 0 otherwise, we can calculate the evidence for hypothesis H_i being true against H_j through the Bayes factor

$$BF_{H_i, H_j} = \frac{P(X|H_i)}{P(X|H_j)}, \quad (4)$$

with $P(X|H_i)$ and $P(X|H_j)$ likelihoods of obtaining the sample X under either hypothesis.

Let $Y = \sum_i^n L_i$ be the random variable counting the number of positive life detections in a sample of size n . Its probability mass function under the null hypothesis H_{null} is that of a binomial distribution:

$$P(Y = k|H_{\text{null}}) = \binom{n}{k} f_{\text{life}}^k (1 - f_{\text{life}})^{n-k}. \quad (5)$$

Under H_1 , Y also follows a binomial distribution, however it is conditioned by $n_\lambda = n(\{F_{\text{NUV},i} \text{ if } F_{\text{NUV},i} > F_{\text{NUV},\text{min}}\}_{i \in [1,n]})$, the number of values of F_{NUV} in the experiment that exceed $F_{\text{NUV},\text{min}}$

$$P(Y = k|H_1) = \binom{n_\lambda}{k} f_{\text{life}}^k (1 - f_{\text{life}})^{n_\lambda - k}. \quad (6)$$

Hence,

$$BF_{H_1, H_{\text{null}}} = \frac{P(Y = k|H_1)}{P(Y = k|H_{\text{null}})} = \frac{\binom{n_\lambda}{k}}{\binom{n}{k}} (1 - f_{\text{life}})^{n_\lambda - n}, \quad (7a)$$

and

$$BF_{H_{\text{null}}, H_1} = \frac{P(Y = k|H_{\text{null}})}{P(Y = k|H_1)} = \frac{\binom{n}{k}}{\binom{n_\lambda}{k}} (1 - f_{\text{life}})^{n - n_\lambda}. \quad (7b)$$

Given a sample of planets, where for some of them we have convincing biosignature detections but remaining agnostic on f_{life} : What evidence for H_1 and H_{null} can we expect to get? The analytical expression for the Bayes factor of this inference problem (Equation 7a) is determined by the unknown variables f_{life} and $F_{\text{NUV},\text{min}}$, as well as by the summary statistic Y (number of biosignature detections). To compute the distribution of evidences, we repeatedly generated samples under H_1 and H_{null} and computed the Bayes factors $BF_{H_1, H_{\text{null}}}$ and $BF_{H_{\text{null}}, H_1}$. We then evaluated the fraction of Monte Carlo runs in which certain evidence thresholds (Jeffreys 1939) were exceeded.

Under a more realistic scenario, the distribution of n_λ depends on additional planetary properties and their evolution, as well as on observational biases and sample selection effects of the survey. We will address these in the following section.

2.3. Exoplanet survey simulations with Bioverse

To assess the diagnostic power of realistic exoplanet surveys, we employed our survey simulator and hypothesis testing framework Bioverse (Bixel & Apai 2021). The general approach is as follows:

- 1. Exoplanet population synthesis:** We populate the Gaia Catalogue of Nearby Stars (Smart et al. 2021) with synthetic exoplanets whose orbital parameters and planetary properties reflect our current understanding of exoplanet demographics (Bergsten et al. 2022; Hardegree-Ullman et al. 2023). Here, we also inject the demographic trend in question - in this case we assign biosignatures according to H_1 , i.e., to planets in the

HZ that have received NUV fluxes above a certain threshold.

2. Survey simulation: We simulate the detection and characterization of these exoplanets with a hypothetical survey, taking into account the survey’s sensitivity, target selection, and observational biases. To model the sensitivity of the information gain of a proposed mission to sample selection and survey strategy, we conduct survey simulations with Bioverse using different sample sizes and survey strategies.

3. Hypothesis testing: We evaluate the likelihood that a given survey would detect a specified demographic trend in the exoplanet population and estimate the precision with which the survey could constrain the parameters of that trend. A common definition of the null hypothesis H_{null} , which is also applied here, is that there is no relationship between the independent variable (here: maximum NUV flux) and the dependent variable (here: biosignature occurrence). The alternative hypothesis H_1 proposes a specific relationship between the independent and dependent variables. Bioverse offers either Bayesian model comparison or non-parametric tests to evaluate the evidence for or against the null hypothesis.

To determine the diagnostic capability of a given survey, Bioverse runs multiple iterations of the simulated survey and calculates the fraction of realizations that successfully reject the null hypothesis. We used this metric, known as the statistical power, to quantify the potential information content of the survey, identify critical design trades, and find strategies that maximize the survey’s scientific return.

2.3.1. Simulated star and planet sample

We generated two sets of synthetic exoplanet populations, one for FGK-type stars and one for M-type stars. The stellar samples are drawn from the Gaia Catalogue of Nearby Stars (Smart et al. 2021) with a maximum Gaia magnitude of 16 and a maximum stellar mass of $1.5 M_{\odot}$. We included stars out to a maximum distance d_{max} that depends on the required planet sample size. Planets were generated and assigned to the synthetic stars following the occurrence rates and size/orbit distributions of Bergsten et al. (2022). Following Bixel & Apai (2021), we considered only transiting EECs with radii $0.8 S^{0.25} < R < 1.4$ that are within the HZ (see Section 2.3.2). The lower limit was suggested as a minimum planet size to retain an atmosphere (Zahnle &

Catling 2017). For all survey simulations and hypothesis tests, we repeated the above in a Monte Carlo fashion to generate randomized ensembles of synthetic star and planet populations (Bixel & Apai 2021).

2.3.2. Habitable zone occupancy and UV flux

To construct a test the UV Threshold Hypothesis, we required that life occurs only on planets with sufficient past UV irradiation exceeding the origins of life threshold $F_{\text{NUV},\text{min}}$. Further, we required this flux to have lasted for a minimum duration ΔT_{min} to allow for a sufficient “origins timescale” (Rimmer 2023). All commonly investigated origins-of-life scenarios require water as a solvent; we thus considered only rocky planets that may sustain liquid water on their surface, i.e., that occupy their host star’s momentary HZ during the above period, as well as at the epoch of observation. To determine HZ occupancy, we took into account the evolution of the host star’s luminosity and HZ boundaries.

The HZ describes a region around a star where a planet with Earth’s atmospheric composition and climate feedbacks can maintain liquid water on its surface (e.g., Ramirez & Kaltenegger 2017; Ramirez 2018; Mol Lous et al. 2022; Spinelli et al. 2023; Tuchow & Wright 2023). Here, we adopted orbital distance estimates that define the HZ as the region between the runaway greenhouse transition, where the stellar instellation cannot anymore be balanced through infrared cooling to space (Ingersoll 1969), and the maximum greenhouse limit, corresponding to the maximum distance at which surface temperatures allowing liquid water can be maintained through a CO_2 greenhouse effect (Kasting 1991; Kasting et al. 1993; Underwood et al. 2003; Kopparapu et al. 2013, 2014). We used the parametrization in Kopparapu et al. (2014) to derive luminosity and planetary mass-dependent distance limits of the HZ a_{inner} and a_{outer} .

To determine HZ occupancy, we interpolated the stellar luminosity evolution grid of Baraffe et al. (1998) using a Clough Tocher interpolant (Nielson 1983; Alfeld 1984, see left panel of Figure 2) to compute the evolution of the inner (runaway greenhouse) and outer (maximum greenhouse) edges as a function of planet mass and stellar spectral type (Kopparapu et al. 2014). Being a local interpolation method, Clough Tocher enables rapid processing while producing a smooth interpolating surface that highlights local trends. From this, we get each planet’s epochs within and outside the HZ.

For the NUV flux, we used the age- and stellar mass-dependent NUV fluxes in the HZ obtained by Richey-Yowell et al. (2023), which considers GALEX UV data in the wavelength range of 177–283 nm. We linearly inter-

polate in their measured grid, where we convert spectral type to stellar mass using the midpoints of their mass ranges (0.75 M_{\odot} for K stars, 0.475 M_{\odot} for early-type M stars, and 0.215 M_{\odot} for late-type M stars). Outside the age and stellar mass range covered in Richey-Yowell et al. (2023), we extrapolate using nearest simplex (see right panel of Figure 2).

We then determined which planets were both in the HZ and had NUV fluxes above $F_{\text{NUV},\text{min}}$. To avoid considering short transitional phases, we require this situation to last for a minimum duration $\Delta T_{\text{min}} \geq 1$ Myr. We assigned the emergence **and persistence** of life to a random fraction f_{life} of all temperate planets fulfilling these requirements. For the probability of a planet having detectable biosignatures, $P(\text{bio})$, the UV Threshold Hypothesis then states

$$H_1 : P(\text{bio}) = \begin{cases} 0, & F_{\text{NUV}} < F_{\text{NUV},\text{min}} \\ f_{\text{life}}, & F_{\text{NUV}} \geq F_{\text{NUV},\text{min}} \\ & \text{and in HZ for } \Delta t \geq 1 \text{ Myr} \end{cases} \quad (8)$$

and the corresponding null hypothesis $H_{\text{null}} : P(\text{bio}) = f_{\text{life}}$, i.e., no correlation with UV flux.

2.3.3. Transit survey simulations

With the synthetic star and planet samples generated, we used Bioverse’s survey module to simulate noisy measurements of key observables with a transit survey. We assumed a hypothetical mission that can target a large planet sample with high photometric precision and conduct a biosignature search on these planets (e.g., Apai et al. 2019, 2022). The simulated survey was designed to measure planetary instellation (for HZ occupancy) with a precision of 5% and host star effective temperature with a precision of 50.0 K. The maximum past NUV flux a planet received can be determined within a precision of 5%. To marginalize over choices of biosignatures and their detectability, which are beyond the scope of this study, we assumed that any inhabited planet would show a biosignature detectable by the survey.

2.3.4. Hypothesis testing

To evaluate the evidence for correctly rejecting the null hypothesis, we employed the Mann-Whitney U test (Mann & Whitney 1947). This is a non-parametric test of the null hypothesis that two independent samples were drawn from a population with the same distribution and is in particular sensitive to one sample being stochastically greater than the other. We used the Mann-Whitney U test to compare the distributions of NUV fluxes of planets with and without biosignatures. The implementation in Bioverse relies on the

`scipy.stats.mannwhitneyu` function (Virtanen et al. 2020) and returns a p-value, for which we set a significance level of $\alpha = 0.05$ to reject the null hypothesis. For every hypothesis test, we repeated randomized survey realizations to estimate the fraction of successful rejections of the null hypothesis, i.e., the statistical power of the survey.

3. RESULTS

3.1. Semi-analytical assessment

In Section 2.2, we computed the probability for true positive evidence for H_1 and H_{null} , respectively (Equations 7a, 7b). Figure 3 shows how these evidences are distributed for sample sizes 10 and 100, and how likely we are to obtain strong evidence ($BF_{H_1, H_j} > 10$). For $n = 10$, strong true evidence for H_1 (H_{null}) can be expected in $\sim 30\%$ ($\sim 40\%$) of all random experiments. In the majority of cases, the outcome of the survey will be inconclusive. The situation improves with larger samples: for $n = 100$, 80% of random samples permit conclusive inference (strong true evidence) under either H_1 or H_{null} .

The expected resulting evidence further depends on the a priori unknown **rate of life’s emergence and persistence** f_{life} and on the NUV flux threshold. Figure 4 illustrates this dependency: For very low values of either parameter, samples drawn under the null or alternative hypotheses are indistinguishable and the Bayesian evidence is always low. Both higher f_{life} and higher NUV flux thresholds increase the probability of obtaining strong evidence. Larger sample sizes enable this at lower values of these parameters.

So far, we have assumed random, uniform distributions of f_{life} , $F_{\text{NUV},\text{min}}$, and F_{NUV} . A high biosignature detection rate f_{life} increases the evidence (see Equation 7a) but a survey strategy cannot influence it. The same is true for $F_{\text{NUV},\text{min}}$, where again higher values increase the evidence as the binomial distribution for H_1 gets increasingly skewed and shifted away from the one for H_{null} . However, one might select exoplanets for which a biosignature test is performed based on a *priori* available contextual information (Catling et al. 2018b) in order to maximize the science yield of investing additional resources. For instance, the distribution of F_{NUV} in the planet sample can be influenced by the survey strategy, and a targeted sampling approach could favor extreme values. We model this by distributing F_{NUV} according to different *Beta* functions and introduce a selectivity parameter $s \in]-1, 1[$ such that $F_{\text{NUV}} \sim \text{Beta}(1/10^s, 1/10^s)$. Figure 5 shows how the probability of obtaining true strong evidence for H_1 scales with selectivity s . For large samples, a high selec-




Figure 2. Interpolated stellar luminosity evolution (left) and evolution of the NUV flux in the HZ (right) as a function of host star mass. Scatter points show age and host star mass of the transiting planets in the synthetic planet sample; crosses denote the estimated NUV values in Richey-Yowell et al. (2019). We show three evolutionary tracks for a threshold flux of $F_{\text{NUV},\text{min}} = 300.0 \text{ erg s}^{-1} \text{ cm}^{-2}$ that occupy the HZ (yellow sections) and exceed the threshold NUV flux (red sections) at different times. Where these sections overlap (white rectangles), the requirements for abiogenesis are met and we assign a biosignature detection with probability f_{life} . Planet 1 is an EEC orbiting a K dwarf that never receives sufficient NUV flux for abiogenesis. Planet 2 and Planet 3 enter the HZ at different times and receive sufficient NUV flux for different durations until their respective host star evolves below the threshold.




Figure 3. Obtaining true strong evidence with different sample sizes. Left: Probability to reach given evidence levels for H_1 and H_{null} under sample sizes $n = 10$ (solid) and $n = 100$ (dashed). Vertical lines denote thresholds for “strong” evidence, $BF_{H_i, H_j} > 10$, and “extreme” evidence, $BF_{H_i, H_j} > 100$. Right: Probability of obtaining true strong evidence for H_1 as a function of sample size n .

tivity ($s \sim 1$) can increase the probability of obtaining true strong evidence from $\sim 70\%$ for $s = 0$ (random uniform distribution) to $> 90\%$.

3.2. Survey simulations with Bioverse

With HZ occupancy as a requirement for abiogenesis, and barring selection biases beyond stellar brightness, the host star distribution of inhabited planets in a simulated transit survey is skewed toward later spectral types. For a fixed planet sample size, the fraction of inhabited planets is highest for planets orbiting M dwarfs due to the higher NUV fluxes in the HZ of these stars

(see Figures 2, 6). Their NUV fluxes are generally highest at early times $\lesssim 100 \text{ Myr}$. These host stars, in particular late subtypes, also provide extended periods of increased NUV emission that overlap with times when some of these planets occupy the HZ (see Figure 2), our requirement for abiogenesis (see Equation 8). Thus – under the UV Threshold Hypothesis – most inhabited transiting planets in the sample orbit M dwarfs.

Here, we are interested in the statistical power of a transit survey with a plausible sample selection and size. In the following, we fix the sample size to 250 and con-




Figure 4. Probability of obtaining true strong evidence for different abiogenesis rates, NUV flux thresholds, and sample sizes. For each of these parameters, higher values increase the probability of yielding strong evidence.




Figure 5. Scaling of the probability of obtaining true strong evidence with sample selectivity. Left: Sampling distribution for different selectivity parameters s . Right: Resulting $P(\text{true strong evidence})$, where f_{life} and $F_{\text{NUV},\text{min}}$ are randomly drawn from a uniform distribution. Sampling more extreme values of F_{NUV} is more likely to yield strong evidence.

sider two different survey strategies targeting FGK and M dwarfs, respectively. We further investigate the sensitivity of the survey to the a priori unknown threshold NUV flux $F_{\text{NUV},\text{min}}$ and the **probability of life emerging and persisting** f_{life} .

3.2.1. Selectivity of simulated transit surveys

In Section 3.1, we showed that the probability of obtaining true strong evidence for the hypothesis that life only originates on planets with a minimum past NUV flux is sensitive to the distribution of sampled past NUV fluxes, i.e., the selectivity of the survey (compare Figure 5). For both surveys targeting M dwarfs and those targeting FGK dwarfs, the maximum NUV distribution is rather unimodal. Applying the approach from Sect. 3.1 of fitting a Beta function to the distribution, we find rather low selectivities (see Figure 6), which is likely detrimental for statistical hypothesis tests.

3.2.2. Expected biosignature pattern

A representative recovery of the injected biosignature pattern is shown in Figure 6. There, we assumed an

abiogenesis rate of $f_{\text{life}} = 0.8$ and a minimum NUV flux of $F_{\text{NUV},\text{min}} = 300.0 \text{ erg s}^{-1} \text{ cm}^{-2}$. All injected biosignatures are assumed to be detected without false positive ambiguity, and the maximum NUV flux is estimated from the host star's spectral type and age with an uncertainty corresponding to the intrinsic scatter in the NUV fluxes in Richey-Yowell et al. (2023). This leads to a distribution of biosignature detections with detections increasingly occurring above a threshold inferred NUV flux. In this example case, the few biosignature detections in the FGK sample lead to a higher evidence than in the M dwarf sample, where the majority of planets are above the threshold NUV flux.

Figure 7 shows the fraction of inhabited planets under the UV Threshold Hypothesis for different threshold NUV fluxes and for the limiting case of a **probability for life's emergence and persistence** of $f_{\text{life}} = 1$. This fraction decreases sharply with increasing threshold flux, as fewer planets receive sufficient NUV flux for abiogenesis. Another effect responsible for this drop is that some planets receive the required NUV flux only




Figure 6. Simulated transit surveys targeting FGK and M stars.

Top: Host stars of all transiting EECs and inhabited planets in a simulated transit survey. In the FGK sample, all EECs and all inhabited planets orbit K dwarfs. In an M dwarf sample of the same size, the fraction of inhabited planets is larger.

Center: Distribution of inferred maximum past NUV flux in transit surveys targeting EECs around FGK and M stars, respectively. The best-fit beta distributions correspond to selectivities of $s_{\text{FGK}} = -0.47$ and $s_{\text{M}} = -0.02$. Red areas show inhabited planets for an abiogenesis rate of $f_{\text{life}} = 0.8$ and a generic threshold NUV flux $F_{\text{NUV},\text{min}} = 300.0 \text{ erg s}^{-1} \text{ cm}^{-2}$.

Bottom: Recovered biosignature detections and non-detections of simulated transit surveys. The dashed line denotes $F_{\text{NUV},\text{min}}$.




Figure 7. Fraction of inhabited planets for different threshold NUV fluxes under the UV Threshold Hypothesis if the abiogenesis rate $f_{\text{life}} = 1$. For all samples, the fraction of inhabited planets drops sharply with increasing threshold NUV flux due to the combined effects of never receiving sufficient NUV flux for abiogenesis or receiving it before entering the HZ.

before entering the HZ – this is especially likely for M dwarfs. For the FGK sample, the fraction of inhabited planets drops at lower threshold fluxes than for the M dwarf sample.

3.2.3. Statistical power and sensitivity to astrophysical parameters

We now investigate the sensitivity of the achieved statistical power of our default transit survey to the a priori unconstrained threshold NUV flux $F_{\text{NUV},\text{min}}$ and the abiogenesis **and persistence** rate f_{life} . Figure 8 shows the statistical power as a function of these parameters for a sample size of $N = 250$. Values of $F_{\text{NUV},\text{min}}$ that lie between the extrema of the inferred maximum NUV flux increase the achieved statistical power of the survey, as in this case the dataset under the alternative hypothesis H_1 differs more from the null hypothesis. Furthermore, **higher** f_{life} **increase** the evidence for H_1 .

Parameter space regions with statistical power above 90% lie at $f_{\text{life}} > 0.5$ and mostly at threshold NUV fluxes of $\sim 200\text{--}400 \text{ erg s}^{-1} \text{ cm}^{-2}$. Notably, the sensitivity of the M dwarf sample extends into the low NUV flux end due to the broader distribution of maximum past NUV fluxes in this sample. Here, the FGK sample is barely sensitive.

4. DISCUSSION

A key question in the quest to understand the origins of life is which natural processes best explain how living matter spontaneously appears from nonliving matter (e.g., Malaterre et al. 2022). Using astronomical methods, this question will likely not be testable for individual planets but rather for ensembles of planets.

The cyanosulfidic scenario for the origins of life (Patel et al. 2015), in particular its predicted existence of a minimum NUV flux required for prebiotic chemistry, offers an opportunity to test an origins of life hypothesis with a statistical transit survey sampling planets with varying NUV flux histories. In the following, we discuss the prospects of testing the UV Threshold Hypothesis in light of our results.

4.1. Sampling strategy for testing a NUV flux threshold

In Sect. 3.1, we show that testing the UV Threshold Hypothesis suffers from ‘nuisance’ parameters that hamper inference through astronomical observations. Here, these parameters are the unspecified value of the NUV threshold hypothesized to exist under H_1 , and the unknown probability of detectable life emerging on a habitable planet f_{life} . While the inference of a planet’s entire UV flux evolution is difficult (e.g., Richey-Yowell et al. 2023), the estimated maximum NUV flux that a planet was exposed to may be used as a proxy, at least if one is interested in a minimum threshold flux and makes the assumption that planetary surfaces offer protection against *too high* UV flux. Indeed, the distribution of the number of planets with detected biosignature in a particular sample of planets with inferred maximum NUV values F_{NUV} depends on both the values of $F_{\text{NUV},\text{min}}$ and f_{life} as shown in equation 6.

In our semi-analytical analysis (Section 3.1), we project a possible test performed by a future observer equipped with a sample of exoplanets with derived past maximum NUV exposure for which biosignature detection has been attempted. This is necessarily reductive as this observer will have more knowledge about experimental conditions and will therefore be able to use this information to guide hypothesis testing. For instance, we have made the choice to consider the total number of detected biosignatures as our summary statistic (Equation 6), which is not sufficient to infer $F_{\text{NUV},\text{min}}$ and f_{life} separately. However, by conditioning the Bayes factor to these variables (Equation 7a), we calculate the probability distribution of the Bayesian evidence in favor of H_1 . In doing so, we may evaluate how evidence depends on the uncertainty over these unknown parameters in general terms, without assuming which particular test a future observer might actually choose to perform over real data when available. From this, we can see that target selection can strongly affect the conclusiveness of a future test of the UV Threshold Hypothesis.

The particular finding that prioritizing extreme values of past NUV flux can enhance statistical power likely clashes with observational constraints, as the composition of the subset of planets that we can observe and




Figure 8. Statistical power as a function of threshold NUV flux and abiogenesis rate. Even for a large sample (here: $N = 250$), a high statistical power of the transit survey requires high **rates of life emerging and persisting** f_{life} . Intermediate values of $F_{\text{NUV},\text{min}}$ are more likely to yield strong evidence than extreme values. **For** $f_{\text{life}} \gtrsim 80\%$, the sensitivity of the M dwarf sample extends into the low NUV flux end.

for which detection of biosignature can be attempted is not independent from their NUV flux history. Hence, for our future observer, selectivity and sample size may be in conflict. This trade-off can be quantified in terms of expected evidence yield, which we have done in Section 3.1. Our analysis shows that regardless of selectivity, sample sizes smaller than 50 likely result in inconclusive tests, and that increasing selectivity towards extreme F_{NUV} offers limited inference gains compared to the uniform case ($s = 0$; Figure 5). For larger samples, however, a narrow distribution of F_{NUV} may prevent inference entirely. We thus argue that selecting a sample with F_{NUV} distributed uniformly or emphasizing extreme values should – barring any practical counterarguments – be considered in any future attempt at testing the UV Threshold Hypothesis. Since the practical implementation of an exoplanet survey can stand in the way of such a selection, the following discussion focuses on the results of our transit survey simulations with Bioverse.

4.2. How planetary context may constrain the UV Threshold Hypothesis

It comes to no surprise that the success rate for testing the UV Threshold Hypothesis is sensitive to the sample size of the survey and to the occurrence of life on temperate exoplanets. As we have shown, the statistical power of this test also depends on the distribution of past NUV fluxes in the sample and on the threshold flux. Optimizing the survey to sample a wide range of NUV flux values, particularly at the extremes, can enhance the likelihood of obtaining strong evidence for or against the hypothesis. Intermediate values of the threshold

NUV flux are more likely to yield strong evidence than extreme values, as the dataset under the alternative hypothesis H_1 differs more from the null hypothesis in this case while still being sufficiently populated. The threshold flux is, of course, a priori unknown and we cannot influence it. If, however, better theoretical predictions for the required NUV flux for abiogenesis become available (Rimmer et al. 2021a), the survey strategy can be further optimized, for instance by targeting planets that are estimated to have received a NUV flux slightly below and above this threshold or by applying a bisection algorithm in a sequential survey (Fields et al. 2023).

4.3. An M dwarf opportunity

An interesting aspect lies in the distribution of host star properties, as different spectral types probe different past NUV flux regimes. FGK stars show a narrow distribution of maximum past NUV fluxes in the HZ, which may – depending on the (unknown) threshold NUV flux – limit the diagnostic power of a survey. In the case of a pure FGK sample, it will essentially only be sensitive to NUV flux thresholds $\sim 200\text{--}400\text{ erg s}^{-1}\text{ cm}^{-2}$, and the chance of detecting biosignatures diminishes rapidly for higher thresholds within this range (see Figure 7). Detected biosignatures in an FGK sample would have little constraining power on testing the UV Threshold Hypothesis; they would either suggest that low NUV fluxes are sufficient for abiogenesis or indicate a different abiogenic pathway (e.g., Westall et al. 2018). A lack of biosignatures in a larger FGK sample would support the UV Threshold Hypothesis.

On the other hand, M dwarfs show a wider distribution of maximum past NUV fluxes in their HZs. While old M dwarfs can be considered low-UV environments, a significant fraction of them emit high NUV fluxes into their HZ during their early stages, in particular later subtypes (Richey-Yowell et al. 2023). This will help to test the high NUV flux end of the UV Threshold Hypothesis; a higher occurrence of biosignatures here would support the hypothesis that a higher NUV flux is favorable or necessary for life. At the same time, a fraction of host stars in our M dwarf sample extends it to lower maximum past NUV fluxes, enabling tests of the low NUV flux end of the hypothesis. The higher and more variable NUV fluxes in M dwarfs thus increase the likelihood of obtaining strong evidence for or against the UV Threshold Hypothesis.

The combination of a lack of UV radiation today, which makes biosignature gases more detectable (Sengura et al. 2005), and a UV-rich past that may have enabled abiogenesis could make M dwarfs the preferred targets for biosignature searches. We note that relevant mission concepts, such as the Large Interferometer for Exoplanets (LIFE, Quanz et al. 2022; Glauser et al. 2024), include M-dwarf systems among their primary targets (Kammerer & Quanz 2018; Carrión-González et al. 2023). Our findings underscore the importance of constraining the UV emission profiles of EEC host stars throughout their evolutionary stages to assess the viability of M-dwarf planets as testbeds for theories on the origins of life (Rimmer et al. 2021a; Ranjan et al. 2023).

4.4. Sensitivity to astrophysical parameters

Our Bioverse simulations that take into account exoplanet demographics, the evolution of habitability and NUV fluxes, and observational biases show that not only the likelihood of a conclusive test of the UV Threshold Hypothesis, but also the likelihood of successful biosignature detection itself is extremely sensitive to the threshold NUV flux if the hypothesis is true. Even if all biosignatures can be detected and the nominal **rate of life's emergence and persistence** is very high, say $f_{\text{life}} = 1$, under the condition that prebiotic chemistry requires a minimum NUV flux *and* liquid water, if the threshold flux turns out to be high the probability of finding life on a randomly selected planet may be very low. As we showed, for high required fluxes the two requirements of simultaneous HZ occupancy and sufficient NUV flux conspire to diminish the fraction of inhabited planets in the sample. Taking the inferred fluxes from Richey-Yowell et al. (2023) at face value (but taking into account intrinsic scatter), a minimum required NUV flux

of $\gtrsim 400 \text{ erg s}^{-1} \text{ cm}^{-2}$ reduces the fraction of inhabited planets to below $\sim 1\%$. This not only calls for a large sample size and a targeted sample selection preferring high expected past NUV fluxes, but also highlight the necessity of continued theoretical and experimental research into the role of UV radiation in prebiotic chemistry (Ranjan et al. 2017b; Rimmer et al. 2018, 2021a).

4.5. Contextual support for potential biosignature detections

The predicted interplay of NUV flux and HZ occupancy in enabling abiogenesis via the cyanosulfidic scenario could in principle be used to add or remove credibility from a tentative biosignature detection. For example, with a strong belief that this scenario is the only viable one for the origins of life, a biosignature detection on a planet orbiting a strongly UV-radiating star may add credibility to the detection. Conversely, a biosignature detection on a planet estimated to have received very little UV radiation would increase the likelihood of a false positive detection. On the other hand, should the detection in the latter case be confirmed, it could be used to falsify the UV Threshold Hypothesis.

Our simulations find no clear criterion for the credibility of a biosignature detection based on spectral type of the host star, as both FGK and M dwarf samples show similar maximum past NUV flux distributions. The abiogenesis rate in both samples show similar trends with the threshold NUV flux, and the fraction of inhabited planets drops at similar threshold fluxes (see Section 3.2). A potentially inhabited planet's host star spectral type may thus not be a strong indicator for the credibility of a biosignature detection in the context of the UV Threshold Hypothesis.

4.6. Overall prospects for testing the UV Threshold Hypothesis

Our results show that the UV Threshold Hypothesis is testable with potential future exoplanet surveys, but that the success of such a test depends on the sample size, the distribution of past NUV fluxes, and several unknown astrophysical nuisance parameters. Even under idealized conditions, obtaining strong evidence for or against the hypothesis likely requires sample sizes on the order of 100 (see Section 3.1). This is true for a future transit survey, the specifics of which we have reflected in our Bioverse simulations (see Section 3.2). However, we have shown that the impacts from the combined requirements of the UV Threshold Hypothesis on the fraction of inhabited planets in a sample are comparable in the non-transiting case.

Given the challenging nature of detecting and characterizing small (Earth-sized) exoplanets, most exoplanet

mission concepts currently under development or considered lack the potential for characterizing large enough samples. Ground-based 25–40-meter class extremely large telescopes are expected to have the capabilities to detect biosignatures on exoplanets like Proxima Centauri b (e.g., Wang et al. 2017; Hawker & Parry 2019; Zhang et al. 2024; Vaughan et al. 2024). Hardegree-Ullman et al. (2024) used Bioverse to determine potential yields for a 10-year direct imaging and high-resolution spectroscopy survey of O₂ on the Giant Magellan Telescope (GMT) and on the Extremely Large Telescope (ELT) and found that between 7 and 19 habitable zone Earth-sized planets could be probed for Earth-like oxygen levels. Such a sample is too small to test the UV Threshold Hypothesis, but it may be synergistic with other detection methods.

The Habitable Worlds Observatory (HWO) is expected to characterize a sample of ~ 25 Earth analogs (Mamajek & Stapelfeldt 2023; Tuchow et al. 2024). Depending on the technical design, *LIFE* is expected to target 25–80 EECs (Kammerer & Quanz 2018; Quanz et al. 2022), which could be just sufficient to constrain the UV Threshold Hypothesis. One current exception is the Nautilus Space Observatory concept (Apai et al. 2019, 2022). Nautilus aims to characterize up to ~ 1000 EEC via transmission spectroscopy, building on an innovative optical technology. To guide the definition of future biosignature surveys, it is important to refine predictions on the role of UV radiation in prebiotic chemistry with both theoretical and experimental work.

4.7. Caveats

Our work is based on a number of assumptions and simplifications that may affect the results and conclusions. We discuss some of these caveats here.

Ranjan et al. (2017b) speculated that UV photochemistry might be rate-limiting for abiogenesis, particularly on planets orbiting M dwarfs, due to their lower baseline UV fluxes. This could delay abiogenesis by orders of magnitude, resulting in a continuous dependence of abiogenesis likelihood on UV flux. However, recent studies challenge this view for the cyanosulfidic scenario. Rimmer et al. (2021a) calculated photochemical timescales on early Earth at 180–300 hours (7.5–12.5 days), significantly shorter than the 1000-year timescale for stochastic geological events (Rimmer 2023). Even with 1000x slower photochemistry on M-dwarf planets, abiogenesis would occur within 20–30 years — geologically negligible compared to stochastic processes — supporting the step-function model.

Nonetheless, alternative abiogenesis pathways or combinations of pathways may exhibit continuous or mixed dependencies on UV flux. While the step-function formalism is justified for the cyanosulfidic scenario, future work should explore UV dependencies across other scenarios to refine predictions for biosignature distributions and testable hypotheses.

4.7.1. The UV Threshold Hypothesis as a narrow step function

A key aspect of the UV Threshold Hypothesis is the proposed step-function dependence of abiogenesis likelihood on UV flux. This approach stems from the chemical constraints governing photochemical pathways, which exhibit a threshold behavior: below a certain flux, competing thermal reactions dominate, preventing abiogenesis, while above the threshold, UV photochemistry proceeds at sufficient rates, and other stochastic processes become rate-limiting.

Ranjan et al. (2017b) speculated that UV photochemistry might be rate-limiting for abiogenesis, particularly on planets orbiting M dwarfs, due to their lower baseline UV fluxes. This could delay abiogenesis by orders of magnitude, resulting in a continuous dependence of abiogenesis likelihood on UV flux. However, recent studies challenge this view for the cyanosulfidic scenario. Rimmer et al. (2021a) calculated photochemical timescales on early Earth at 180–300 hours (7.5–12.5 days), significantly shorter than the 1000-year timescale for stochastic geological events (Rimmer 2023). Even with 1000x slower photochemistry on M-dwarf planets, abiogenesis would occur within 20–30 years — geologically negligible compared to stochastic processes — supporting the step-function model.

Nonetheless, alternative abiogenesis pathways or combinations of pathways may exhibit continuous or mixed dependencies on UV flux. While the step-function formalism is justified for the cyanosulfidic scenario, future work should explore UV dependencies across other scenarios to refine predictions for biosignature distributions and testable hypotheses.

4.7.2. Existence of an atmosphere-crust interface

By its nature, cyanosulfidic scenario relies on rock surfaces exposed to the planetary atmosphere. Water worlds that have their entire planetary surface covered by oceans contradict this requirement and do not allow for the wet-dry cycling inherent to this origin of life scenario. The competition of tectonic stress with grav-

844 itational crustal spreading (Melosh 2011) sets the max-
 845 imum possible height of mountains, which in the solar
 846 system does not exceed ~ 20 km. Such mountains will
 847 be permanently underwater on water worlds. Another
 848 impediment to wet-dry cycles may be tidal locking of
 849 the planet as it stalls stellar tide-induced water move-
 850 ment and diurnal irradiation variability (e.g., Ranjan
 851 et al. 2017b). However, recent dynamical models sug-
 852 gest tidally locked planets to undergo rapid drift of their
 853 sub-stellar point (Revol et al. 2024).

854 4.7.3. Stellar flares

855 Our assumptions on past UV flux neglect the contri-
 856 bution of stellar flares, which may be hypothesized as an
 857 alternative source of UV light (Buccino et al. 2007; Ran-
 858 jan et al. 2017a). This concerns mainly ultracool dwarfs,
 859 due to their low quiescent emission and high pre-main
 860 sequence stellar activity (Buccino et al. 2007; West et al.
 861 2008). However, recent work indicates that the majority
 862 of stars show inadequate activity levels for a sufficient
 863 contribution through flares (Glazier et al. 2020; Ducrot
 864 et al. 2020; Günther et al. 2020).

865 4.7.4. Atmosphere transmission

866 We do not take into account absorption of UV radia-
 867 tion by the planetary atmosphere. Theoretical work sug-
 868 gests that the atmosphere of prebiotic Earth was largely
 869 transparent at NUV wavelengths with the only known
 870 source of attenuation being Rayleigh scattering (Ran-
 871 jan & Sasselov 2017; Ranjan et al. 2017a). We thus
 872 approximated surface UV flux using top-of-atmosphere
 873 fluxes. If there are planets in a sample that do not have
 874 a transparent atmosphere at NUV wavelengths and re-
 875 quire higher fluxes for abiogenesis, the fraction of inhab-
 876 ited planets in the sample will be lower. However, these
 877 planets will not pollute the below-threshold subsample,
 878 as they will not be able to host life under the UV Thresh-
 879 old Hypothesis. Exoplanet surveys focusing on highly
 880 irradiated planets offer an opportunity to constrain the
 881 typical oxidation state of rocky exoplanets, providing
 882 insights into the average composition of their secondary
 883 atmospheres (Lichtenberg & Miguel 2024). This is par-
 884 ticularly relevant for prebiotic worlds, as varying oxi-
 885 dation states significantly perturb the classical habit-
 886 able zone concept (Nicholls et al. 2024) and also influ-
 887 ence surface UV levels through changing atmospheric
 888 transmission. Optimally, the atmospheric composition
 889 of young rocky protoplanets will be probed to constrain
 890 the possible range of atmospheric and mantle oxidation
 891 states during early planetary evolution by future direct
 892 imaging concepts (Cesario et al. 2024).

893 5. CONCLUSIONS

894 We propose that specific origins-of-life scenarios may
 895 leave a detectable imprint on the distribution of biosig-
 896 natures in exoplanet populations. We have investigated
 897 the potential of upcoming exoplanet surveys to test the
 898 hypothesis – motivated by the cyanosulfidic origins-of-
 899 life scenario – that a minimum past NUV flux is required
 900 for abiogenesis. To this end, we first employed a semi-
 901 analytical Bayesian analysis to estimate probabilities of
 902 obtaining strong evidence for or against this hypothesis.
 903 We then used the Bioverse framework to assess the diag-
 904 nostic power of realistic transit surveys, taking into ac-
 905 count exoplanet demographics, time-dependency of hab-
 906 itability and NUV fluxes, observational biases, and tar-
 907 get selection.

908 Our main findings are:

- 909 1. The UV Threshold Hypothesis of the cyanosulfidic
 910 scenario for the origins of life should lead to a cor-
 911 relation between past NUV flux and current occur-
 912 rence of biosignatures that may be observationally
 913 testable.
- 914 2. The required sample size for detecting this correla-
 915 tion depends on the abiogenesis rate on temperate
 916 exoplanets and the distribution of host star prop-
 917 erties in the sample; in particular their maximum
 918 past NUV fluxes. Samples smaller than 50 planets
 919 are unlikely to yield conclusive results.
- 920 3. Under the UV Threshold Hypothesis, the fraction
 921 of inhabited planets in a transit survey is sensi-
 922 tive to the threshold NUV flux and is expected to
 923 drop sharply for required fluxes above a few hun-
 924 dred $\text{erg s}^{-1} \text{cm}^{-2}$.
- 925 4. If the predicted UV correlation exists, obtain-
 926 ing strong evidence for the hypothesis is likely
 927 ($\gtrsim 80\%$) for sample sizes ≥ 100 if the abiogen-
 928 esis rate is high ($\gtrsim 50\%$) and if no very high NUV
 929 fluxes are required. A survey strategy that targets
 930 extreme values of inferred past NUV irradiation
 931 increases the diagnostic power.
- 932 5. Samples of planets orbiting M dwarfs overall
 933 yield higher chances of successfully testing the
 934 UV Threshold Hypothesis. They may also be more
 935 likely to yield biosignature detections under this
 936 hypothesis.

937 Overall, our work demonstrates that future exoplanet
 938 surveys have the potential to test the hypothesis that
 939 a minimum past NUV flux is required for abiogenesis.
 940 More generally, we found that models of the origins of
 941 life provide hypotheses that may be testable with these
 942 surveys. Conducting realistic survey simulations with

representative samples is important to identify testable science questions, support trade studies, help define science cases for future missions, and guide further theoretical and experimental work on the origins of life. Our work highlights the importance of understanding the context in which a biosignature detection is made, which can not only help to assess the credibility of the detection but also to test competing hypotheses on the origins of life on Earth and beyond.

ACKNOWLEDGMENTS

The authors thank Kevin Heng, Dominik Hintz, and Rhys Seeburger for insightful discussions. This material is based upon work supported by the National Aeronautics and Space Administration under Agreement No. 80NSSC21K0593 for the program “Alien Earths”. The results reported herein benefited from collaborations and/or information exchange within NASA’s Nexus for Exoplanet System Science (NExSS) research coordination network sponsored by NASA’s Science Mission Directorate. This work has made use of data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement. T.L. was supported by the Branco Weiss Foundation, the Netherlands eScience Center (PROTEUS project, NLESC.OEC.2023.017), and the Alfred P. Sloan Foundation (AETHer project, G202114194).

REFERENCES

- Alfeld, P. 1984, *Computer Aided Geometric Design*, 1, 169, doi: [10.1016/0167-8396\(84\)90029-3](https://doi.org/10.1016/0167-8396(84)90029-3)
- Apai, D., Milster, T. D., Kim, D. W., et al. 2019, *AJ*, 158, 83, doi: [10.3847/1538-3881/ab2631](https://doi.org/10.3847/1538-3881/ab2631)
- Apai, D., Milster, T. D., Kim, D., et al. 2022, in *Optical Manufacturing and Testing XIV*, Vol. 12221 (SPIE), 59–71, doi: [10.1117/12.2633184](https://doi.org/10.1117/12.2633184)
- Baraffe, I., Chabrier, G., Allard, F., & Hauschildt, P. H. 1998, *Astronomy and Astrophysics*, v.337, p.403-412 (1998), 337, 403
- Baross, J. A., & Hoffman, S. E. 1985, *Origins Life Evol Biosphere*, 15, 327, doi: [10.1007/BF01808177](https://doi.org/10.1007/BF01808177)
- Bergsten, G. J., Pascucci, I., Mulders, G. D., Fernandes, R. B., & Koskinen, T. T. 2022, *AJ*, 164, 190, doi: [10.3847/1538-3881/ac8fea](https://doi.org/10.3847/1538-3881/ac8fea)

AUTHOR CONTRIBUTIONS

M.S., D.A., and S.R. conceived the project, planned its implementation, and interpreted the results. M.S. developed the planetary evolution component to Bioverse, carried out the hypothesis tests and statistical analyses, and wrote the manuscript. D.A. leads the “Alien Earths” program through which this project is funded, helped to guide the strategy of the project, and provided text contributions. A.A. carried out the semi-analytical computations regarding the correlation of past UV flux and biosignature occurrence. S.R. advised on planetary NUV flux evolution and the cyanosulfidic scenario of the origins of life. R.F. wrote the initial draft of the Introduction and advised on the evolutionary biology aspects of the project. K.H.-U. contributed to the Bioverse software development and simulations. T.L. supported the selection of testable hypotheses and provided text contributions to the initial draft. S.M. advised on the scope of the project and supported the selection of testable hypotheses. All authors provided comments and suggestions on the manuscript.

REPRODUCIBILITY

All code required to reproduce our results, figures, and this article itself is available at <https://github.com/matiscke/originsoflife>.

- Bixel, A., & Apai, D. 2021, *The Astronomical Journal*, 161, 228, doi: [10.3847/1538-3881/abe042](https://doi.org/10.3847/1538-3881/abe042)
- Brasier, M. D., Matthewman, R., McMahon, S., & Wacey, D. 2011, *Astrobiology*, 11, 725, doi: [10.1089/ast.2010.0546](https://doi.org/10.1089/ast.2010.0546)
- Buccino, A. P., Lemarchand, G. A., & Mauas, P. J. D. 2007, *Icarus*, 192, 582, doi: [10.1016/j.icarus.2007.08.012](https://doi.org/10.1016/j.icarus.2007.08.012)
- Carrión-González, Ó., Kammerer, J., Angerhausen, D., et al. 2023, *A&A*, 678, A96, doi: [10.1051/0004-6361/202347027](https://doi.org/10.1051/0004-6361/202347027)
- Catling, D. C., Krissansen-Totton, J., Kiang, N. Y., et al. 2018a, *Astrobiology*, 18, 709, doi: [10.1089/ast.2017.1737](https://doi.org/10.1089/ast.2017.1737)
- Catling, D. C., Krissansen-Totton, J., Kiang, N. Y., et al. 2018b, *Astrobiology*, 18, 709, doi: [10.1089/ast.2017.1737](https://doi.org/10.1089/ast.2017.1737)

- Cesario, L., Lichtenberg, T., Alei, E., et al. 2024, Large Interferometer For Exoplanets (LIFE). XIV. Finding Terrestrial Protoplanets in the Galactic Neighborhood, arXiv, doi: [10.48550/arXiv.2410.13457](https://doi.org/10.48550/arXiv.2410.13457)
- Deamer, D., Damer, B., & Kompanichenko, V. 2019, *Astrobiology*, 19, 1523, doi: [10.1089/ast.2018.1979](https://doi.org/10.1089/ast.2018.1979)
- Deamer, D. W., & Georgiou, C. D. 2015, *Astrobiology*, 15, 1091, doi: [10.1089/ast.2015.1338](https://doi.org/10.1089/ast.2015.1338)
- Ducrot, E., Gillon, M., Delrez, L., et al. 2020, *A&A*, 640, A112, doi: [10.1051/0004-6361/201937392](https://doi.org/10.1051/0004-6361/201937392)
- Fields, B., Gupta, S., & Sandora, M. 2023, *International Journal of Astrobiology*, 22, 583, doi: [10.1017/S1473550423000150](https://doi.org/10.1017/S1473550423000150)
- Fox, S., & Strasdeit, H. 2013, *Astrobiology*, 13, 578, doi: [10.1089/ast.2012.0934](https://doi.org/10.1089/ast.2012.0934)
- Glauser, A. M., Quanz, S. P., Hansen, J., et al. 2024, in *Optical and Infrared Interferometry and Imaging IX*, Vol. 13095 (SPIE), 354–374, doi: [10.1117/12.3019090](https://doi.org/10.1117/12.3019090)
- Glazier, A. L., Howard, W. S., Corbett, H., et al. 2020, *ApJ*, 900, 27, doi: [10.3847/1538-4357/aba4a6](https://doi.org/10.3847/1538-4357/aba4a6)
- Günther, M. N., Zhan, Z., Seager, S., et al. 2020, *AJ*, 159, 60, doi: [10.3847/1538-3881/ab5d3a](https://doi.org/10.3847/1538-3881/ab5d3a)
- Hardegree-Ullman, K. K., Apai, D., Bergsten, G. J., Pascucci, I., & López-Morales, M. 2023, *AJ*, 165, 267, doi: [10.3847/1538-3881/acd1ec](https://doi.org/10.3847/1538-3881/acd1ec)
- Hardegree-Ullman, K. K., Apai, D., Haffert, S. Y., et al. 2024, *Bioverse: GMT and ELT Direct Imaging and High-Resolution Spectroscopy Assessment* $\S_{\text{Surveying Exo-Earth O}_2 \text{ and Testing the Habitable Zone Oxygen Hypothesis}}$, <https://arxiv.org/abs/2405.11423v1>
- Hawker, G. A., & Parry, I. R. 2019, *MNRAS*, 484, 4855, doi: [10.1093/mnras/stz323](https://doi.org/10.1093/mnras/stz323)
- Ingersoll, A. P. 1969, *Journal of Atmospheric Sciences*, 26, 1191, doi: [10.1175/1520-0469\(1969\)026<1191:TRGAHO>2.0.CO;2](https://doi.org/10.1175/1520-0469(1969)026<1191:TRGAHO>2.0.CO;2)
- Jeffreys, H. 1939, *Theory of Probability*
- Kammerer, J., & Quanz, S. P. 2018, *A&A*, 609, A4, doi: [10.1051/0004-6361/201731254](https://doi.org/10.1051/0004-6361/201731254)
- Kasting, J. F. 1991, *Icarus*, 94, 1, doi: [10.1016/0019-1035\(91\)90137-I](https://doi.org/10.1016/0019-1035(91)90137-I)
- Kasting, J. F., Whitmire, D. P., & Reynolds, R. T. 1993, *Icarus*, 101, 108, doi: [10.1006/icar.1993.1010](https://doi.org/10.1006/icar.1993.1010)
- Kite, E. S., & Ford, E. B. 2018, *ApJ*, 864, 75, doi: [10.3847/1538-4357/aad6e0](https://doi.org/10.3847/1538-4357/aad6e0)
- Kopparapu, R. K., Ramirez, R. M., SchottelKotte, J., et al. 2014, *ApJL*, 787, L29, doi: [10.1088/2041-8205/787/2/L29](https://doi.org/10.1088/2041-8205/787/2/L29)
- Kopparapu, R. K., Ramirez, R., Kasting, J. F., et al. 2013, *ApJ*, 765, 131, doi: [10.1088/0004-637X/765/2/131](https://doi.org/10.1088/0004-637X/765/2/131)
- Lichtenberg, T., & Miguel, Y. 2024, *Super-Earths and Earth-like Exoplanets*, arXiv, doi: [10.48550/arXiv.2405.04057](https://doi.org/10.48550/arXiv.2405.04057)
- Malaterre, C., Jeancolas, C., & Nghe, P. 2022, *Astrobiology*, 22, 851, doi: [10.1089/ast.2021.0162](https://doi.org/10.1089/ast.2021.0162)
- Mamajek, E., & Stapelfeldt, K. 2023, 55, 116.07
- Mann, H. B., & Whitney, D. R. 1947, *The Annals of Mathematical Statistics*, 18, 50
- Melosh, H. J. 2011, *Planetary Surface Processes*, Cambridge Planetary Science (Cambridge: Cambridge University Press), doi: [10.1017/CBO9780511977848](https://doi.org/10.1017/CBO9780511977848)
- Mol Lous, M., Helled, R., & Mordasini, C. 2022, *Nat Astron*, 6, 819, doi: [10.1038/s41550-022-01699-8](https://doi.org/10.1038/s41550-022-01699-8)
- Mulkidjanian, A. Y., Bychkov, A. Y., Dibrova, D. V., Galperin, M. Y., & Koonin, E. V. 2012, *Proceedings of the National Academy of Sciences*, 109, E821, doi: [10.1073/pnas.1117774109](https://doi.org/10.1073/pnas.1117774109)
- Nicholls, H., Lichtenberg, T., Bower, D. J., & Pierrehumbert, R. 2024, *Magma Ocean Evolution at Arbitrary Redox State*, arXiv, doi: [10.48550/arXiv.2411.19137](https://doi.org/10.48550/arXiv.2411.19137)
- Nielson, G. M. 1983, *Math. Comp.*, 40, 253, doi: [10.1090/S0025-5718-1983-0679444-7](https://doi.org/10.1090/S0025-5718-1983-0679444-7)
- Noack, L., Höning, D., Rivoldini, A., et al. 2016, *Icarus*, 277, 215, doi: [10.1016/j.icarus.2016.05.009](https://doi.org/10.1016/j.icarus.2016.05.009)
- Patel, B. H., Percivalle, C., Ritson, D. J., Duffy, C. D., & Sutherland, J. D. 2015, *Nature Chem*, 7, 301, doi: [10.1038/nchem.2202](https://doi.org/10.1038/nchem.2202)
- Quanz, S. P., Ottiger, M., Fontanet, E., et al. 2022, *A&A*, 664, A21, doi: [10.1051/0004-6361/202140366](https://doi.org/10.1051/0004-6361/202140366)
- Ramirez, R. M. 2018, *Geosciences*, 8, 280, doi: [10.3390/geosciences8080280](https://doi.org/10.3390/geosciences8080280)
- Ramirez, R. M., & Kaltenegger, L. 2017, *ApJL*, 837, L4, doi: [10.3847/2041-8213/aa60c8](https://doi.org/10.3847/2041-8213/aa60c8)
- Ranjan, S., Nayak, P. K., Pineda, J. S., & Narang, M. 2023, *The Astronomical Journal*, 166, 70, doi: [10.3847/1538-3881/ace32d](https://doi.org/10.3847/1538-3881/ace32d)
- Ranjan, S., & Sasselov, D. D. 2016, *Astrobiology*, 16, 68, doi: [10.1089/ast.2015.1359](https://doi.org/10.1089/ast.2015.1359)
- . 2017, *Astrobiology*, 17, 169, doi: [10.1089/ast.2016.1519](https://doi.org/10.1089/ast.2016.1519)
- Ranjan, S., Wordsworth, R., & Sasselov, D. D. 2017a, *Astrobiology*, 17, 687, doi: [10.1089/ast.2016.1596](https://doi.org/10.1089/ast.2016.1596)
- . 2017b, *ApJ*, 843, 110, doi: [10.3847/1538-4357/aa773e](https://doi.org/10.3847/1538-4357/aa773e)
- Rapf, R. J., & Vaida, V. 2016, *Phys. Chem. Chem. Phys.*, 18, 20067, doi: [10.1039/C6CP00980H](https://doi.org/10.1039/C6CP00980H)
- Revol, A., Bolmont, É., Sastre, M., et al. 2024, *A&A*, 691, L3, doi: [10.1051/0004-6361/202451532](https://doi.org/10.1051/0004-6361/202451532)
- Richey-Yowell, T., Shkolnik, E. L., Schneider, A. C., et al. 2019, *ApJ*, 872, 17, doi: [10.3847/1538-4357/aafa74](https://doi.org/10.3847/1538-4357/aafa74)
- . 2023, *ApJ*, 951, 44, doi: [10.3847/1538-4357/acd2dc](https://doi.org/10.3847/1538-4357/acd2dc)

- 1127 Rimmer, P. B. 2023, in *Conflicting Models for the Origin of*
 1128 *Life* (John Wiley & Sons, Ltd), 407–424,
 1129 doi: [10.1002/9781119555568.ch16](https://doi.org/10.1002/9781119555568.ch16)
- 1130 Rimmer, P. B., Ranjan, S., & Rugheimer, S. 2021a,
 1131 *Elements*, 17, 265, doi: [10.2138/gselements.17.4.265](https://doi.org/10.2138/gselements.17.4.265)
- 1132 Rimmer, P. B., Thompson, S. J., Xu, J., et al. 2021b,
 1133 *Astrobiology*, 21, 1099, doi: [10.1089/ast.2020.2335](https://doi.org/10.1089/ast.2020.2335)
- 1134 Rimmer, P. B., Xu, J., Thompson, S. J., et al. 2018, *Science*
 1135 *Advances*, 4, eaar3302, doi: [10.1126/sciadv.aar3302](https://doi.org/10.1126/sciadv.aar3302)
- 1136 Sassellov, D. D., Grotzinger, J. P., & Sutherland, J. D. 2020,
 1137 *Science Advances*, 6, eaax3419,
 1138 doi: [10.1126/sciadv.aax3419](https://doi.org/10.1126/sciadv.aax3419)
- 1139 Schlecker, M., Apai, D., Lichtenberg, T., et al. 2024,
 1140 *Planet. Sci. J.*, 5, 3, doi: [10.3847/PSJ/acf57f](https://doi.org/10.3847/PSJ/acf57f)
- 1141 Segura, A., Kasting, J. F., Meadows, V., et al. 2005,
 1142 *Astrobiology*, 5, 706, doi: [10.1089/ast.2005.5.706](https://doi.org/10.1089/ast.2005.5.706)
- 1143 Smart, R. L., Sarro, L. M., Rybizki, J., et al. 2021, *A&A*,
 1144 649, A6, doi: [10.1051/0004-6361/202039498](https://doi.org/10.1051/0004-6361/202039498)
- 1145 Spinelli, R., Borsa, F., Ghirlanda, G., Ghisellini, G., &
 1146 Haardt, F. 2023, *The Ultraviolet Habitable Zone of*
 1147 *Exoplanets*, doi: [10.48550/arXiv.2303.16229](https://doi.org/10.48550/arXiv.2303.16229)
- 1148 Tuchow, N., Stark, C., & Mamajek, E. 2024, *HPIC: The*
 1149 *Habitable Worlds Observatory Preliminary Input*
 1150 *Catalog*, arXiv, doi: [10.48550/arXiv.2402.08038](https://doi.org/10.48550/arXiv.2402.08038)
- 1151 Tuchow, N. W., & Wright, J. T. 2023, *ApJ*, 944, 71,
 1152 doi: [10.3847/1538-4357/acb054](https://doi.org/10.3847/1538-4357/acb054)
- 1153 Underwood, D. R., Jones, B. W., & Sleep, P. N. 2003,
 1154 *International Journal of Astrobiology*, 2, 289,
 1155 doi: [10.1017/S1473550404001715](https://doi.org/10.1017/S1473550404001715)
- 1156 Vaughan, S. R., Birkby, J. L., Thatte, N., et al. 2024,
 1157 *MNRAS*, 528, 3509, doi: [10.1093/mnras/stae242](https://doi.org/10.1093/mnras/stae242)
- 1158 Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2020,
 1159 *Nat Methods*, 17, 261, doi: [10.1038/s41592-019-0686-2](https://doi.org/10.1038/s41592-019-0686-2)
- 1160 Walker, S. I., Bains, W., Cronin, L., et al. 2018,
 1161 *Astrobiology*, 18, 779, doi: [10.1089/ast.2017.1738](https://doi.org/10.1089/ast.2017.1738)
- 1162 Wang, J., Mawet, D., Ruane, G., Hu, R., & Benneke, B.
 1163 2017, *AJ*, 153, 183, doi: [10.3847/1538-3881/aa6474](https://doi.org/10.3847/1538-3881/aa6474)
- 1164 West, A. A., Hawley, S. L., Bochanski, J. J., et al. 2008,
 1165 *AJ*, 135, 785, doi: [10.1088/0004-6256/135/3/785](https://doi.org/10.1088/0004-6256/135/3/785)
- 1166 Westall, F., Hickman-Lewis, K., Hinman, N., et al. 2018,
 1167 *Astrobiology*, 18, 259, doi: [10.1089/ast.2017.1680](https://doi.org/10.1089/ast.2017.1680)
- 1168 Zahnle, K. J., & Catling, D. C. 2017, *ApJ*, 843, 122,
 1169 doi: [10.3847/1538-4357/aa7846](https://doi.org/10.3847/1538-4357/aa7846)
- 1170 Zhang, H., Wang, J., & Plummer, M. K. 2024, *AJ*, 167, 37,
 1171 doi: [10.3847/1538-3881/ad109e](https://doi.org/10.3847/1538-3881/ad109e)