

STATISTICS

Hypothesis Testing PART 1

Birol Aslanyürek

Rev: 27.11.2024

Hypothesis Testing

Statistics

Hypothesis Testing

A **hypothesis** is an assertion about a parameter of a population or two populations. There are two types of opposite statistical hypotheses:

1. Null Hypothesis (H_0)
2. Alternative Hypothesis (H_a) or (H_1)

Let θ be a population parameter and θ_0 be an asserted value of the parameter. Then, we can build three types of (H_0, H_a) hypothesis.

1)	$H_0: \theta = \theta_0$	2)	$H_0: \theta = \theta_0 \text{ or } H_0: \theta \leq \theta_0$	3)	$H_0: \theta = \theta_0 \text{ or } H_0: \theta \geq \theta_0$
	$H_a: \theta \neq \theta_0$		$H_a: \theta > \theta_0$		$H_a: \theta < \theta_0$

Two-Tailed Hypothesis

One-Tailed Hypothesis

Hypothesis Testing

Statistics

Hypothesis Testing

H_0 and H_a never intersect. H_0 always contains equality operator ($=, \leq, \geq$), while H_a does not.

Example: It is claimed that the average life expectancy of women in Turkey is more than 72 years.

$H_0: \mu = 72$ or $\mu \leq 72$ ➡ The average life expectancy of women in Turkey is **not** more than 72 years.

$H_a: \mu > 72$ ➡ The average life expectancy of women in Turkey is more than 72 years.

Hypothesis Testing

Statistics

There are four possible situations in the process of hypothesis testing.

Decision	Exact Situation	
	H_0 is True	H_0 is False
Accept H_0	Correct Decision ($1 - \alpha$)	II.Type Error (β)
Reject H_0	I.Type Error (α)	Correct Decision ($1 - \beta$)

α : Significance level. It is the probability of rejecting H_0 when H_0 is true.

$1 - \alpha$: It states the confidence level of a test. It is the probability of accepting H_0 when H_0 is true.

$1 - \beta$: The power of the test. It is the probability of rejecting H_0 when H_0 is false.

Hypothesis Testing

Statistics

The Steps of Hypothesis Testing

The hypothesis testing is performed under the assumption that H_0 is correct. The steps of a hypothesis testing are:

1. Determine hypothesis H_0 and H_a .
2. Specify α .
3. Select the test statistic.
4. Select the decision rule (determine rejection regions).
5. Compute the test statistics by using the sample.
6. Make a decision about (reject or fail to reject) H_0 .



Hypothesis Testing

Statistics

Hypothesis Testing for Population Mean μ :

A) For Normal Distributed Populations with Known Variance

- 1) Consider that $H_0: \mu = \mu_0$ for α .
 $H_a: \mu \neq \mu_0$



$$P\left(\mu_0 - z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \leq \bar{x} \leq \mu_0 + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}\right) = 1 - \alpha \quad P\left(-z_{\alpha/2} \leq Z_0 \leq z_{\alpha/2}\right) = 1 - \alpha$$

The gray-shaded region is the 'fail to reject' region, while the other parts are 'reject' regions.

Hypothesis Testing

Statistics


Hypothesis Testing for Population Mean μ :

A) For Normal Distributed Populations with Known Variance

- 1) Consider that $H_0: \mu = \mu_0$ for α .
 $H_a: \mu \neq \mu_0$

Decision Rule:

If $Z_0 < -z_{\alpha/2}$ or $Z_0 > z_{\alpha/2}$, reject H_0 . Otherwise, we fail to reject H_0 .



$$P(-z_{\alpha/2} \leq Z_0 \leq z_{\alpha/2}) = 1 - \alpha$$

Hypothesis Testing

Statistics


Hypothesis Testing for Population Mean μ :

A) For Normal Distributed Populations with Known Variance


2) Consider that $H_0: \mu = \mu_0$ or $\mu \leq \mu_0$ for α .
 $H_a: \mu > \mu_0$

Test Statistic: $Z_0 = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}}$

If H_0 is true, ...



$$P\left(\bar{x} \leq \mu_0 + z_\alpha \frac{\sigma}{\sqrt{n}}\right) = 1 - \alpha$$



$$P(Z_0 \leq z_\alpha) = 1 - \alpha$$

The gray-shaded region is the 'fail to reject' region, the other part is 'reject' region.

Hypothesis Testing

Statistics


Hypothesis Testing for Population Mean μ :

A) For Normal Distributed Populations with Known Variance

2) Consider that $H_0: \mu = \mu_0$ or $\mu \leq \mu_0$ for α .
 $H_a: \mu > \mu_0$

Test Statistic: $Z_0 = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}}$

If H_0 is true, ...



Decision Rule:

If $Z_0 > z_\alpha$, reject H_0 . Otherwise, we fail to reject H_0 .

$$P(Z_0 \leq z_\alpha) = 1 - \alpha$$

Hypothesis Testing

Statistics


Hypothesis Testing for Population Mean μ :

A) For Normal Distributed Populations with Known Variance


3) Consider that $H_0: \mu = \mu_0$ or $\mu \geq \mu_0$ for α .
 $H_a: \mu < \mu_0$

Test Statistic: $Z_0 = \frac{\bar{x} - \mu_0}{\sigma/\sqrt{n}}$

If H_0 is true, ...



$$P\left(\mu_0 - z_\alpha \frac{\sigma}{\sqrt{n}} \leq \bar{x}\right) = 1 - \alpha$$



$$P(-z_\alpha \leq Z_0) = 1 - \alpha$$

The gray-shaded region is the 'fail to reject' region, the other part is 'reject' region.

Hypothesis Testing

Statistics


Hypothesis Testing for Population Mean μ :

A) For Normal Distributed Populations with Known Variance

3) Consider that $H_0: \mu = \mu_0$ or $\mu \geq \mu_0$ for α .
 $H_a: \mu < \mu_0$

Test Statistic: $Z_0 = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}}$

If H_0 is true, ...



Decision Rule:

If $Z_0 < -z_\alpha$, reject H_0 . Otherwise, we fail to reject H_0 .

$$P(-z_\alpha \leq Z_0) = 1 - \alpha$$

Hypothesis Testing

Statistics

Example 1: It is claimed that the IQ scores of undergraduate students in our country have a normal distribution with a mean of 117 and a variance of 160. The IQ scores of 12 randomly selected students have been measured to research this claim for the mean. The following scores are obtained:

103, 96, 112, 130, 111, 125, 89, 100, 104, 117, 99, 118.

Test whether the population mean is 117 ($\alpha = 0.05$).

Ans: $Z_0 = -2.28$, reject that $\mu = 117$ (H_0)

Hypothesis Testing

Statistics

Hypothesis Testing for Population Mean μ :

We have investigated hypothesis testing of μ for the normally distributed population with known variance. Except for test statistic, the same analysis is valid for the populations with unknown variances and/or the populations with unknown distributions.

for normal distributed population
with unknown variance



Test Statistic:

$$t_0 = \frac{\bar{x} - \mu_0}{S/\sqrt{n}}$$

for unknown distributed population
with known variance ($n \geq 30$)



Test Statistic:

$$z_0 = \frac{\bar{x} - \mu_0}{\sigma/\sqrt{n}}$$

for unknown distributed population
with unknown variance
($n \geq 30$)



Test Statistic:

$$z_0 = \frac{\bar{x} - \mu_0}{S/\sqrt{n}}$$

Hypothesis Testing

Statistics

Test Statistic	Hypothesis testing for the population mean (μ)
$z_0 = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$	For the Normal distribution and known population variance
$z_0 = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$	For the unknown distribution ($n \geq 30$) and known population variance
$z_0 = \frac{\bar{X} - \mu_0}{s/\sqrt{n}}$	For the unknown distribution ($n \geq 30$) and unknown population variance
Hypothesis	Decision Rule
1. $H_0: \mu = \mu_0$ $H_a: \mu \neq \mu_0$	If $z_0 > z_{\alpha/2}$ or $z_0 < -z_{\alpha/2}$, then reject H_0 .
2. $H_0: \mu = \mu_0$ or $\mu \geq \mu_0$ $H_a: \mu < \mu_0$	If $z_0 < -z_{\alpha}$, then reject H_0 .
3. $H_0: \mu = \mu_0$ or $\mu \leq \mu_0$ $H_a: \mu > \mu_0$	If $z_0 > z_{\alpha}$, then reject H_0 .
$t_0 = \frac{\bar{X} - \mu_0}{s/\sqrt{n}}$	For the Normal distribution and unknown population variance
Hypothesis	Decision Rule
1. $H_0: \mu = \mu_0$ $H_a: \mu \neq \mu_0$	If $t_0 > t_{n-1, \alpha/2}$ or $t_0 < -t_{n-1, \alpha/2}$, then reject H_0 .
2. $H_0: \mu = \mu_0$ or $\mu \geq \mu_0$ $H_a: \mu < \mu_0$	If $t_0 < -t_{n-1, \alpha}$, then reject H_0 .
3. $H_0: \mu = \mu_0$ or $\mu \leq \mu_0$ $H_a: \mu > \mu_0$	If $t_0 > t_{n-1, \alpha}$, then reject H_0 .

Hypothesis Testing

Statistics

Example 2: It is claimed that the IQ scores of undergraduate students in our country have a normal distribution with a mean of 117. The IQ scores of 12 randomly selected students have been measured to research this claim for the mean. The following scores are obtained:

103, 96, 112, 130, 111, 125, 89, 100, 104, 117, 99, 118.

Test whether the population mean is 117 ($\alpha = 0.05$).

Ans: $t_0 = -2.35$, reject that $\mu = 117$ (H_0)

Hypothesis Testing


Statistics

Hypothesis Testing for Population Variance σ^2 or Standard Deviation σ (for normally distributed populations)

1) Consider that $H_0: \sigma = \sigma_0$ for α .
 $H_a: \sigma \neq \sigma_0$

Test Statistic: $\chi_0^2 = \frac{(n-1)S^2}{\sigma_0^2}$

If H_0 is true, ...



Decision Rule:

If $\chi_0^2 < \chi_{n-1, 1-\alpha/2}^2$ or $\chi_0^2 > \chi_{n-1, \alpha/2}^2$,
reject H_0 . Otherwise, we fail to reject H_0 .

$$P\left(\chi_{n-1, 1-\alpha/2}^2 \leq \chi_0^2 \leq \chi_{n-1, \alpha/2}^2\right) = 1 - \alpha$$

The shaded-regions are the 'reject' regions, while the other part is the 'fail to reject' region.


Hypothesis Testing

Statistics

Hypothesis Testing for Population Variance σ^2 or Standard Deviation σ (for normally distributed populations)

2) Consider that $H_0: \sigma = \sigma_0$ or $\sigma \leq \sigma_0$ for α . Test Statistic: $\chi_0^2 = \frac{(n-1)S^2}{\sigma_0^2}$
 $H_a: \sigma > \sigma_0$

If H_0 is true, ...



$$P(\chi_0^2 \leq \chi_{n-1, \alpha}^2) = 1 - \alpha$$

Decision Rule:

If $\chi_0^2 > \chi_{n-1, \alpha}^2$, reject H_0 . Otherwise, we fail to reject H_0 .

The shaded-region is the 'reject' region, the other parts are the 'fail to reject' regions.


Hypothesis Testing

Statistics

Hypothesis Testing for Population Variance σ^2 or Standard Deviation σ (for normally distributed populations)

3) Consider that $H_0: \sigma = \sigma_0$ or $\sigma \geq \sigma_0$ for α . Test Statistic: $\chi_0^2 = \frac{(n-1)S^2}{\sigma_0^2}$
 $H_a: \sigma < \sigma_0$

If H_0 is true, ...



$$P(\chi_{n-1, 1-\alpha}^2 \leq \chi_0^2) = 1 - \alpha$$

Decision Rule:

If $\chi_0^2 < \chi_{n-1, 1-\alpha}^2$, reject H_0 .
Otherwise, we fail to reject H_0 .

The shaded-region is the 'reject' region, the other parts are the 'fail to reject' regions.

Hypothesis Testing

Statistics

Example 3: The crankshafts whose diameter are 4 cm are produced in a factory. The administrators of the factory do not want that the variance of diameters is greater than 0.25 cm to protect their production standards. If the variance is greater than the specified value, they will reorganize the production procedure. A sample of size 20 crankshafts is randomly selected. If the diameter of the crankshafts is assumed to be normally distributed and the sample variance is 0.32 cm, is a new production procedure required or not? ($\alpha = 0.05$).

Ans: $\chi_0^2 = 24.32$, fail to reject that $\sigma^2 \leq 0.25$ (H_0)


Hypothesis Testing

Statistics

Hypothesis Testing for Population Proportion p ($n \geq 30$)

1) Consider that $H_0: p = p_0$ for α .
 $H_a: p \neq p_0$

Test Statistic: $z_0 = \frac{\hat{p} - p_0}{\sqrt{p_0 q_0 / n}}$



$$P(-z_{\alpha/2} \leq Z_0 \leq z_{\alpha/2}) = 1 - \alpha$$

Decision Rule:

If $Z_0 < -z_{\alpha/2}$ or $Z_0 > z_{\alpha/2}$, reject H_0 . Otherwise, we fail to reject H_0 .

The gray-shaded region is the 'fail to reject' region, while the other parts are 'reject' regions.


Hypothesis Testing

Statistics

Hypothesis Testing for Population Proportion p ($n \geq 30$)

2) Consider that $H_0: p = p_0$ or $p \leq p_0$ for α . Test Statistic: $z_0 = \frac{\hat{p} - p_0}{\sqrt{p_0 q_0 / n}}$
 $H_a: p > p_0$

If H_0 is true, ...



$$P(Z_0 \leq z_\alpha) = 1 - \alpha$$

Decision Rule:

If $Z_0 > z_\alpha$, reject H_0 . Otherwise, we fail to reject H_0 .

The gray-shaded region is the 'fail to reject' region, the other part is 'reject' region.


Hypothesis Testing

Statistics

Hypothesis Testing for Population Proportion p ($n \geq 30$)

3) Consider that $H_0: p = p_0$ or $p \geq p_0$ for α . Test Statistic: $z_0 = \frac{\hat{p} - p_0}{\sqrt{p_0 q_0 / n}}$
 $H_a: p < p_0$

If H_0 is true, ...



$$P(-z_\alpha \leq Z_0) = 1 - \alpha$$

Decision Rule:

If $Z_0 < -z_\alpha$, reject H_0 . Otherwise, we fail to reject H_0 .

The gray-shaded region is the 'fail to reject' region, the other part is 'reject' region.

Hypothesis Testing

Statistics

Example 4: A drug company claimed that a newly produced drug has a side effect on at most 10% of the patients. The drug is tested on 300 patients, and it is observed that the side effect appeared in 45 patients. Is it very significant proof to reject the company's claim?

Remark: If a hypothesis is claimed for $\alpha = 0.01$, we say that 'it is very significant proof'. If α is selected as 0.05, we say that 'it is significant proof'.

Ans: $Z_0 = 2.89$, reject that $p \leq 0.1$ (H_0). It's very significant proof to reject the claim.

Hypothesis Testing

Statistics

Hypothesis Testing for the Difference of Population Means ($\mu_1 - \mu_2$)

A) For Normal Distributed Populations with Known Variance

To compare mean of the two populations, we can use the following hypothesis:

$$\begin{array}{ll} H_0: \mu_1 = \mu_2 & \text{or} \\ H_a: \mu_1 \neq \mu_2 & H_0: \mu_1 - \mu_2 = 0 \\ & H_a: \mu_1 - \mu_2 \neq 0 \end{array}$$

For more general analysis, we can use

$$\begin{array}{l} H_0: \mu_1 - \mu_2 = \delta_0 \\ H_a: \mu_1 - \mu_2 \neq \delta_0 \end{array}$$

Hypothesis Testing


Statistics

Hypothesis Testing for the Difference of Population Means ($\mu_1 - \mu_2$)

1) Consider that $H_0: \mu_1 - \mu_2 = \delta_0$ for α .
 $H_a: \mu_1 - \mu_2 \neq \delta_0$

Test Statistic:
$$Z_0 = \frac{(\bar{x}_1 - \bar{x}_2) - \delta_0}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$$

If H_0 is true, ...



$$P(-z_{\alpha/2} \leq Z_0 \leq z_{\alpha/2}) = 1 - \alpha$$

Decision Rule:

If $Z_0 < -z_{\alpha/2}$ or $Z_0 > z_{\alpha/2}$, **reject** H_0 . Otherwise, we **fail to reject** H_0 .

The gray-shaded region is the 'fail to reject' region, while the other parts are 'reject' regions.

Hypothesis Testing

Statistics

As seen, a hypothesis test can be performed like the mean of a single population. Thus, we can derive the following test statistics and decision rules similarly.

Test Statistic	Hypothesis testing for the difference of population mean ($\mu_1 - \mu_2$)
$z_0 = \frac{(\bar{X}_1 - \bar{X}_2) - \delta_0}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$	For the Normal distribution and known populations variances
$z_0 = \frac{(\bar{X}_1 - \bar{X}_2) - \delta_0}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$	For the unknown distribution ($n_1 \geq 30$ and $n_2 \geq 30$) and known populations variances
$z_0 = \frac{(\bar{X}_1 - \bar{X}_2) - \delta_0}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$	For the unknown distribution ($n_1 \geq 30$ and $n_2 \geq 30$) and unknown populations variances
Hypothesis	Decision Rule
1. $H_0: \mu_1 - \mu_2 = \delta_0$ $H_a: \mu_1 - \mu_2 \neq \delta_0$	If $z_0 > z_{\alpha/2}$ or $z_0 < -z_{\alpha/2}$, then reject H_0 .
2. $H_0: \mu_1 - \mu_2 = \delta_0$ or $\mu_1 - \mu_2 \geq \delta_0$ $H_a: \mu_1 - \mu_2 < \delta_0$	If $z_0 < -z_{\alpha}$, then reject H_0 .
3. $H_0: \mu_1 - \mu_2 = \delta_0$ or $\mu_1 - \mu_2 \leq \delta_0$ $H_a: \mu_1 - \mu_2 > \delta_0$	If $z_0 > z_{\alpha}$, then reject H_0 .

Hypothesis Testing

Statistics

$t_0 = \frac{(\bar{X}_1 - \bar{X}_2) - \delta_0}{S_k \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}$ $v = n_1 + n_2 - 2$	For the Normal distribution and unknown populations variances ($\sigma_1^2 = \sigma_2^2$)
$t_0 = \frac{(\bar{X}_1 - \bar{X}_2) - \delta_0}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$ $v = \frac{\left(\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}\right)^2}{\frac{(s_1^2/n_1)^2}{n_1 - 1} + \frac{(s_2^2/n_2)^2}{n_2 - 1}}$	For the Normal distribution and unknown populations variances ($\sigma_1^2 \neq \sigma_2^2$)
Hypothesis	Decision Rule
1. $H_0: \mu_1 - \mu_2 = \delta_0$ $H_a: \mu_1 - \mu_2 \neq \delta_0$	If $t_0 > t_{v, \alpha/2}$ or $t_0 < -t_{v, \alpha/2}$, then reject H_0 .
2. $H_0: \mu_1 - \mu_2 = \delta_0$ or $\mu_1 - \mu_2 \geq \delta_0$ $H_a: \mu_1 - \mu_2 < \delta_0$	If $t_0 < -t_{v, \alpha}$, then reject H_0 .
3. $H_0: \mu_1 - \mu_2 = \delta_0$ or $\mu_1 - \mu_2 \leq \delta_0$ $H_a: \mu_1 - \mu_2 > \delta_0$	If $t_0 > t_{v, \alpha}$, then reject H_0 .

Hypothesis Testing

Statistics

Example 5: A research is made to investigate the nicotine content of two cigarette brands. A sample of 50 cigarettes in the first type has a mean of 2.61 mg and a standard deviation of 0.12 mg as well as a sample of 40 cigarettes in the second type has a mean of 2.38 mg and a standard deviation of 0.14 mg. Compare the hypothesis $\mu_1 - \mu_2 = 0.20$ to the alternative hypothesis $\mu_1 - \mu_2 \neq 0.20$ in the level of significance 0.05.

Ans: $Z_0 = -6.94$, reject that $\mu_1 - \mu_2 = 0.20$ (H_0)

Hypothesis Testing

Statistics

Example 6: To compare two diet programs, a group of 9 people applied for the first diet program, and another group of 12 people applied for the second program during a month. It is observed that the people in the first group and the second group have lost an average of 7.2 kg and 6.9 kg, respectively. Their standard deviations are measured as 0.59 kg and 0.82 kg, respectively. By assuming that the population distributions are normally distributed, for $\alpha=0.05$, test the hypothesis “there is no significant difference between diet programs” with the alternative hypothesis “the first diet program is more efficient”.

Ans: $t_0 = 0.97$, fail to reject that $\mu_1 - \mu_2 = 0$ (H_0), there is no significant proof that there is no difference between diet programs

Hypothesis Testing

Statistics



Hypothesis Testing for the Difference of Population Proportions ($p_1 - p_2$) ($n_1 \geq 30, n_2 \geq 30$)

To compare the proportion of the two populations, we can use the following hypothesis:

$$H_0: p_1 - p_2 = \delta_0$$

$$H_a: p_1 - p_2 \neq \delta_0$$

If $n_1 \geq 30$ and $n_2 \geq 30$, we know that the following random variable has approximately standard normal distribution:

$$Z = \frac{(\hat{p}_1 - \hat{p}_2) - (p_1 - p_2)}{\sqrt{\frac{\hat{p}_1 \hat{q}_1}{n_1} + \frac{\hat{p}_2 \hat{q}_2}{n_2}}}$$

If we use weighted means $\hat{p}_w$ and $\hat{q}_w$ instead of $\hat{p}_1, \hat{q}_1, \hat{p}_2, \hat{q}_2$, the following test statistic is used:

$$Z_0 = \frac{(\hat{p}_1 - \hat{p}_2) - \delta_0}{\sqrt{\hat{p}_w \hat{q}_w \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}}, \quad \hat{p}_w = \frac{n_1 \hat{p}_1 + n_2 \hat{p}_2}{n_1 + n_2}, \quad \hat{q}_w = 1 - \hat{p}_w$$

Hypothesis Testing

Statistics

We can derive the following test statistics and decision rules similarly.

Test Statistic	Hypothesis testing for the difference of the population proportion ($p_1 - p_2$)
$z_0 = \frac{(\hat{p}_1 - \hat{p}_2) - p_0}{\sqrt{\hat{p}_w \hat{q}_w \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}}$	For ($n_1 \geq 30$ and $n_2 \geq 30$)
Hypothesis	Decision Rule
1. $H_0: p_1 - p_2 = \delta_0$ $H_a: p_1 - p_2 \neq \delta_0$	If $z_0 > z_{\alpha/2}$ or $z_0 < -z_{\alpha/2}$, then reject H_0 .
2. $H_0: p_1 - p_2 = \delta_0$ or $p_1 - p_2 \geq \delta_0$ $H_a: p_1 - p_2 < \delta_0$	If $z_0 < -z_{\alpha}$, then reject H_0 .
3. $H_0: p_1 - p_2 = \delta_0$ or $p_1 - p_2 \leq \delta_0$ $H_a: p_1 - p_2 > \delta_0$	If $z_0 > z_{\alpha}$, then reject H_0 .

Hypothesis Testing

Statistics

Example 7: It is seen that 21 families have a villa in 220 families which are randomly selected in Istanbul, and 18 families have a villa in 177 families which are randomly selected in Ankara. Is there a significant difference between the proportions for the two cities regarding having a villa?

Ans: $z_0 = -0.23$, fail to reject that $p_1 - p_2 = 0$ (H_0), there is no significant proof that there is no difference between the proportions for the two cities regarding having a villa.

Hypothesis Testing

Statistics

Hypothesis Testing for the Rate of Population Variances (σ_1^2 / σ_2^2)

Assume that both populations have a normal distribution. We can use the following hypothesis:

$$\begin{array}{ll} H_0: \sigma_1^2 = \sigma_2^2 & \text{or} & H_0: \sigma_1^2 / \sigma_2^2 = 1 \\ H_a: \sigma_1^2 \neq \sigma_2^2 & & H_a: \sigma_1^2 / \sigma_2^2 \neq 1 \end{array}$$

We know that

$$F_{n_1-1, n_2-1} = \frac{S_1^2 \sigma_2^2}{S_2^2 \sigma_1^2}$$

When H_0 is true ($\sigma_1^2 = \sigma_2^2$), the test statistic becomes:

$$F_0 = \frac{S_1^2}{S_2^2}$$


Hypothesis Testing

Statistics

Hypothesis Testing for the Rate of Population Variances (σ_1^2/σ_2^2) (for normal distributed populations)

Consider that $H_0: \sigma_1^2/\sigma_2^2 = 1$ for α .
If H_0 is true, ... $H_a: \sigma_1^2/\sigma_2^2 \neq 1$

Test Statistic: $F_0 = \frac{S_1^2}{S_2^2}$



Decision Rule:

If $F_0 < F_{n_1-1, n_2-1, 1-\alpha/2}$ or $F_0 > F_{n_1-1, n_2-1, \alpha/2}$, **reject** H_0 .
Otherwise, we **fail to reject** H_0 .

$$P\left(F_{n_1-1, n_2-1, 1-\alpha/2} \leq F_0 \leq F_{n_1-1, n_2-1, \alpha/2}\right) = 1 - \alpha$$

The shaded-regions are the '**reject**' regions, while the other part is the '**fail to reject**' region.

Hypothesis Testing

Statistics

We can derive the following test statistics and decision rules similarly.

Test Statistic	Hypothesis testing for the rate of populations variances (σ_1^2/σ_2^2)
$F_0 = \frac{S_1^2}{S_2^2}$	For the Normal distribution
Hypothesis	Decision Rule
1. $H_0: \sigma_1^2/\sigma_2^2 = 1$ $H_a: \sigma_1^2/\sigma_2^2 \neq 1$	If $F_0 > F_{n_1-1, n_2-1, \alpha/2}$ or $F_0 < F_{n_1-1, n_2-1, 1-\alpha/2}$, then reject H_0 .
2. $H_0: \sigma_1^2/\sigma_2^2 = 1$ or $\sigma_1^2/\sigma_2^2 \geq 1$ $H_a: \sigma_1^2/\sigma_2^2 < 1$	If $F_0 < F_{n_1-1, n_2-1, 1-\alpha}$, then reject H_0 .
3. $H_0: \sigma_1^2/\sigma_2^2 = 1$ or $\sigma_1^2/\sigma_2^2 \leq 1$ $H_a: \sigma_1^2/\sigma_2^2 > 1$	If $F_0 > F_{n_1-1, n_2-1, \alpha}$, then reject H_0 .

Hypothesis Testing

Statistics

Example 8: In a soap factory, two machines fill the boxes whose capacities are 300 gr with soap powder. The randomly selected two samples of 10 boxes that are filled by these machines are weighted. If their standard deviations are 19 gr and 12 gr, respectively, test the hypothesis “there is no significant difference between two machines regarding the variance” with the alternative hypothesis “the first machine has more variance”. ($\alpha = 0.05$)

Ans: $F_0 = 2.51$, fail to reject that $\sigma_1^2 = \sigma_2^2 = 0$ (H_0), there is no significant proof that there is no difference between two machines regarding the variance.

Hypothesis Testing

Statistics


Hypothesis Testing for the Means of Paired Populations (μ_D)

A) For Normal Distributed Populations

1) Consider that $H_0: \mu_D = \delta_0$ for α .
 $H_a: \mu_D \neq \delta_0$

Test Statistic: $t_0 = \frac{\bar{D} - \delta_0}{\frac{S_D}{\sqrt{n}}}$

If H_0 is true, ...



Decision Rule:

If $t_0 < -t_{\alpha/2, n-1}$ or $t_0 > t_{\alpha/2, n-1}$,
reject H_0 . Otherwise, we fail to reject H_0 .

$$P(-t_{\alpha/2, n-1} \leq t_0 \leq t_{\alpha/2, n-1}) = 1 - \alpha$$

The gray-shaded region is the 'fail to reject' region, while the other parts are 'reject' regions.

Hypothesis Testing

Statistics


Hypothesis Testing for the Means of Paired Populations (μ_D)

B) For Any Populations, $n \geq 30$

1) Consider that $H_0: \mu_D = \delta_0$ for α .
 $H_a: \mu_D \neq \delta_0$

Test Statistic: $z_0 = \frac{\bar{D} - \delta_0}{\frac{s_D}{\sqrt{n}}}$

If H_0 is true, ...



$$P(-z_{\alpha/2} \leq z_0 \leq z_{\alpha/2}) = 1 - \alpha$$

Decision Rule:

If $z_0 < -z_{\alpha/2}$ or $z_0 > z_{\alpha/2}$, reject H_0 . Otherwise, we fail to reject H_0 .

The gray-shaded region is the 'fail to reject' region, while the other parts are 'reject' regions.

Hypothesis Testing

Statistics

We can derive the following test statistics and decision rules similarly.

Test Statistic	Hypothesis testing for the means of paired populations (μ_D)
$t_0 = \frac{\bar{D} - \delta_0}{S_D / \sqrt{n}}$	For a normal distribution
Hypothesis	Decision Rule
1. $H_0: \mu_D = \delta_0$ $H_a: \mu_D \neq \delta_0$	If $t_0 > t_{\alpha/2, n-1}$ or $t_0 < -t_{\alpha/2, n-1}$, then reject H_0
2. $H_0: \mu_D = \delta_0$ or $\mu_D \geq \delta_0$ $H_a: \mu_D < \delta_0$	If $t_0 < -t_{\alpha, n-1}$, then reject H_0
3. $H_0: \mu_D = \delta_0$ or $\mu_D \leq \delta_0$ $H_a: \mu_D > \delta_0$	If $t_0 > t_{\alpha, n-1}$, then reject H_0
$Z_0 = \frac{\bar{D} - \delta_0}{S_D / \sqrt{n}}$	For any distribution ($n \geq 30$)
Hypothesis	Decision Rule
1. $H_0: \mu_D = \delta_0$ $H_a: \mu_D \neq \delta_0$	If $Z_0 > Z_{\alpha/2}$ or $Z_0 < -Z_{\alpha/2}$, then reject H_0
2. $H_0: \mu_D = \delta_0$ or $\mu_D \geq \delta_0$ $H_a: \mu_D < \delta_0$	If $Z_0 < -Z_{\alpha}$, then reject H_0
3. $H_0: \mu_D = \delta_0$ or $\mu_D \leq \delta_0$ $H_a: \mu_D > \delta_0$	If $Z_0 > Z_{\alpha}$, then reject H_0

Hypothesis Testing

Statistics

Example 9: To test the effect of an online tutorial on the students, 10 students are randomly chosen and their grades are measured before and after the online tutorial.

Students	After the online tutorial	Before the online tutorial
Student_1	62	61
Student_2	71	76
Student_3	66	70
Student_4	87	91
Student_5	69	46
Student_6	77	80
Student_7	45	37
Student_8	61	73
Student_9	89	90
Student_10	65	70

Test the hypothesis “the online tutorial increases the student grades” with significance level $\alpha = 0.05$. (Assume that populations have normal distribution.)

Hypothesis Testing

Statistics

Students	X_{1i}	X_{2i}	D_i	$D_i - \bar{D}$	$(D_i - \bar{D})^2$
Student_1	62	61	1	1.2	1.44
Student_2	71	76	-5	-4.8	23.04
Student_3	66	70	-4	-3.8	14.44
Student_4	87	91	-4	-3.8	14.44
Student_5	69	46	23	23.2	538.24
Student_6	77	80	-3	-2.8	7.84
Student_7	45	37	8	8.2	67.24
Student_8	61	73	-12	-11.8	139.24
Student_9	89	90	-1	-0.8	0.64
Student_10	65	70	-5	-4.8	23.04

SUM:

-2

829.6

$$H_0: \mu_D \leq 0$$

$$H_a: \mu_D > 0$$

$$\bar{D} = \frac{\sum_{i=1}^n D_i}{n} = -0.2$$

$$S_D = \sqrt{\frac{\sum_{i=1}^n (D_i - \bar{D})^2}{n-1}} \cong 9.60$$

$$t_0 = \frac{\bar{D} - \delta_0}{S_D / \sqrt{n}} = \frac{-0.2 - 0}{9.60 / \sqrt{10}} \cong -0.066$$

$$t_{n-1, \alpha} = t_{10-1, 0.05} = 1.833$$

Since $t_0 \leq t_{n-1, \alpha}$, we **fail to reject** H_0 .
We have no significant proof to reject that the online tutorials have not increased the grades.

Hypothesis Testing

Statistics

p-value


Consider a two-tailed hypothesis testing. *p-value* is a probability which can be determined from the following expression:

$$P(-z_0 \leq z \leq z_0) = 1 - p_value$$

where z_0 is the computed test statistic. We can also make a decision about hypothesis testing by using the *p-value*:

If $p_{value} < \alpha$ $\Rightarrow$ reject H_0

If $p_{value} \geq \alpha$ $\Rightarrow$ fail to reject H_0



Hypothesis Testing

Statistics

***p*-value**

For one-tailed hypothesis testing, the *p*-value can be determined from the following expressions:



$$P(z \leq z_0) = 1 - p_value$$

$$P(-Z_0 \leq z) = 1 - p_value$$

where z_0 is the computed test statistic. We can also make a decision about hypothesis testing by using the *p*-value:

If $p_value < \alpha \Rightarrow$ Reject H_0

If $p_value \geq \alpha \Rightarrow$ fail to reject H_0



Hypothesis Testing

Statistics

Similarly, p -value can be calculated for t , χ^2 or F distributions.

Example 10: Find p values for Ex1, Ex2 and Ex 4.

Ans: Ex1) p -value = 0.0226, Ex2) p -value = 0.0414, Ex4) p -value = 0.0019

Example 11: Solve other examples through the computation of p values.