

SPECTRA: Stellar Parameter Estimation and Calculation Tools for Research and Analysis

João Paulo Almeida da Silva Matos ^{1,2¶}, Maria Jaqueline Vasconcelos ², and Adriano Hoth Cerqueira ²

¹ Department of Physics, São Paulo State University (UNESP), Institute of Biosciences, Humanities and Exact Sciences, São José do Rio Preto, SP, 15054-000, Brazil ² Laboratório de Astrofísica Teórica e Observacional, Universidade Estadual de Santa Cruz (UESC), Ilhéus, Brasil ¶ Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

Determining fundamental stellar parameters — such as mass, age, luminosity, effective temperature (T_{eff}), radius, and distance — is central to observational and theoretical astrophysics ([Serenelli et al., 2021](#)). **SPECTRA**¹ (*Stellar Parameter Estimation and Calculation Tools for Research and Analysis*, v1.0.0_build_160726) is an open-source Python tool designed to optimize and automate stellar mass determination and diagnostic modeling for stars in clusters, with a particular focus on young, low-mass stars.

The software integrates data preprocessing, missing data imputation via k -Nearest Neighbors (k -NN) and iterative techniques, theoretical isochrone fitting (IsocFit), machine-learning regression modeling (Mass-Magnitude Modeling), and interactive visualization tools ([Baraffe et al., 2015](#); [Serenelli et al., 2021](#)).

Statement of need

In Astrophysics, estimating the mass of single stars outside binary systems relies heavily on theoretical stellar evolution models and mass-luminosity or mass-magnitude relations (MLR/MMR) ([Benedict et al., 2016](#); [Serenelli et al., 2021](#)). Taking advantage of age and metallicity homogeneity of stars belonging to clusters, isochrone or stellar track fitting on color magnitude diagrams proves to be a useful tool to obtain simultaneously stellar ages and mass. However, it depends on a well defined main sequence which in turn depends on high quality data and on the age of the cluster. Younger clusters have a great number of pre main sequence (PMS) stars that lie above the main sequence making the fitting more difficult. Moreover, PMS stars exhibit high observational uncertainties due to differential interstellar extinction, age dispersion, and incomplete data vectors ([Herczeg & Hillenbrand, 2015](#)). Data quality and constraints, differences between stellar models affect the final results. Stellar structure and evolutionary models differ in the adopted input physics and this impact the isochrone fitting results ([Lebreton et al., 2014](#)). On the other hand, MLR or MMR are model independent and rely on gathering the best available photometric and spectroscopic data for systems that can provide direct mass values as, for example, double-lined eclipsing binary systems. Using MLR/MMR requires goodness-of-fit methods that can be statistical in nature or that can rely on the use of machine learning techniques. SPECTRA uses both isochrone fitting and MMR relations to calculate mass values of stars belonging to stellar clusters.

SPECTRA addresses these challenges by offering a unified, user-friendly desktop application built

¹The development of this software and its underlying research framework were initiated and primarily executed at the Universidade Estadual de Santa Cruz (UESC), with final updates and infrastructure migration supported by the Universidade Estadual Paulista (UNESP).

39 with `ttkbootstrap`. It enables researchers and educators to: 1. Automatically preprocess,
40 filter, and impute incomplete photometric datasets. 2. Execute automated local interpolation
41 over theoretical evolutionary grids (e.g., Siess et al. 2000; BHAC15) via `IsocFit` (Baraffe
42 et al., 2015). > [como você vai tratar estas citações e as referências?] 3. Train, optimize
43 (via Grid Search and K -fold cross-validation), and statistically evaluate machine learning
44 regression algorithms (e.g., Random Forests, Support Vector Regressors, k -NN) to construct
45 semi-empirical mass-magnitude relations. 4. Obtain stellar parameters including mass, age,
46 effective temperature and luminosity.

47 State of the field

48 Several computational tools exist within the astronomical ecosystem for stellar parameter
49 estimation and cluster analysis: * Machine learning tools applied to stellar parameter derivation,
50 such as CARMENES transfer-learning routines (Bello-García et al., 2023), target specific
51 spectral types (M dwarfs) rather than open clusters. * Empirical relations based on low-mass
52 field binaries (Benedict et al., 2016; Mann et al., 2019) provide benchmarks for main-sequence
53 objects, but lack integrated GUI workflows for cluster-wide missing-data imputation and
54 pre-main-sequence fits.

55 SPECTRA fills this gap by combining GUI-driven visual diagnostics (e.g., Hertzsprung-Russell and
56 diagnostic residual plots), automated missing-data imputation, and a model-ranking framework
57 based on goodness-of-fit metrics ($RMSE$, MAE , R^2 , and AIC).

58 Software design and workflow

59 SPECTRA is written in Python 3 and relies on core packages including `numpy`, `pandas`, `scikit-`
60 `learn`, `statsmodels`, and `ttkbootstrap` for user interface rendering (Gomes, 2024). The
61 application's architecture decouples GUI control logic from heavy computational backends,
62 providing a modular design structured around three main analytical pathways.

63 Figure 1 illustrates the high-level execution flow and operational logic of SPECTRA:




Figure 1: Execution workflow and modular architecture of SPECTRA.

Execution Pipeline & Module Description

1. **Initialization & Environment Setup:** Upon launching `main.py`, SPECTRA performs an automated first-run verification check (Gomes, 2024). If local theoretical evolutionary grids (e.g., Siess et al. 2000; BHAC15) or model weights are missing, the system automatically fetches and caches them to ensure reproducible offline execution (Baraffe et al., 2015; Gomes, 2024).

2. **Core Analytical Workflows:** From the primary desktop window, users can direct multi-catalog photometry (e.g., *Gaia* DR3, TESS) into three specialized processing modules (Babusiaux et al., 2023; Gomes, 2024; Paegert et al., 2021; Stassun et al., 2019):

- **Isochrone Fitting Module (IsocFit):** Implements local grid-search minimization to map observed stars onto theoretical evolutionary tracks (Gomes, 2024). The Euclidean distance δ in the HR diagram space is minimized via:

$$\delta = \sqrt{(T_{\text{eff}} - T_{\text{eff},0})^2 + (L - L_0)^2}$$

where $(T_{\text{eff},0}, L_0)$ are interpolated target parameters, followed by refined local track interpolation across adjacent stellar mass and age boundaries (Gomes, 2024).

- **Mass-Magnitude Modeling Module (RMM):** Builds semi-empirical relations by training candidate regression algorithms (e.g., k -NN, Random Forests, Support Vector Regressors, Bayesian Linear Regression) over theoretical stellar grids (Gomes, 2024). Hyperparameters are optimized via K -fold cross-validation and Grid Search (Gomes, 2024). Candidate models are evaluated and automatically ranked using a normalized multi-metric score S :

$$S = \hat{\mathcal{M}} \cdot \mathcal{P}^T$$

where $\hat{\mathcal{M}}$ represents the min-max normalized metrics matrix and $\mathcal{P}$ corresponds to an unit vector, assigning equal relative importance to $RMSE$, MAE , $(1 - R^2)$, and AIC to compute the composite metric score via matrix multiplication (Gomes, 2024).

- **Mathematical Modeling Module:** Provides advanced preprocessing, dataset filtering, missing attribute imputation (e.g., k -NN and iterative stochastic techniques), feature correlation analysis, Principal Component Analysis (PCA), and residual diagnostic checks (homoscedasticity and normality of errors) (Gomes, 2024).

3. **Output Artifact Generation:** All generated diagnostics—including annotated HR/Color-Magnitude diagrams, statistical summary tables (.csv), regression report logs, and diagnostic residual plots—are automatically organized and saved in the localized outputs/repository for direct export into publications (Gomes, 2024).

Research impact & empirical case study

The practical effectiveness of SPECTRA was validated through a comprehensive study of young stellar clusters covering distinct evolutionary stages: Orion Nebula Cluster (ONC, ~ 1 Myr) (Hillenbrand, 1997), Upper Scorpius (~ 5 – 11 Myr) (Pécaut et al., 2012; Rebull et al., 2018), h Persei / NGC 869 (~ 13 Myr), Pleiades (~ 112 Myr) (Dahm, 2015; Gossage et al., 2018; Lodieu et al., 2019), and NGC 2516 (~ 150 Myr).

Methodological Validation

Using a control sample of isolated eclipsing binary stars from Benedict et al. (2016), SPECTRA's k -NN regression model derived from the BHAC15 isochrone grid (Baraffe et al., 2015) achieved high predictive accuracy ($R^2 = 0.96$, $P_{\text{skedasticity}} = 0.12$) with a median residual uncertainty of 3.8%.

Cluster Analysis Results

- **Young PMS Clusters (ONC & Upper Scorpius):** For clusters rich in pre-main-sequence objects with significant age dispersion (Hillenbrand, 1997; Pécaut et al., 2012), IsocFit successfully identified low-mass PMS objects, whereas k -NN regression provided robust mass estimates with less dependency on bolometric corrections (Pécaut & Mamajek, 2013).
- **Well-Defined Main-Sequence Clusters (Pleiades & NGC 2516):** On clusters with established main sequences (Lodieu et al., 2019), k -NN RMM modeling using *Gaia* G -band and V -band absolute magnitudes yielded near-perfect alignment with literature values ($R^2 > 0.99$, $RMSE < 0.01M_{\odot}$), demonstrating superior performance over traditional isochrone grid edge-effects.

Figure 2 illustrates a comparative evaluation of mass estimates derived by SPECTRA for member stars of the Pleiades cluster using both the traditional grid-interpolation method (IsocFit - BHAC15, left panels) and the machine-learning Mass-Magnitude relation (KNN - BHAC15, right panels), benchmarked against literature values from TESS V8.2 (Stassun et al., 2019), Lodieu et al. (2019) (Lodieu et al., 2019), and Delfosse et al. (2000) (Benedict et al., 2016).




Figure 2: Comparison of stellar mass estimates for the Pleiades cluster obtained via IsocFit (left column) and KNN - BHAC15 (right column) against reference datasets from TESS V8.2, Lodieu et al. (2019), and Delfosse et al. (2000). The dashed line represents the ideal 1:1 identity relation.

As shown in Figure 2, the IsocFit - BHAC15 method exhibits greater dispersion around the 1:1 identity line, particularly for stars with $M > 0.7M_{\odot}$. This behavior is attributed to the convergence of evolutionary tracks near the Zero-Age Main Sequence (ZAMS) at the age of the Pleiades (~ 112 Myr), which creates localized degeneracies during grid minimization[cite: 1]. Conversely, the semi-empirical KNN - BHAC15 model demonstrates exceptional predictive stability, showing near-perfect alignment ($R^2 \approx 1.0$) when cross-validated against the photometrically derived masses of Lodieu et al. (2019). When benchmarked against empirical

binary masses from Benedict et al. (2016), the k -NN model reproduces stellar masses with minimal bias up to the applicability threshold of the empirical K -band relation ($M \lesssim 0.75M_{\odot}$) [cite: 1].

A more detailed astrophysical discussion and complete parameter catalogues derived for each cluster will be presented in a forthcoming paper (Gomes et al., in prep.).

By reducing manual preprocessing time and providing rigorous statistical diagnostic reports, SPECTRA supports both observational research in stellar rotation/dynamics and undergraduate/graduate instruction in stellar astrophysics (Gomes, 2024).

AI usage disclosure

No generative AI tools were used in the algorithmic development or code implementation of the SPECTRA package.

Acknowledgements

The primary development, algorithmic design, and empirical cluster validations of SPECTRA were conducted at the Laboratório de Astrofísica Teórica e Observacional (LATO) at the Universidade Estadual de Santa Cruz (UESC), whose financial and structural support during the initial phases of this research was fundamental. The primary author also acknowledges the current institutional support provided by the Universidade Estadual Paulista (UNESP) during the final stages and preparation of this software manuscript, as well as the entire open-source Python scientific community. The work of MJV and AHC is partially funded by the UESC's projects 2023.0003119-74 and 2025.0033125-91.

References

- Babusiaux, C., Fabricius, C., Khanna, S., Muraveva, T., Reylé, C., Spoto, F., Vallenari, A., Luri, X., Arenou, F., Álvarez, M. A., Anders, F., Antoja, T., Balbinot, E., Barache, C., Bauchet, N., Bossini, D., Busonero, D., Cantat-Gaudin, T., Carrasco, J. M., ... Weiler, M. (2023). *Gaia* Data Release 3: Catalogue validation. *Astronomy & Astrophysics*, 674, A32. <https://doi.org/10.1051/0004-6361/202243790>
- Baraffe, I., Homeier, D., Allard, F., & Chabrier, G. (2015). New evolutionary models for pre-main sequence and main sequence low-mass stars down to the hydrogen-burning limit. *Astronomy & Astrophysics*, 577, A42. <https://doi.org/10.1051/0004-6361/201425481>
- Bello-García, A., Passegger, V. M., Ordieres-Meré, J., Schweitzer, A., Caballero, J. A., González-Marcos, A., Ribas, I., Reiners, A., Quirrenbach, A., Amado, P. J., Béjar, V. J. S., Cifuentes, C., Henning, Th., Kaminski, A., Luque, R., Montes, D., Morales, J. C., Pedraz, S., Tabernero, H. M., & Zechmeister, M. (2023). The CARMENES search for exoplanets around M dwarfs: A deep transfer learning method to determine t_{eff} and $[M/H]$ of target stars. *Astronomy & Astrophysics*, 673, A105. <https://doi.org/10.1051/0004-6361/202243934>
- Benedict, G. F., Henry, T. J., Franz, O. G., McArthur, B. E., Wasserman, L. H., Jao, W.-C., Cargile, P. A., Dieterich, S. B., Bradley, A. J., Nelan, E. P., & Whipple, A. L. (2016). THE SOLAR NEIGHBORHOOD. XXXVII. THE MASS-LUMINOSITY RELATION FOR MAIN-SEQUENCE M DWARFS*. *The Astronomical Journal*, 152(5), 141. <https://doi.org/10.3847/0004-6256/152/5/141>
- Dahm, S. E. (2015). REEXAMINING THE LITHIUM DEPLETION BOUNDARY IN THE PLEIADES AND THE INFERRED AGE OF THE CLUSTER. *The Astrophysical Journal*, 813(2), 108. <https://doi.org/10.1088/0004-637X/813/2/108>

- 173 Gomes, J. P. M. D. (2024). *Determinação da massa de estrelas jovens em aglomerados*
174 *estelares: Uma abordagem observacional e estatística* [Trabalho de Conclusão de Curso
175 (Bacharelado em Física)]. Universidade Estadual de Santa Cruz (UESC).
- 176 Gossage, S., Conroy, C., Dotter, A., Choi, J., Rosenfield, P., Cargile, P., & Dolphin, A. (2018).
177 Age Determinations of the Hyades, Praesepe, and Pleiades via MESA Models with Rotation.
178 *The Astrophysical Journal*, 863(1), 67. <https://doi.org/10.3847/1538-4357/aad0a0>
- 179 Herczeg, G. J., & Hillenbrand, L. A. (2015). EMPIRICAL ISOCHRONES FOR LOW MASS
180 STARS IN NEARBY YOUNG ASSOCIATIONS. *The Astrophysical Journal*, 808(1), 23.
181 <https://doi.org/10.1088/0004-637X/808/1/23>
- 182 Hillenbrand, L. A. (1997). On the Stellar Population and Star-Forming History of the Orion
183 Nebula Cluster. *The Astronomical Journal*, 113, 1733. <https://doi.org/10.1086/118389>
- 184 Lebreton, Y., Goupil, M.-J., & Montalbán, J. (2014). Stellar evolutionary models: Current
185 status and challenges. *Astrophysics and Space Science*, 109, 23–38. <https://doi.org/10.1007/s10509-014-1848-1>
- 187 Lodieu, N., Pérez-Garrido, A., Smart, R. L., & Silvotti, R. (2019). A 5D view of the
188 α Per, Pleiades, and Praesepe clusters. *Astronomy & Astrophysics*, 628, A66. <https://doi.org/10.1051/0004-6361/201935533>
- 190 Mann, A. W., Dupuy, T., Kraus, A. L., Gaidos, E., Ansdell, M., Ireland, M., Rizzuto,
191 A. C., Hung, C.-L., Dittmann, J., Factor, S., Feiden, G., Martinez, R. A., Ruíz-
192 Rodríguez, D., & Chia Thao, P. (2019). How to Constrain Your M Dwarf. II. The
193 Mass–Luminosity–Metallicity Relation from 0.075 to 0.70 Solar Masses. *The Astrophysical*
194 *Journal*, 871(1), 63. <https://doi.org/10.3847/1538-4357/aaf3bc>
- 195 Paegert, M., Stassun, K. G., Collins, K. A., Pepper, J., Torres, G., Jenkins, J., Twicken, J. D.,
196 & Latham, D. W. (2021). *TESS Input Catalog versions 8.1 and 8.2: Phantoms in the 8.0*
197 *Catalog and How to Handle Them*. arXiv. <https://doi.org/10.48550/ARXIV.2108.04778>
- 198 Pecaú, M. J., & Mamajek, E. E. (2013). INTRINSIC COLORS, TEMPERATURES, AND
199 BOLOMETRIC CORRECTIONS OF PRE-MAIN-SEQUENCE STARS. *The Astrophysical*
200 *Journal Supplement Series*, 208(1), 9. <https://doi.org/10.1088/0067-0049/208/1/9>
- 201 Pecaú, M. J., Mamajek, E. E., & Bubar, E. J. (2012). A REVISED AGE FOR UPPER
202 SCORPIUS AND THE STAR FORMATION HISTORY AMONG THE F-TYPE MEMBERS
203 OF THE SCORPIUS-CENTAURUS OB ASSOCIATION. *The Astrophysical Journal*, 746(2),
204 154. <https://doi.org/10.1088/0004-637X/746/2/154>
- 205 Rebull, L. M., Stauffer, J. R., Cody, A. M., Hillenbrand, L. A., David, T. J., & Pinsonneault,
206 M. (2018). Rotation of Low-mass Stars in Upper Scorpius and ρ Ophiuchus with K2. *The*
207 *Astronomical Journal*, 155(5), 196. <https://doi.org/10.3847/1538-3881/aab605>
- 208 Serenelli, A., Weiss, A., Aerts, C., Angelou, G. C., Baroch, D., Bastian, N., Beck, P. G.,
209 Bergemann, M., Bestenlehner, J. M., Czekala, I., Elias-Rosa, N., Escorza, A., Van Eylen,
210 V., Feuillet, D. K., Gandolfi, D., Gieles, M., Girardi, L., Lebreton, Y., Lodieu, N., ... Zwintz,
211 K. (2021). Weighing stars from birth to death: Mass determination methods across the
212 HRD. *The Astronomy and Astrophysics Review*, 29(1), 4. <https://doi.org/10.1007/s00159-021-00132-9>
- 214 Stassun, K. G., Oelkers, R. J., Paegert, M., Torres, G., Pepper, J., Lee, N. D., Collins, K.,
215 Latham, D. W., Muirhead, P. S., Chittidi, J., Rojas-Ayala, B., Fleming, S. W., Rose, M.
216 E., Tenenbaum, P., Ting, E. B., Kane, S. R., Barclay, T., Bean, J. L., Brassuer, C. E., ...
217 Winn, J. N. (2019). The Revised TESS Input Catalog and Candidate Target List. *The*
218 *Astronomical Journal*, 158(4), 138. <https://doi.org/10.3847/1538-3881/ab3467>