

A PCA plot showing the first two principal components, PC1 (x-axis) and PC2 (y-axis). The x-axis ranges from -4 to 8, and the y-axis ranges from -4 to 4. Dashed lines indicate the PC1 = 0 and PC2 = 0 axes. The plot displays a large number of data points, many of which are colored (blue, orange, green, red, yellow, pink, grey) and have black outlines. A yellow star with a black outline is located at approximately PC1 = -1.2 and PC2 = -1.2, highlighting a specific cluster of points. The points are distributed across the plot, with some clusters being more distinct than others.

A PCA plot showing the first two principal components, PC1 (x-axis) and PC2 (y-axis). The x-axis ranges from -4 to 8, and the y-axis ranges from -4 to 4. The plot displays a large number of data points colored by cluster. A yellow star with a black outline is located at approximately PC1 = -1.2 and PC2 = -1.2, marking a specific point of interest within a cluster of blue, green, and pink points. Other notable points include a large blue point at PC1 ≈ -3.8, PC2 ≈ -1.2; a large orange point at PC1 ≈ -0.5, PC2 ≈ 3.2; a large red point at PC1 ≈ 1.2, PC2 ≈ 1.2; and a large green point at PC1 ≈ 1.8, PC2 ≈ -1.2. Dashed lines indicate the PC1 = 0 and PC2 = 0 axes.

A PCA plot showing the first two principal components, PC1 (x-axis) and PC2 (y-axis). The x-axis ranges from -4 to 8, and the y-axis ranges from -4 to 4. Dashed lines indicate the PC1 = 0 and PC2 = 0 axes. The plot displays a large number of data points, many of which are colored (blue, orange, green, red, yellow, pink, grey) and have black outlines. A yellow star with a black outline is located at approximately PC1 = -1.2 and PC2 = -1.2, highlighting a specific cluster of points. The points are distributed across the plot, with some clusters being more distinct than others.