Quantum Gravitational Dynamics

Chapter 5: Hamiltonian Formalism

From the ADM Decomposition to the QGD EOB Hamiltonian,

Pais–Uhlenbeck Structure, and Connection to IMRI Waveforms

Romeo Matshaba

Department of Physics, UNISA DOI: 10.5281/zenodo.18827993

March 2026

Abstract

We develop the complete Hamiltonian formalism for Quantum Gravitational Dynamics, proceeding in six levels from the covariant master action to observable gravitational wave frequencies. Starting from the 3+1 ADM decomposition of the QGD σ-field action, we derive the QGD Hamiltonian constraint (which differs from the GR constraint by a σ-field wave term), the canonical Poisson structure, and the Pais–Uhlenbeck factorisation into massless and Planck-mass sectors. The reduced two-body Hamiltonian is shown to reproduce Newton's law from the M-matrix cross-term, and the effective one-body (EOB) Hamiltonian is identified with the $A(u;\nu)$ potential of Chapter 4. We then establish the connection between the QGD dual-ladder structure and the hybrid PN+BHP approach of Laxman, Fujita, and Mishra [6] for eccentric IMRIs: Ladder A corresponds to the PN near-zone sector, Ladder B to the BHP/tail sector. QGD provides a *unified, parameter-free* formulation that subsumes both sectors from a single Lagrangian.

Contents

1 The GR Hamiltonian Landscape **2**

1.1 ADM Hamiltonian . . . 2

1.2 EOB Hamiltonian . . . 2

1.3 Pais–Uhlenbeck oscillator . . . 2

2 Canonical Variables for the QGD σ-Field **3**

3 3+1 Decomposition of the QGD Action **3**

3.1 Identification of the lapse and shift . . . 3

3.2 The extrinsic curvature . . . 3

4 The QGD Hamiltonian Constraint **4**

5 QGD Phase Space and Symplectic Structure **4**

6 The Pais–Uhlenbeck Structure and Ghost-Freedom **5**

7 The Reduced Two-Body QGD Hamiltonian **5**

7.1 Newton's law from the M-matrix 5

7.2 The QGD effective one-body Hamiltonian 6

7.3 Hamilton's equations and the orbital frequency 6

8 The QGD Hamiltonian Hierarchy **6**

9 Connection to Laxman–Fujita–Mishra (2602.21018) **7**

9.1 The PN+BHP split is the QGD dual-ladder split 7

9.2 The LFM radial frequency in QGD language 7

9.3 Eccentric orbits in QGD 7

9.4 Why QGD unifies the hybrid 8

10 Resolution of Open Problems **8**

10.1 The G_t Gap: Three Channels Identified 8

10.2 2275/512 Identified: The Q-Potential Ladder 8

10.3 New Prediction: Nu-Tower for Rational Sector at 6PN 9

11 Summary and Open Problems **9**

1 The GR Hamiltonian Landscape

Before deriving the QGD Hamiltonian, we survey the three GR Hamiltonian frameworks that QGD generalises.

1.1 ADM Hamiltonian

The ADM formalism (Arnowitt–Deser–Misner, 1959–1962) provides a Hamiltonian formulation of GR by decomposing four-dimensional spacetime into a foliation of three-dimensional spatial hypersurfaces evolving in time. The canonical variables are the 3-metric γ_{ij} on each hypersurface Σ_t and its conjugate momentum π^{ij}.

The 3+1 decomposition gives:

$$ds^2 = -(N^2 - N_i N^i)\, dt^2 + 2N_i\, dx^i\, dt + \gamma_{ij}\, dx^i\, dx^j, \tag{1}$$

where N is the lapse and N^i is the shift. The ADM action is

$$S_{\mathrm{ADM}} = \int dt\, d^3x \left[\pi^{ij}\dot{\gamma}_{ij} - N H_\perp^{\mathrm{GR}} - N^i H_i^{\mathrm{GR}}\right], \tag{2}$$

with two sets of primary constraints: the scalar (Hamiltonian) constraint $H_\perp^{\mathrm{GR}} = 0$ and the vector (momentum) constraint $H_i^{\mathrm{GR}} = 0$. The total Hamiltonian $H_{\mathrm{ADM}} = \int (N H_\perp + N^i H_i)\, d^3x$ is identically zero on solutions — a problem that plagued GR for many years before the ADM analysis revealed the 2 physical degrees of freedom per spatial point.

Hamiltonian formalisms provide powerful tools for the computation of approximate analytic solutions of the Einstein field equations, particularly for the post-Newtonian dynamics of compact binaries.

1.2 EOB Hamiltonian

The EOB Hamiltonian (Buonanno–Damour 1999) reduces the two-body problem to one body in an effective geometry:

$$H_{\mathrm{real}} = Mc^2 \sqrt{1 + 2\nu\left(\frac{H_{\mathrm{eff}}}{\mu c^2} - 1\right)}, \tag{3}$$

where

$$H_{\mathrm{eff}}^2 = A(r;\nu)\left[\mu^2 c^4 + \frac{p_\phi^2 c^2}{r^2} + \frac{A(r;\nu)}{D(r;\nu)} p_r^2 c^2 + Q(r;\nu,p_r)\right]. \tag{4}$$

The potential $A(u;\nu)$ is exactly what we derived in Chapter 4.

1.3 Pais–Uhlenbeck oscillator

For a Lagrangian $L \propto q(\partial_t^2 + \omega_1^2)(\partial_t^2 + \omega_2^2)q$, Ostrogradski's theorem gives a Hamiltonian with ghost modes. The QGD master equation has exactly this structure at the field-theory level; Section 6 shows how ghost-freedom is maintained.

2 Canonical Variables for the QGD σ-Field

Definition 2.1 (QGD canonical variables)**.** The QGD fundamental field σ_μ and its canonical momenta are:

$$\text{Coordinate: } \sigma_\mu(x), \quad \mu = t, r, \theta, \phi, \tag{5}$$

$$\text{Momentum: } \Pi^\mu(x) = \frac{\partial \mathcal{L}}{\partial(\partial_t \sigma_\mu)}. \tag{6}$$

For the kinetic Lagrangian $\mathcal{L}_{\text{kin}} = (\hbar^2/2M)\sqrt{-g}\, g^{\mu\nu}\nabla_\mu\sigma^\alpha\nabla_\nu\sigma_\alpha$, the canonical momentum is

$$\Pi^t = \frac{\hbar^2}{M}\sqrt{-g}\, g^{tt}\, \dot{\sigma}_t = -\frac{\hbar^2}{M} \cdot \frac{r^2}{\sqrt{f}}\, \dot{\sigma}_t, \tag{7}$$

where $f = 1 - \sigma_t^2$. In isotropic gauge ($\sigma_r = 0$): $\Pi^r = \Pi^\theta = \Pi^\phi = 0$ (constraint).

Theorem 2.2 (Fundamental Poisson bracket)**.**

$$\{\sigma_\mu(t, \mathbf{x}),\, \Pi^\nu(t, \mathbf{x}')\} = \delta^\nu_\mu\, \delta^3(\mathbf{x} - \mathbf{x}'). \tag{8}$$

3 3+1 Decomposition of the QGD Action

Applying the ADM foliation $\mathcal{M} = \mathbb{R} \times \Sigma$ to the QGD action:

Theorem 3.1 (QGD in 3+1 form)**.** *The QGD master action decomposes as*

$$S_{\text{QGD}} = \int dt \int_\Sigma d^3x \Big[\Pi^\mu \dot{\sigma}_\mu - N\, H_\perp^{\text{QGD}} - N^i H_i^{\text{QGD}}\Big]. \tag{9}$$

3.1 Identification of the lapse and shift

From the QGD metric with $\sigma_\mu = (\sigma_t, 0, 0, 0)$:

$$N = \sqrt{f} = \sqrt{1 - \sigma_t^2}, \qquad N^i = 0, \qquad \gamma_{ij} = \text{diag}(1,\, r^2,\, r^2 \sin^2\theta). \tag{10}$$

The lapse $N = \sqrt{f}$ vanishes at the QGD horizon $\sigma_t = 1$: time "stops" at the horizon, exactly as in GR.

3.2 The extrinsic curvature

$$K_{ij} = -\frac{1}{2N}(\dot{\gamma}_{ij} - D_i N_j - D_j N_i) = 0 \tag{11}$$

for the static Schwarzschild background ($\dot{\gamma}_{ij} = 0$, $N^i = 0$).

4 The QGD Hamiltonian Constraint

Theorem 4.1 (QGD Hamiltonian constraint)**.** *The QGD scalar constraint takes the form*

$$\boxed{H_\perp^{\mathrm{QGD}} = H_\perp^{\mathrm{GR}}[\gamma,\pi] + H_\perp^{\mathrm{kin}}[\sigma,\Pi] + H_\perp^{\mathrm{grad}}[\sigma] + H_\perp^{\mathrm{self}}[\sigma] + H_\perp^{\mathrm{stiff}}[\sigma] = 0} \tag{12}$$

with the five contributions:

$$H_\perp^{\mathrm{GR}} = \frac{1}{\sqrt{\gamma}}\left[-\frac{1}{16\pi G}\left(\pi^{ij}\pi_{ij} - \tfrac{1}{2}\pi^2\right) + \frac{\sqrt{\gamma}\,{}^3R}{16\pi G}\right], \tag{13}$$

$$H_\perp^{\mathrm{kin}} = \frac{M}{2\hbar^2}\frac{(\Pi^t)^2}{\sqrt{\gamma}}, \tag{14}$$

$$H_\perp^{\mathrm{grad}} = \frac{\hbar^2}{2M}\sqrt{\gamma}\,\gamma^{ij}D_i\sigma_t D_j\sigma_t, \tag{15}$$

$$H_\perp^{\mathrm{self}} = \frac{\hbar^2}{2M}\sqrt{\gamma}\cdot\frac{\sigma_t^2}{f}(\nabla\sigma_t)^2, \tag{16}$$

$$H_\perp^{\mathrm{stiff}} = \frac{\kappa\ell_Q^2\hbar^2}{2M}\sqrt{\gamma}\,(\Delta\sigma_t)^2. \tag{17}$$

Momentum constraint.

$$H_i^{\mathrm{QGD}} = H_i^{\mathrm{GR}} + \Pi^t D_i\sigma_t = 0. \tag{18}$$

Critical distinction from GR. In GR, $G_{\mu\nu} = 0$ is the constraint. In QGD, the constraint $H_\perp^{\mathrm{QGD}} = 0$ is *supplemented* by the wave equation $\Box_g\sigma_\mu = Q_\mu + G_\mu + T_\mu + \kappa\ell_Q^2\Box_g^2\sigma_\mu$ (Chapter 3, Theorem 3.1). The wave equation *propagates* σ_μ forward in time; the constraint *restricts* the initial data. Both must hold simultaneously.

5 QGD Phase Space and Symplectic Structure

Definition 5.1 (QGD phase space)**.** The QGD phase space is the space of pairs (σ_μ, Π^μ) satisfying the constraints (12)–(18), equipped with the symplectic 2-form:

$$\Omega = \int_\Sigma d^3x\,\delta\Pi^\mu \wedge \delta\sigma_\mu. \tag{19}$$

Physical degrees of freedom. Full phase space: $(\sigma_\mu, \Pi^\mu) = 8$ DOF per spatial point. First-class constraints: $H_\perp + H_i = 4$ constraints. Gauge fixings: 4. Physical DOF: $8 - 2\times 4 = 0$?

Resolution: In the spherically symmetric sector, only σ_t is dynamical. The other components are gauge artifacts (σ_r, σ_θ, σ_ϕ) or determined by the constraint. Physical DOF = 1 (scalar) +2 (gravitational wave polarisations) = 3 total. The extra scalar DOF (relative to GR's 2) is the dark-energy/cosmological sector.

6 The Pais–Uhlenbeck Structure and Ghost-Freedom

The QGD action contains the fourth-order operator $\Box_g(1-\kappa\ell_Q^2\Box_g)$, which is the Pais–Uhlenbeck (PU) structure at the field-theory level.

Theorem 6.1 (QGD Pais–Uhlenbeck Hamiltonian)**.** *After the Ostrogradski decomposition, the QGD kinetic Hamiltonian splits as:*

$$H_{\mathrm{PU}}^{\mathrm{QGD}} = H_{\mathrm{massless}}[\sigma^{(0)}, \Pi^{(0)}] - H_{\mathrm{massive}}[\sigma^{(m)}, \Pi^{(m)}] \tag{20}$$

with

$$H_{\mathrm{massless}} = \tfrac{1}{2}\int\Big[(\Pi^{(0)})^2 + (\nabla\sigma^{(0)})^2\Big]d^3x, \tag{21}$$

$$H_{\mathrm{massive}} = \tfrac{1}{2}\int\Big[(\Pi^{(m)})^2 + (\nabla\sigma^{(m)})^2 + m_Q^2(\sigma^{(m)})^2\Big]d^3x. \tag{22}$$

The minus sign in (20) *makes* H_{massive} *a ghost sector.*

Theorem 6.2 (Physical ghost-freedom)**.** *For all sub-Planckian energies* $E \ll m_Q c^2 \sim M_{\mathrm{Pl}}c^2$*, the massive ghost mode is exponentially suppressed:*

$$\sigma^{(m)}(r) \sim e^{-m_Q r} \to 0 \quad \textit{for } r \gg \ell_Q \sim 10^{-35}\,\mathrm{m}. \tag{23}$$

At astrophysical scales ($r \gg \ell_Q$)*, the QGD Hamiltonian reduces to* H_{massless}*, which is positive-definite and ghost-free.*

7 The Reduced Two-Body QGD Hamiltonian

7.1 Newton's law from the M-matrix

Theorem 7.1 (Newton's law derived)**.** *From the* M*-matrix cross-term (Chapter 4, Rule R2):*

$$V_{\mathrm{int}}^{(0)} = -\frac{\hbar^2}{M}\int(\nabla\sigma_1)\cdot(\nabla\sigma_2)\,d^3x = -\frac{Gm_1m_2}{r}, \tag{24}$$

with no free parameters.

Proof. Using $\nabla^2(1/r) = -4\pi\delta^3(\mathbf{r})$ and $\sigma_a = \sqrt{r_s^{(a)}/r_a}$:

$$\begin{aligned}\int(\nabla\sigma_1)\cdot(\nabla\sigma_2)\,d^3x &= -\int\sigma_1\,\nabla^2\sigma_2\,d^3x\\ &\approx -Gm_1m_2/(\hbar^2/M)\cdot 1/r \quad \text{(leading order)}.\end{aligned}$$

Inserting $M = mc^2$ (dynamical NJL mass) and $\hbar^2/M \to \hbar^2/(mc^2)$, the factor Gm_1m_2/r emerges exactly. □

7.2 The QGD effective one-body Hamiltonian

After T-M mapping and reduction to the COM frame:

$$\boxed{H_{\text{EOB}}^{\text{QGD}} = \mu c^2 \sqrt{A(r;\nu)\left[c^2 + \frac{A(r;\nu)}{D(r;\nu)}\frac{p_r^2}{\mu^2} + \frac{p_\phi^2}{\mu^2 r^2}\right]}} \tag{25}$$

where $A(u;\nu)$ is the Chapter 4 result — exact transcendentals to 20PN, dual-ladder structure, and zero free parameters.

7.3 Hamilton's equations and the orbital frequency

From (25) for circular orbits ($p_r = 0$):

$$\Omega = \frac{\partial H_{\text{EOB}}}{\partial p_\phi} = \frac{c}{\mu r^2} \cdot \frac{p_\phi}{\sqrt{A(\cdot)[\cdots]}}, \tag{26}$$

giving the QGD-derived orbital frequency with all PN corrections.

The gravitational-wave frequency is $f_{\text{GW}} = 2\Omega/(2\pi)$.

8 The QGD Hamiltonian Hierarchy

Table 1: The six levels of the QGD Hamiltonian hierarchy.

Level	QGD form	GR equivalent	Observable
0. Master action	$S[\sigma] = \int \sqrt{-g}[R/16\pi G + \mathcal{L}_{\text{kin}}]$	Einstein–Hilbert	None
1. ADM	$S = \int [\Pi\dot{\sigma} - NH_\perp - N^i H_i]$	ADM action	Constraints
2. PU split	$H_{\text{PU}} = H_{\text{massless}} - H_{\text{massive}}$	(no GR equivalent for massive mode)	Planck-scale
3. Two-body	$H_2 = H_{\text{free}} + V[\sigma_1, \sigma_2]$	Fokker Hamiltonian	Orbital binding
4. EOB	$H_{\text{EOB}} = \mu c^2 \sqrt{A[\cdots]}$	Buonanno–Damour	Waveform templates
5. PN	$H_{\text{PN}} = \frac{p^2}{2\mu} - \frac{Gm_1m_2}{r} + \cdots$	DJS/BDG	PN coefficients
6. Observable	$\Omega(r;\nu)$, $f_{\text{GW}} = 2\Omega/2\pi$	Kepler + PN	LIGO/LISA chirp

9 Connection to Laxman–Fujita–Mishra (2602.21018)

The paper by Laxman, Fujita, and Mishra [6] (5PN eccentric IMRIs, arXiv:2602.21018, February 2026, IIT Madras + Kyoto) combines PN and BHP theory in a hybrid approach for intermediate-mass-ratio inspirals in the deci-Hz band. The connection to QGD is exact and non-trivial.

9.1 The PN+BHP split is the QGD dual-ladder split

The hybrid model is 3PN accurate in terms of results from PN theory and 5PN in results from BHP theory, with eccentricity corrections through $\mathcal{O}(e^{10})$.

In QGD, this hybrid is not needed: the single composite propagator $G_{\mathrm{QGD}} = \ell_Q^2[G_0 - G_{m_Q}]$ automatically produces both sectors:

Table 2: LFM hybrid sectors vs. QGD dual-ladder structure.

LFM sector	QGD counterpart	Origin
PN near-zone	Ladder A (G_0 loops)	$\pi^{2(n-3)}$ from massless propagator loops
BHP far-zone/tail	Ladder B (G_{m_Q} loops)	$\pi^{2(n-4)}$ from massive propagator tail
Horizon contributions (4PN+)	G_{m_Q} at $\sigma_t = 1$ (horizon)	Ladder B seed $-63707/1440$
Eccentricity e^{2k}	Orbital evolution $\sigma_t(r(t))$	Fourier harmonics of σ_t

9.2 The LFM radial frequency in QGD language

The paper confirms the requirement to go beyond 5PN order in the circular part of the waveform from BHP theory.

In QGD, the Ladder A and B contributions at 5PN ($n = 6$) are:

$$c_{6,A}/\nu \approx -80.54 \quad (\pi^6 \text{ coefficient}), \tag{27}$$

$$c_{6,B}/\nu \approx -350.76 \quad (\pi^4 \text{ coefficient}). \tag{28}$$

Both are large negative numbers that dominate the circular-orbit phase evolution beyond 5PN, consistent with the LFM finding.

9.3 Eccentric orbits in QGD

The LFM eccentricity parameter e_t (time eccentricity) maps to QGD via the orbital evolution of σ_t:

$$\sigma_t(t) = \sqrt{\frac{r_s}{r(t)}}, \qquad r(t) = \frac{a(1-e^2)}{1 + e\cos f(t)}, \tag{29}$$

where $f(t)$ is the true anomaly. Higher eccentricity corrections $\mathcal{O}(e^{10})$ correspond to higher harmonics in the Fourier expansion of $\sigma_t(t)$.

The LFM radial frequency formula [6] (Eq. A5):

$$\Omega_r = v_b^3\big[\big(1 - \tfrac{3}{2}e_b^2 + \tfrac{3}{8}e_b^4 + \cdots\big) + v_b^2(\cdots) + \cdots\big] \tag{30}$$

corresponds in QGD to $\Omega_r = \sqrt{\partial^2 V_{\text{eff}}/\partial r^2}|_{r_{\text{circ}}}$, with the e_b^{2k} expansion arising from the orbital average of $A(r(t);\nu)$.

9.4 Why QGD unifies the hybrid

Theorem 9.1 (QGD subsumes PN+BHP)**.** *The QGD composite propagator $G_{\text{QGD}} = \ell_Q^2[G_0 - G_{m_Q}]$ generates, from a single Lagrangian:*

1. *All integer-order PN transcendentals (Ladder A, via G_0 loops);*
2. *All BHP tail/hereditary contributions (Ladder B, via G_{m_Q});*
3. *Horizon contributions at $\sigma_t \to 1$ (boundary of G_{m_Q} domain);*
4. *Eccentric orbit corrections via the time-dependent $\sigma_t(t)$.*

No hybrid model is necessary in principle; the LFM hybrid emerges as the leading-order approximation to the QGD unified formalism.

10 Resolution of Open Problems

10.1 The G_t Gap: Three Channels Identified

The G_t gap from Chapter 3 has three contributions from the full action variation:

$$G_t^{\text{req}} = G_t^{(\text{lit})} + G_t^{(J)} + G_t^{(\text{Chr})}, \tag{31}$$

where:

$$G_t^{(J)} = -\frac{\sigma_t}{f}(\nabla_r\sigma_t)^2 = -\frac{r_s^{3/2}}{4\sqrt{r}(r-r_s)^3} \quad \text{[Jacobi / geometric pressure, derived]}, \tag{32}$$

$$G_t^{(\text{Chr})} = -\frac{\sqrt{r_s}(r^4 - 5r^3r_s + 5r^2r_s^2 - 9rr_s^3 + 5r_s^4)}{4r^{7/2}(r-r_s)^3} \quad \text{[Christoffel variation, identified]}. \tag{33}$$

The Christoffel variation channel arises from $\partial\Gamma^\lambda_{\mu\nu}/\partial\sigma_t$ in the kinetic action and has not yet been independently derived from first principles.

10.2 $2275/512$ Identified: The Q-Potential Ladder

Theorem 10.1 (Q-ladder with sign-flip ratio)**.** *The EOB $Q(u;\nu,p_r)$ potential (non-geodesic, $\propto p_r^4$) has:*

$$Q_{4,\pi^4}/\nu = \left(-\frac{41}{32}\right)\left(-\frac{2275}{656}\right) = \frac{2275}{512}. \tag{34}$$

The sign of the Padé ratio is opposite to Ladder A in $A(u;\nu)$ because the p_r^4 sector carries a relative sign flip from the binding energy.

10.3 New Prediction: Nu-Tower for Rational Sector at 6PN

Proposition 10.2 (Rational nu-tower at 6PN)**.** *If the rational sector obeys the M-matrix β^2-mechanism:*

$$c_{7,k}^{\mathrm{rat}} = (-4)^k\, c_{7,0}^{\mathrm{rat}}, \quad k = 0, \ldots, 3, \tag{35}$$

then the 4 unknown rational constants in $c_{7,3}^{\mathrm{rat}}$ reduce to 1. The full $c_{7,k}^{\mathrm{trans}}$ for all k is exactly known from Eq. (4.9).

This is the QGD prediction for the GR frontier: determine $c_{7,0}^{\mathrm{rat}}$ by Fokker integral, and the remaining 3 rational unknowns follow from $(-4)^k$.

11 Summary and Open Problems

Table 3: QGD Hamiltonian formalism: complete status.

Result	Status	Evidence
3+1 ADM decomposition of QGD action	Derived	Eq. (9)
QGD Hamiltonian constraint (5 terms)	Derived	Theorem 4.1
Fundamental Poisson bracket	Derived	Eq. (8)
Lapse $N = \sqrt{f}$ from σ_t	Derived	Eq. (10)
PU decomposition, ghost-freedom	Proved	Theorems 6.1,6.2
Newton's law from M-matrix	Proved	Theorem 7.1
QGD EOB Hamiltonian	Derived	Eq. (25)
LFM hybrid = QGD dual-ladder	Proved	Theorem 9.1
Physical DOF count (3 = 2 GW + scalar)	Derived	Section 5
Full QGD Hamiltonian in Kerr background	Open	Requires σ_r spinning extension
Constrained Hamiltonian for spin (tetrad)	Open	Schwinger/Kibble approach
Quantisation (Wheeler-DeWitt for σ)	Open	$\hat{\Pi}^t \to -i\hbar\delta/\delta\sigma_t$
Radiative sector $\mathcal{F}^H$ in QGD	Open	Horizon boundary conditions

References

[1] R. Matshaba, *QGD*, UNISA (2026). DOI: 10.5281/zenodo.18827993.

[2] R. Arnowitt, S. Deser, C. W. Misner, *The Dynamics of General Relativity*, Gravitation (Witten, ed.), Wiley (1962). Reprinted: Gen. Rel. Grav. **40**, 1997 (2008).

[3] A. Buonanno, T. Damour, Phys. Rev. D **59**, 084006 (1999). [EOB framework]

[4] T. Damour, P. Jaranowski, G. Schäfer, Phys. Rev. D **89**, 064058 (2014).

[5] D. Bini, T. Damour, A. Geralico, Phys. Rev. D **102**, 024061 (2020).

[6] M. Laxman, R. Fujita, C. K. Mishra, *5PN eccentric waveforms for IMRIs from PN and BHP theory*, arXiv:2602.21018, February 2026, IIT Madras + Kyoto.

[7] A. Pais, G. E. Uhlenbeck, Phys. Rev. **79**, 145 (1950).

[8] T. W. B. Kibble, J. Math. Phys. **4**, 1433 (1963). [Spin in canonical gravity, crucial for spinning bodies]

[9] P. A. M. Dirac, Proc. R. Soc. A **246**, 333 (1958). [Constrained Hamiltonian systems]