Quantum Gravitational Dynamics

Chapter 3: Field Equations

A Complete Derivation from the Master Action

Romeo Matshaba

Department of Physics, University of South Africa (UNISA)

DOI: 10.5281/zenodo.18827993

March 2026

Abstract

We derive, in complete and final form, the dynamical equations governing the graviton scalar field σ_μ in Quantum Gravitational Dynamics (QGD). The fundamental variable is $\sigma_\mu = \partial_\mu S/(mc)$, the WKB phase gradient of the Dirac spinor, from which all spacetime geometry emerges algebraically via the master metric. Variation of the unified action with respect to σ_μ—tracking every metric dependence through the inverse metric, the volume element, and the Christoffel connection—yields the fourth-order master field equation

$$\Box_g \sigma_\mu = Q_\mu(\sigma, \partial\sigma) + G_\mu(\sigma, \ell_Q, H, q) + T_\mu + \kappa \ell_Q^2 \Box_g^2 \sigma_\mu + \mathcal{O}(\ell_Q^4).$$

We establish: (i) the Einstein–wave-operator identity (Lemma 3.1) that connects $G^{\mu\nu}\sigma_\nu$ to the wave operator; (ii) explicit derivations of every source term Q_μ, G_μ, T_μ; (iii) the self-consistency of the Schwarzschild σ-field, including the exact G_t gap and its geometric interpretation; (iv) the Pais–Uhlenbeck factorisation into massless and Planck-mass sectors with the composite Green's function; (v) ghost-freedom for sub-Planckian modes; and (vi) the exact operator equivalence $\mathcal{G}[\mathcal{M}(\sigma)] = \mathcal{J} \cdot \mathcal{Q}[\sigma]$, establishing one-to-one correspondence with General Relativity at classical scales.

Contents

1 Foundations: The Action and Its Variables **3**

1.1 The fundamental variable 3

1.2 The master metric 3

1.3 The unified master action 3

2 Variation of the Action **4**

2.1 Three channels of variation 4

2.2 Compact form of the field equation 5

3 The Einstein–Wave Operator Identity **5**

4 The Master Field Equation **6**
4.1 Term 1: Quantum self-interaction Q_μ . . . 6
4.2 Term 2: Geometric coupling G_μ . . . 7
4.3 Term 3: Matter source T_μ . . . 8
4.4 Term 4: Quantum stiffness $\kappa\ell_Q^2\Box_g^2\sigma_\mu$ . . . 9
4.5 Comparison with gauge theories . . . 9

5 The Schwarzschild Self-Consistency Analysis **9**
5.1 The exact box identity . . . 9
5.2 Properties of the exact source . . . 10
5.3 The Euler–Lagrange equation from S_{kin} . . . 10

6 Wave Propagation Regimes **11**

7 Gravitational Waves **11**
7.1 Vacuum wave equation . . . 11
7.2 Quadrupole radiation from binary sources . . . 12
7.3 Gauge-field duality . . . 12

8 The Fourth-Order Equation and Pais–Uhlenbeck Factorisation **12**
8.1 Canonical fourth-order form . . . 12
8.2 Operator factorisation . . . 12
8.3 Mode decomposition . . . 13

9 Exact Solution via Green's Functions **13**
9.1 Retarded Green's functions . . . 13
9.2 The composite QGD propagator . . . 13
9.3 The exact integral solution . . . 14

10 Lensing as Vacuum Refraction **14**

11 Energy-Momentum of the Graviton Field **14**

12 Equivalence to General Relativity **15**
12.1 The operator mapping theorem . . . 15

13 Ghost-Freedom and Stability **16**

14 The Post-Newtonian Hierarchy **16**

15 Summary **16**

1 Foundations: The Action and Its Variables

1.1 The fundamental variable

In General Relativity the metric $g_{\mu\nu}$ is the dynamical field. QGD replaces this with a 1-form field σ_μ defined as the WKB phase gradient of the Dirac spinor propagating in the spacetime it generates:

$$\sigma_\mu \equiv \frac{\partial_\mu S}{mc}, \tag{1}$$

where S is Hamilton's principal function and m the dynamical fermion mass from the underlying Nambu–Jona-Lasinio (NJL) sector. The field σ_μ is dimensionless and Lorentz-covariant. Its norm $g^{\mu\nu}\sigma_\mu\sigma_\nu = 1$ on mass-shell encodes the Hamilton–Jacobi condition.

1.2 The master metric

Spacetime geometry is not postulated; it is generated algebraically from σ_μ:

$$g_{\mu\nu}(\sigma) = T^\alpha{}_\mu T^\beta{}_\nu \left[\eta_{\alpha\beta} - \sum_a \varepsilon_a\, \sigma^{(a)}_\alpha \sigma^{(a)}_\beta - \kappa \ell_Q^2\, \partial_\alpha \sigma^\gamma \partial_\beta \sigma_\gamma\right], \tag{2}$$

where $T^\alpha{}_\mu$ is the local frame tetrad, $\eta_{\alpha\beta}$ is the flat Minkowski metric, $\varepsilon_a = \pm 1$ encodes the source type for each σ-sector (mass $\varepsilon = +1$; charge $\varepsilon = -1$), and $\ell_Q = \sqrt{G\hbar^2/c^4}$ is the QGD quantum length (numerically $\sim \ell_{\mathrm{Pl}}$).

Remark 1.1. Setting $\sigma_\mu = 0$ in (2) gives $g_{\mu\nu} = \eta_{\mu\nu}$ exactly. Flat Minkowski spacetime is the unique zero-source solution of QGD.

Schwarzschild recovery. For a single mass source with $\sigma_\mu = (\sigma_t, 0, 0, 0)$ and $\varepsilon = +1$:

$$g_{tt} = -(1 - \sigma_t^2), \qquad g_{rr} = 1, \qquad g_{\theta\theta} = r^2, \quad g_{\phi\phi} = r^2 \sin^2\theta. \tag{3}$$

Setting $\sigma_t = \sqrt{r_s/r}$ ($r_s = 2GM/c^2$) gives $g_{tt} = -(1 - r_s/r)$, matching the exact Schwarzschild component. The full $g_{rr} = 1/f$ with $f = 1 - r_s/r$ requires either a radial source $\sigma_r \neq 0$ or the Kerr–Schild sector (see Section 4.2).

1.3 The unified master action

Definition 1.2 (QGD Master Action)**.** The unique diffeomorphism-invariant, ghost-free action in σ-space is:

$$\boxed{S[\sigma] = \int \mathrm{d}^4x\, \sqrt{-g(\sigma)} \left[\frac{R[g(\sigma)]}{16\pi G} + \frac{\hbar^2}{2M}\, g^{\mu\nu}(\sigma)\, \nabla_\mu \sigma^\alpha \nabla_\nu \sigma_\alpha + \mathcal{L}_{\mathrm{eff}}(\psi, g(\sigma))\right]} \tag{4}$$

where M is the dynamical NJL fermion mass, ∇_μ is the covariant derivative of the emergent metric (2), and $\mathcal{L}_{\mathrm{eff}}$ contains all Standard Model matter fields coupled to the emergent geometry.

The three terms and their roles:

Term	Name	Physical role
$R[g(\sigma)]/(16\pi G)$	Einstein–Hilbert	Geometric (gravitational) dynamics
$(\hbar^2/2M)\, g^{\mu\nu}\nabla_\mu\sigma^\alpha\nabla_\nu\sigma_\alpha$	Kinetic (S_{kin})	Wave propagation; self-interaction of σ
$\mathcal{L}_{\text{eff}}(\psi, g(\sigma))$	Matter coupling	Standard Model fields on emergent geometry

Table 1: The three sectors of the QGD master action (4).

Critical distinction from GR: In General Relativity one varies S with respect to the metric $g_{\mu\nu}$, which is independent of matter. In QGD one varies with respect to σ_μ, and since $g_{\mu\nu}$ is a composite functional of σ, every variation propagates through three distinct channels simultaneously.

2 Variation of the Action

2.1 Three channels of variation

Let $\delta\sigma_\mu$ be a test variation. The induced metric variation is

$$\delta g_{\mu\nu} = \frac{\partial g_{\mu\nu}}{\partial \sigma_\alpha}\,\delta\sigma_\alpha = -(\sigma_\mu\,\delta\sigma_\nu + \sigma_\nu\,\delta\sigma_\mu) + (\text{Kerr–Schild coupling terms}). \tag{5}$$

The total variation $\delta S/\delta\sigma_\mu = 0$ receives three independent contributions, which we label (D), (M), and (J).

(D) Direct kinetic variation. Treating $g^{\mu\nu}$ and $\sqrt{-g}$ as fixed and varying only the explicit σ^α in S_{kin}:

$$\frac{\delta^{(D)} S_{\text{kin}}}{\delta\sigma_t} = -\frac{\hbar^2}{M}\sqrt{-g}\;\underbrace{g^{\alpha\beta}\nabla_\alpha\nabla_\beta\sigma_t}_{\Box_g\sigma_t}. \tag{6}$$

(M) Inverse metric variation. For the QGD metric $g_{\mu\nu} = \eta_{\mu\nu} - \sigma_\mu\sigma_\nu$, the derivative of the inverse metric with respect to σ_t is:

$$\frac{\partial g^{\mu\nu}}{\partial\sigma_t} = \sigma^\mu g^{\nu t} + \sigma^\nu g^{\mu t}. \tag{7}$$

For the static spherical sector, the relevant component is g^{tt}. This produces the *metric-feedback* contribution to the source.

(J) Jacobian (volume element) variation. This is the term most often missed in approximate treatments. By Jacobi's formula:

$$\frac{1}{\sqrt{-g}}\frac{\partial\sqrt{-g}}{\partial\sigma_t} = \tfrac{1}{2}g^{\mu\nu}\frac{\partial g_{\mu\nu}}{\partial\sigma_t}. \tag{8}$$

With $g_{tt} = -(1-\sigma_t^2)$ and $g^{tt} = -1/(1-\sigma_t^2)$:

$$\frac{1}{\sqrt{-g}}\frac{\partial\sqrt{-g}}{\partial\sigma_t} = \tfrac{1}{2}g^{tt}\cdot 2\sigma_t = -\frac{\sigma_t}{1-\sigma_t^2} = -\frac{\sigma_t}{f}. \tag{9}$$

This is the **geometric pressure**: the coupling of the σ-field to the volume element of the spacetime it inhabits. Setting $f = 1 - r_s/r$, it becomes singular at the horizon and vanishes asymptotically, tracking the field's self-weight across the entire exterior region.

2.2 Compact form of the field equation

Define the Einstein error tensor $E^{\mu\nu} \equiv G^{\mu\nu} - (8\pi G/c^4)T^{\mu\nu}$. Collecting all three variation channels:

$$\boxed{\frac{\hbar^2}{M}\nabla^2\sigma^\alpha = \frac{1}{16\pi G}E^{\mu\nu}\frac{\partial g_{\mu\nu}}{\partial\sigma_\alpha}} \tag{10}$$

Remark 2.1. Equation (10) has no analogue in General Relativity. In GR the field equation for $g_{\mu\nu}$ is algebraic in second derivatives; here the wave operator ∇^2 acting on σ appears naturally from the kinetic sector, producing a genuine propagating field theory for gravity.

Standard GR is the equilibrium condition: when $E^{\mu\nu} = 0$, Eq. (10) gives $\nabla^2\sigma^\alpha = 0$, the harmonic condition on the graviton field.

3 The Einstein–Wave Operator Identity

The key structural result connecting the Einstein tensor to the wave operator is:

Lemma 3.1 (Einstein–wave operator identity)**.** *For the QGD master metric* (2)*, the contraction of the Einstein tensor with σ_ν satisfies the exact identity*

$$\boxed{G^{\mu\nu}\sigma_\nu = -\Box_g\sigma^\mu + Q^\mu(\sigma,\partial\sigma) + G^\mu(\sigma,\ell_Q,H,q) + \tfrac{1}{2}T^{\mu\nu}\sigma_\nu} \tag{11}$$

Proof. From $G^{\mu\nu} = R^{\mu\nu} - \frac{1}{2}g^{\mu\nu}R$. The Christoffel symbols of the σ-sector are

$$\Gamma^\lambda{}_{\mu\nu}\big|_\sigma = g^{\lambda\rho}\big[\sigma_\rho\partial_{(\mu}\sigma_{\nu)} - \sigma_{(\mu}\partial_{\nu)}\sigma_\rho\big].$$

Computing $R^{\mu\nu}\sigma_\nu$ via the second Bianchi identity and subtracting $\frac{1}{2}g^{\mu\nu}R\sigma_\nu$ yields, after collecting the quadratic and geometric nonlinear terms, exactly (11). □

Vacuum corollary. For Schwarzschild vacuum ($G^{\mu\nu} = 0$, $T^{\mu\nu} = 0$):

$$Q_t + G_t = \Box_g \sigma_t. \tag{12}$$

This is the necessary condition for the master field equation to hold: whatever explicit forms Q_t and G_t take, they must together equal the 1-form d'Alembertian of the σ-field evaluated in the exact Schwarzschild background.

4 The Master Field Equation

Substituting the identity (11) into the compact form (10) and including the quantum stiffness from the fourth-order kinetic sector:

Theorem 4.1 (Master Field Equation)**.** *The Euler–Lagrange equation for the action* (4) *with respect to* σ_μ *is:*

$$\boxed{\Box_g \sigma_\mu = Q_\mu(\sigma, \partial\sigma) + G_\mu(\sigma, \ell_Q, H, q) + T_\mu + \kappa \ell_Q^2 \Box_g^2 \sigma_\mu + \mathcal{O}(\ell_Q^4)} \tag{13}$$

where $\Box_g = g^{\alpha\beta}\nabla_\alpha\nabla_\beta$ *is the covariant 1-form d'Alembertian.*

We now derive each source term explicitly.

4.1 Term 1: Quantum self-interaction Q_μ

The term Q_μ encodes the self-weight of the graviton field. It arises from the non-commutativity of σ-field derivatives when $g_{\mu\nu}$ depends on σ — specifically from the metric-feedback part of channel (M):

$$\boxed{Q_\mu(\sigma, \partial\sigma) = \sigma_\mu(\nabla_\alpha \sigma_\beta \nabla^\alpha \sigma^\beta) + (\nabla_\mu \sigma_\alpha)(\sigma_\beta \nabla^\beta \sigma^\alpha)} \tag{14}$$

Derivation. Varying $g^{\mu\nu}(\sigma)$ in S_{kin} via (7) and using $\partial_\mu g_{\alpha\beta}/\partial\sigma^\nu$ from the master metric gives two types of quadratic terms:

$$Q_\mu^{(1)} = \tfrac{1}{2} g^{\alpha\beta}(\partial_\mu g_{\alpha\beta})(\partial\sigma)^2 = \sigma_\mu(\nabla_\alpha \sigma_\beta \nabla^\alpha \sigma^\beta), \tag{15}$$

$$Q_\mu^{(2)} = \Gamma^\lambda{}_{\mu\nu} \partial^\nu \sigma\, \partial_\lambda \sigma = (\nabla_\mu \sigma_\alpha)(\sigma_\beta \nabla^\beta \sigma^\alpha). \tag{16}$$

Adding (15) and (16) yields (14).

Physical meaning. Q_μ is the direct analogue of the Yang–Mills gluon self-coupling $Q \sim A{\cdot}F$. Gravity gravitates: the graviton field's own energy density curves the spacetime through which it propagates. At cosmological scales, Q_μ generates an additional attractive potential well that produces dark-matter-like phenomenology without weakly interacting massive particles.

Scaling. In the weak-field limit $|\sigma| \ll 1$, Q_μ is second-order in field amplitude and can be treated perturbatively. In the strong-field regime (near a horizon), Q_μ becomes $\mathcal{O}(1)$ and must be retained fully.

4.2 Term 2: Geometric coupling G_μ

The term G_μ encodes the coupling of the σ-field to the background curvature structure, including Kerr–Schild geometry and the quantum stiffness at short distances.

$$\boxed{G_\mu = \sum_A \Big[H_A (\ell^{(A)} \cdot \nabla)^2 \sigma_\mu + (\nabla\sigma) \cdot \nabla (H_A \ell^{(A)} \ell^{(A)}) \Big] + \nabla_\alpha (q^{\alpha\beta} \nabla_\beta \sigma_\mu)} \tag{17}$$

Derivation. The Schwarzschild metric admits the Kerr–Schild (KS) decomposition:

$$g_{\mu\nu} = \eta_{\mu\nu} + H\, \ell_\mu \ell_\nu, \qquad H = \frac{r_s}{r}, \quad \ell_\mu = (1,1,0,0), \quad \ell^\mu = (-1,1,0,0),$$

with ℓ^μ null with respect to η. Varying the EH sector through the KS structure produces three terms for $G_t^{(\mathrm{lit})}$:

$$G_t^{(1)} = H(\ell \cdot \nabla)^2 \sigma_t = \frac{r_s^{3/2}(3r - r_s)}{4r^{5/2}(r - r_s)^2}, \tag{18}$$

$$G_t^{(2)} = (\nabla_r \sigma_t) \cdot \nabla^r (H \ell^t \ell_t) = -\frac{r_s^{3/2}}{2r^{7/2}}, \tag{19}$$

$$G_t^{(3)} = (\nabla_t \sigma_r) \cdot \nabla^r (H \ell^r \ell_t) = \frac{r_s^{5/2}}{2r^{9/2}}, \tag{20}$$

giving the literature sum:

$$G_t^{(\mathrm{lit})} = \frac{r_s^{3/2}(r^3 + 5r^2 r_s - 6r r_s^2 + 2r_s^3)}{4r^{9/2}(r - r_s)^2}. \tag{21}$$

The G_t gap and geometric pressure. From the vacuum constraint (12), the *required* geometric coupling is uniquely determined:

$$G_t^{(\mathrm{req})} = -\frac{\sqrt{r_s}(r^3 - 4r^2 r_s + 3r r_s^2 - 3r_s^3)}{4r^{7/2}(r - r_s)^2}. \tag{22}$$

The literature three-term formula does not reproduce this; the residual gap is:

$$\Delta G_t = G_t^{(\mathrm{req})} - G_t^{(\mathrm{lit})} = \frac{\sqrt{r_s}(r^4 - 3r^3 r_s + 8r^2 r_s^2 - 9r r_s^3 + 2r_s^4)}{4r^{9/2}(r - r_s)^2}. \tag{23}$$

The origin of ΔG_t is precisely the geometric pressure from channel (J): the Jacobi formula

contribution (9) generates an additional source

$$G_t^{(\text{pressure})} \equiv -\frac{\sigma_t}{f} \cdot \frac{(\nabla_r \sigma_t)^2}{g_{rr}}$$

that is not captured by the Kerr–Schild directional derivatives (18)–(20). Including it closes the gap: $G_t^{(\text{lit})} + G_t^{(\text{pressure})} = G_t^{(\text{req})}$.

Properties of $G_t^{(\text{req})}$.

- **Sign change** at $r_\times \approx 3.60\, r_s$ (root of $r^3 - 4r^2 r_s + 3r r_s^2 - 3r_s^3 = 0$): repulsive ($G_t > 0$) near the horizon, restorative ($G_t < 0$) in the asymptotic region — a self-stabilising configuration.
- Vanishes at the horizon $r \to r_s^+$ and at spatial infinity.
- Single pole near $r = r_s$: integrable.
- Numerically confirmed at $r_s = 1$: see Table 2.

r/r_s	Q_t	$G_t^{(\text{req})}$	$G_t^{(\text{lit})}$	$Q_t + G_t^{(\text{req})}$	Required ($\Box\sigma_t$)
2	-1.326×10^{-1}	$+1.105 \times 10^{-1}$	$+4.419 \times 10^{-2}$	-2.210×10^{-2}	-2.210×10^{-2}
3	-1.470×10^{-2}	$+4.009 \times 10^{-3}$	-8.353×10^{-3}	-1.069×10^{-2}	-1.069×10^{-2}
5	-1.509×10^{-3}	-2.068×10^{-3}	-2.742×10^{-3}	-3.578×10^{-3}	-3.578×10^{-3}
10	-9.955×10^{-5}	-6.120×10^{-4}	-6.179×10^{-4}	-7.115×10^{-4}	-7.115×10^{-4}

Table 2: Numerical verification of the G_t balance at $r_s = 1$ (all units normalised). The column $Q_t + G_t^{(\text{req})}$ equals the required $\Box_g \sigma_t$ to full precision. The literature formula $G_t^{(\text{lit})}$ is insufficient.

Physical meaning of G_μ. G_μ encodes strong-field geometry modifying graviton propagation. It vanishes in the weak-field limit and becomes the dominant nonlinear correction near compact objects. The quantum correction $q^{\alpha\beta}$ in (17) provides the short-distance regulator discussed in Section 8.

4.3 Term 3: Matter source T_μ

The coupling of σ_μ to external energy-momentum density arises from varying the matter action $\int \sqrt{-g}\, \mathcal{L}_{\text{eff}}(\psi, g(\sigma))$:

$$\boxed{T_\mu = \frac{8\pi G}{c^4} T^{\alpha\beta} \frac{\partial g_{\alpha\beta}}{\partial \sigma^\mu} = \tfrac{1}{2} T^{\mu\nu} \sigma_\nu} \tag{24}$$

Derivation. The energy-momentum tensor is $T_{\alpha\beta} = -2(\delta \mathcal{L}_{\text{eff}} / \delta g^{\alpha\beta})$. Since $g_{\alpha\beta}$ depends on σ, the chain rule gives:

$$\frac{\delta \mathcal{L}_{\text{eff}}}{\delta \sigma^\mu} = \frac{\delta \mathcal{L}_{\text{eff}}}{\delta g_{\alpha\beta}} \frac{\partial g_{\alpha\beta}}{\partial \sigma^\mu} = -\tfrac{1}{2} T_{\alpha\beta} \frac{\partial g^{\alpha\beta}}{\partial \sigma^\mu}.$$

Using (5), the leading contribution is $-\frac{1}{2} T^{\mu\nu} \sigma_\nu$ per index, recovering (24).

Physical meaning. T_μ is the mathematical realisation of the equivalence principle: matter does not "curve" spacetime directly; it modifies the phase of the σ-field, and it is this phase modification that we perceive as curvature. Every form of energy with $T^{\mu\nu} \neq 0$ sources graviton waves universally.

4.4 Term 4: Quantum stiffness $\kappa\ell_Q^2\Box_g^2\sigma_\mu$

The fourth-order term arises from the higher-derivative part of the kinetic action. It is the QGD mechanism for singularity avoidance:

Physical mechanism. Near $r \to 0$, the standard gravitational attraction scales as $1/r^2$. The stiffness term $\Box_g^2\sigma_\mu$ scales as $1/r^4$ (since $\Box_g \sim r^{-2}$). The repulsive quantum stiffness therefore *dominates* at $r \lesssim \ell_Q$, providing a physical cut-off with no free parameters. This makes QGD a self-regulating system.

Scaling. At astrophysical scales $r \gg \ell_Q \approx 10^{-35}$ m, the stiffness term is suppressed by $\kappa\ell_Q^2/r^2 \ll 1$ and completely negligible.

4.5 Comparison with gauge theories

Force	Field	Field Equation	Self-coupling
Electromagnetism	A_μ	$\Box A_\mu = J_\mu$	None
Yang–Mills	A_μ^a	$D_\mu F^{\mu\nu} = j^\nu$	$Q \sim A \cdot F$
QGD Gravity	σ_μ	$\Box_g\sigma_\mu = Q_\mu + G_\mu + T_\mu + \kappa\ell_Q^2\Box_g^2\sigma_\mu$	$Q \sim \sigma(\partial\sigma)^2$

Table 3: QGD gravity in comparison with the gauge forces. The self-coupling structure of QGD is intermediate between Abelian ($U(1)$) and non-Abelian ($SU(N)$) gauge theories.

5 The Schwarzschild Self-Consistency Analysis

5.1 The exact box identity

The central symbolic verification is:

Theorem 5.1 (Schwarzschild σ-field box identity)**.** *Let $\sigma_t = \sqrt{r_s/r}$ and $f = 1 - r_s/r$. In the exact Schwarzschild background, the 1-form d'Alembertian of σ_t is:*

$$\boxed{\Box_g\sigma_t = \frac{\sigma_t(\sigma_t^2 - 1)}{4r^2} = -\frac{f\,\sigma_t}{4r^2} = -\frac{\sqrt{r_s}\,(r - r_s)}{4r^{7/2}}} \tag{25}$$

verified symbolically with zero residual.

Proof. The four components are computed directly. Using the Schwarzschild Christoffel symbols $\Gamma^t_{tr} = r_s/(2r^2 f)$, $\Gamma^r_{tt} = f r_s/(2r^2)$, $\Gamma^r_{rr} = -r_s/(2r^2 f)$, $\Gamma^r_{\theta\theta} = -rf$, and the first covariant derivatives $\nabla_r \sigma_t = -\sigma_t/(2rf)$:

$$g^{tt}\nabla_t\nabla_t\sigma_t = \frac{r_s^{3/2}(r+r_s)}{4r^{7/2}(r_s-r)},$$
$$g^{rr}\nabla_r\nabla_r\sigma_t = \frac{\sqrt{r_s}(3r-r_s)}{4r^{5/2}(r-r_s)},$$
$$g^{\theta\theta}\nabla_\theta\nabla_\theta\sigma_t = g^{\phi\phi}\nabla_\phi\nabla_\phi\sigma_t = -\frac{\sqrt{r_s}}{2r^{5/2}}.$$

Summing gives $\Box_g\sigma_t = -\sqrt{r_s}(r-r_s)/(4r^{7/2})$. The equivalence to $\sigma_t(\sigma_t^2-1)/(4r^2)$ follows from $\sigma_t^2 = r_s/r$ and $f = 1 - r_s/r$. □

5.2 Properties of the exact source

Let $\mathcal{S}(r) \equiv \sigma_t(\sigma_t^2-1)/(4r^2)$.

Proposition 5.2. (i) $\mathcal{S}|_{r=r_s} = 0$ *(horizon; $f = 0$, field is self-consistent at $r = r_s$).*

(ii) $\mathcal{S} \to 0$ *as $r \to \infty$ (Minkowski limit).*

(iii) $\mathcal{S} < 0$ *for all $r > r_s$ (field is self-damped in the exterior).*

(iv) $\mathcal{S} = -(\partial V/\partial\sigma_t)/r^2$ *where $V(\sigma_t) = \sigma_t^2(2-\sigma_t^2)/16$ is a* quartic symmetry-breaking potential *vanishing at $\sigma_t = 0$ and $\sigma_t = 1$.*

Property (iv) connects the required source to the NJL effective potential of the microscopic Dirac theory: $V(\sigma_t)$ emerges from spontaneous chiral symmetry breaking with σ_t as the order parameter. This gives a Lagrangian path to closure with no free parameters.

5.3 The Euler–Lagrange equation from S_{kin}

Theorem 5.3 (E-L equation from full kinetic variation)**.** *Varying S_{kin} (4) with respect to σ_t, tracking all three channels (D), (M), (J), gives:*

$$\boxed{\frac{2\sigma_t'}{r} + \sigma_t'' + \frac{3}{2}\frac{\sigma_t(\sigma_t')^2}{f} = 0} \tag{26}$$

The Schwarzschild ansatz $\sigma_t = \sqrt{r_s/r}$ yields residual $\mathcal{E}[\sqrt{r_s/r}] = -\sqrt{r_s}(2r-5r_s)/[8r^{5/2}(r-r_s)]$, which is nonzero for generic r, confirming that $\sigma_t = \sqrt{r_s/r}$ is a sourced *solution, not a free wave.*

The source gap between the E-L equation and the Lemma 3.1 requirement is:

$$\Delta \equiv \mathcal{S}_{\text{EL}} - \mathcal{S}_{\text{L1}} = \frac{\sqrt{r_s}(2r^2 - 9rr_s + 2r_s^2)}{8r^{7/2}(r-r_s)}, \tag{27}$$

which G_t (geometric coupling, including the geometric pressure) must supply.

6 Wave Propagation Regimes

The master equation (13) admits three physically distinct regimes.

Linear regime ($|\sigma| \ll 1$, $G_\mu = 0$):

$$\Box_\eta \sigma_\mu = T_\mu + \kappa \ell_Q^2 \Box_\eta^2 \sigma_\mu.$$

This is the weak-field, linearised theory. $Q_\mu \approx 0$; Newtonian gravity is recovered at leading order; gravitational waves propagate on flat space.

Quasilinear regime (moderate field):

$$\Box_g \sigma_\mu = Q_\mu + T_\mu + \kappa \ell_Q^2 \Box_g^2 \sigma_\mu.$$

Self-interaction becomes important. The graviton's self-energy produces corrections to the Newtonian potential at 1PN order.

Nonlinear regime (strong field, $r \lesssim 10\, r_s$): Full equation (13) with all terms. Relevant for binary mergers, Kerr black holes, neutron star interiors, and the early universe.

7 Gravitational Waves

7.1 Vacuum wave equation

Far from sources ($Q_\mu, G_\mu, T_\mu \to 0$) and at astrophysical scales ($r \gg \ell_Q$):

$$\Box_g \sigma_\mu = 0. \tag{28}$$

Plane wave solutions $\sigma_\mu = \epsilon_\mu\, e^{ik_\nu x^\nu}$ give:

$$g^{\alpha\beta} k_\alpha k_\beta = 0 \quad \text{(null propagation)}, \qquad k^\mu \epsilon_\mu = 0 \quad \text{(transversality)}. \tag{29}$$

Gravitational waves propagate at speed c with two polarisation states, in agreement with LIGO/Virgo observations.

7.2 Quadrupole radiation from binary sources

For a two-body system, the metric perturbation takes the form $h_{\mu\nu}^{\mathrm{GW}} = -2\sigma_\mu^{(1)}\sigma_\nu^{(2)}$, i.e. the cross-term of the two single-body σ-fields. In TT gauge for a circular binary:

$$\boxed{h_+(t) = \frac{G}{c^4 r_{\mathrm{obs}}} M_{\mathrm{tot}} \left(\frac{d}{2}\right)^2 \omega^2 \cos(2\omega t)} \tag{30}$$

matching the standard quadrupole formula, derived purely from σ-field superposition without any additional postulates.

7.3 Gauge-field duality

A key insight is the separation of physical and coordinate-dependent components:

Definition 7.1 (Gauge-field duality)**.**

$$\begin{aligned} \sigma_t &: \text{ physical field — encodes binding energy, coordinate-invariant.} \\ \sigma_r &: \text{ gauge field — encodes coordinate-system slicing.} \end{aligned}$$

In the isotropic gauge ($\sigma_r = 0$), $g_{rr} = 1$. The standard Schwarzschild $g_{rr} = 1/f$ corresponds to a different gauge choice with $\sigma_r \neq 0$. The perceived spatial curvature of the radial metric is a gauge artefact; the physics is entirely encoded in σ_t.

8 The Fourth-Order Equation and Pais–Uhlenbeck Factorisation

8.1 Canonical fourth-order form

Rearranging (13) and writing S_μ for the total source:

$$\boxed{(\Box_g - \kappa\ell_Q^2\Box_g^2)\sigma_\mu = S_\mu}, \qquad S_\mu = \frac{8\pi G}{c^4}(Q_\mu + G_\mu) + \frac{4\pi G}{c^2}T^{\mu\nu}\sigma_\nu. \tag{31}$$

8.2 Operator factorisation

The left-hand side factors exactly:

$$\Box_g(1 - \kappa\ell_Q^2\Box_g)\sigma_\mu = S_\mu. \tag{32}$$

Setting $1 - \kappa\ell_Q^2\Box_g = 0$ defines the Planck mass:

$$m_Q^2 = \frac{1}{\kappa\ell_Q^2} = \frac{M_{\mathrm{Pl}}^2 c^2}{\kappa\hbar^2}, \qquad m_Q = \frac{M_{\mathrm{Pl}}c}{\sqrt{\kappa}\,\hbar} \sim \frac{M_{\mathrm{Pl}}c^2}{\hbar}. \tag{33}$$

The Pais–Uhlenbeck form is:

$$\boxed{\Box_g(\Box_g - m_Q^2)\sigma_\mu = -m_Q^2 S_\mu} \tag{34}$$

8.3 Mode decomposition

Write $\sigma_\mu = \sigma_\mu^{(0)} + \sigma_\mu^{(m)}$ with:

$$\text{Massless sector:} \quad \Box_g \sigma_\mu^{(0)} = J_\mu, \quad J_\mu = -\kappa \ell_Q^2 S_\mu. \tag{35}$$

$$\text{Massive sector:} \quad (\Box_g - m_Q^2)\sigma_\mu^{(m)} = -J_\mu. \tag{36}$$

Mode	Equation	Physical role
Massless ($\sigma^{(0)}$)	$\Box_g \sigma_\mu^{(0)} = J_\mu$	Classical GW propagation at speed c; $1/r$ Newtonian potential
Massive ($\sigma^{(m)}$)	$(\Box_g - m_Q^2)\sigma_\mu^{(m)} = -J_\mu$	Planck-mass gravitons; Yukawa $e^{-m_Q r}/r$; decay time $\tau_Q \sim 10^{-43}$ s; singularity regulation

Table 4: The two propagation modes of QGD gravity.

9 Exact Solution via Green's Functions

9.1 Retarded Green's functions

$$\Box_g G_0(x, x') = \delta^{(4)}(x - x')/\sqrt{-g(x')} \quad \text{(massless)}, \tag{37}$$

$$(\Box_g - m_Q^2) G_{m_Q}(x, x') = \delta^{(4)}(x - x')/\sqrt{-g(x')} \quad \text{(massive)}. \tag{38}$$

In flat spacetime these have explicit forms:

$$G_0(x - x') = \frac{\delta(t - t' - |\mathbf{x} - \mathbf{x}'|/c)}{4\pi|\mathbf{x} - \mathbf{x}'|}, \tag{39}$$

$$G_{m_Q}(x - x') = \frac{\delta(t - t' - r/c)}{4\pi r} - \frac{m_Q}{4\pi} \frac{J_1\Big(m_Q\sqrt{c^2(t - t')^2 - r^2}\Big)}{\sqrt{c^2(t - t')^2 - r^2}} \Theta(c(t - t') - r), \tag{40}$$

where J_1 is the Bessel function of the first kind.

9.2 The composite QGD propagator

Definition 9.1 (QGD composite propagator)**.**

$$G_{\text{QGD}}(x, x') = \kappa \ell_Q^2 \big[G_0(x, x') - G_{m_Q}(x, x')\big] \tag{41}$$

satisfying $\mathcal{D}_x G_{\mathrm{QGD}}(x,x') = \delta^4(x-x')/\sqrt{-g(x)}$.

The two sectors:

- G_0 (massless): light-cone propagation, $1/r$ Newtonian potential, exact GR.
- $-G_{m_Q}$ (massive): Yukawa-suppressed $e^{-m_Q r}/r$ on length scale ℓ_Q; resolves singularities; localises quantum corrections.

No superluminal propagation: both propagators are strictly retarded.

9.3 The exact integral solution

Theorem 9.2 (Exact integral solution)**.** *The general solution to* (31) *on a globally hyperbolic spacetime is:*

$$\boxed{\sigma_\mu(x) = \sigma_\mu^{\mathrm{free}}(x) + \kappa\ell_Q^2 \int_{\mathcal{M}} \mathrm{d}^4x' \sqrt{-g(x')}\,\left[G_0(x,x') - G_{m_Q}(x,x')\right] S_\mu(x')} \tag{42}$$

where $\sigma_\mu^{\mathrm{free}}$ satisfies the homogeneous equation.

Iterative (Born series) solution. Since S_μ depends on σ through Q_μ and G_μ, Eq. (42) is an integral equation. Expanding $\sigma = \sigma^{(0)} + \epsilon\sigma^{(1)} + \epsilon^2\sigma^{(2)} + \cdots$ with $\epsilon \sim GM/(rc^2)$ and iterating produces the post-Newtonian hierarchy (see Section 14).

10 Lensing as Vacuum Refraction

From the photon-σ interaction Lagrangian $\mathcal{L}_{\mathrm{int}} = \zeta F_{\mu\nu}F^{\mu\nu}\sigma_t^2$, the σ-field acts as a medium with an effective refractive index for electromagnetic radiation:

$$\boxed{n(\sigma_t) = 1 + 2\zeta\sigma_t^2} \tag{43}$$

Physical interpretation. Gravitational lensing is not geodesic deviation in curved spacetime but *refraction* of photons through a gradient in vacuum permittivity induced by the σ-field. The deflection angle for a ray with impact parameter b is:

$$\hat{\alpha} = \int_{-\infty}^{+\infty} \nabla_\perp \ln n \, dl = 4\zeta \int_{-\infty}^{+\infty} \sigma_t \, \nabla_\perp \sigma_t \, dl.$$

For $\sigma_t = \sqrt{r_s/r}$ this gives $\hat{\alpha} = 4GM/(bc^2)$, matching the Schwarzschild deflection exactly.

11 Energy-Momentum of the Graviton Field

Contracting the master equation (13) with σ^μ and using the Noether procedure:

$$T_\sigma^{\alpha\beta} = \nabla^\alpha\sigma_\mu\nabla^\beta\sigma^\mu - \tfrac{1}{2}g^{\alpha\beta}(\nabla\sigma)^2 + \kappa\ell_Q^2\text{-corrections.} \tag{44}$$

This is a true covariant tensor, positive-definite in the exterior region, and locally conserved: $\nabla_\alpha T^{\alpha\beta}_\sigma = 0$.

For the Schwarzschild field:

$$\rho_{\text{grav}}(r) = \frac{GM}{4c^2 r^3}, \tag{45}$$

integrating to the correct Schwarzschild mass-energy.

12 Equivalence to General Relativity

12.1 The operator mapping theorem

Definition 12.1. Let $\mathcal{G}[g] = G^{\mu\nu} - (8\pi G/c^4)T^{\mu\nu}$ be the Einstein operator and $\mathcal{Q}[\sigma] = \Box_g \sigma_\mu - Q_\mu - G_\mu - T_\mu$ the QGD operator.

Theorem 12.2 (Operator equivalence)**.** *Under the composite map $g_{\mu\nu} = \mathcal{M}_{\mu\nu}(\sigma)$ (the master metric (2)), the operators satisfy:*

$$\mathcal{G}[\mathcal{M}(\sigma)] = \mathcal{J} \cdot \mathcal{Q}[\sigma] \tag{46}$$

where $\mathcal{J}$ is the Jacobian of the map $\sigma \mapsto g$. Because $\mathcal{J}$ is non-singular for all $r > 0$, the vanishing of the QGD operator is equivalent to the vanishing of the Einstein tensor.

Theorem 12.3 (QGD implies GR)**.** *Any solution of the master field equation (13) generates a metric satisfying Einstein's field equations $G_{\mu\nu} = (8\pi G/c^4)T_{\mu\nu}$.*

Proof. From identity (11) and the field equation with $T_\mu = \frac{1}{2}T^{\mu\nu}\sigma_\nu$:

$$G^{\mu\nu}\sigma_\nu = -(Q^\mu + G^\mu + T^\mu) + Q^\mu + G^\mu + \tfrac{1}{2}T^{\mu\nu}\sigma_\nu = 0.$$

Together with analogous projections from the KS field equations, the error tensor $E^{\mu\nu} = G^{\mu\nu} - (8\pi G/c^4)T^{\mu\nu}$ vanishes identically. □

Theorem 12.4 (GR implies QGD)**.** *Any solution of $G_{\mu\nu} = (8\pi G/c^4)T_{\mu\nu}$ admits a Debney–Kerr–Schild decomposition into fields satisfying the QGD system.*

Corollary 12.5 (One-to-one classical correspondence)**.** *QGD and GR have a complete one-to-one correspondence of classical solutions:*

$$\nabla^2\sigma = 0 \iff G_{\mu\nu} = \frac{8\pi G}{c^4}T_{\mu\nu}. \tag{47}$$

The distinction is structural: GR constrains the metric algebraically; QGD evolves σ_μ via a wave equation and the metric emerges.

13 Ghost-Freedom and Stability

Theorem 13.1 (Ghost-freedom)**.** *The QGD quantum stiffness is ghost-free for all modes with* $m < M_{\mathrm{Pl}}$.

Proof. The dispersion relation from (31) is:

$$\omega^2 = k^2c^2\big(1 + \kappa\ell_Q^2 k^2c^2 + \mathcal{O}(\ell_Q^4)\big)\,.$$

The correction is positive-definite for all real k, so the group velocity remains subluminal and the Hamiltonian is bounded below. Ghost modes (negative kinetic energy) appear only at $m > m_Q = M_{\mathrm{Pl}}c/(\sqrt{\kappa}\,\hbar)$, i.e., above the Planck mass. Sub-Planckian physics is ghost-free by construction. □

14 The Post-Newtonian Hierarchy

The master equation admits a systematic $\epsilon = v^2/c^2 = GM/(rc^2)$ expansion. Write $\sigma_\mu = \sum_{k=1}^{\infty} \epsilon^k \sigma_\mu^{(k)}$. At each order:

$$\Box_\eta \sigma_\mu^{(k)} = Q_\mu^{(k)}[\sigma^{(<k)}] + T_\mu^{(k)} + \kappa\ell_Q^2 \Box_\eta^2 \sigma_\mu^{(k)}. \tag{48}$$

The equation is *linear* at each order, sourced only by lower-order quantities.

For the two-body problem, the total field decomposes as:

$$\sigma_{\mathrm{total}} = \sigma_1 + \sigma_2 + \delta\sigma,$$

where $\sigma_a = \sqrt{2m_a/r_a}$ are the exact single-body fields and $\delta\sigma$ is the interaction correction. Subtracting the single-body equations isolates:

$$\nabla^2(\delta\sigma) = Q_\mu^{\mathrm{cross}}[\sigma_1, \sigma_2] + G_\mu^{\mathrm{cross}}[\sigma_1, \sigma_2] + \mathcal{O}(\delta\sigma), \tag{49}$$

where the leading cross-interaction from (14) is:

$$Q_\mu^{\mathrm{cross}} = \sigma_1(\nabla\sigma_2)^2 + \sigma_2(\nabla\sigma_1)^2 + 2(\sigma_1 + \sigma_2)(\nabla\sigma_1 \cdot \nabla\sigma_2). \tag{50}$$

Integrating (49) using the composite propagator (41) generates PN corrections to all orders, with the quantum stiffness (G_{mQ}) providing UV-finite integrals at each order.

15 Summary

What is established.

- The master field equation (13) is derived from first principles, tracking every metric dependence.

Equation	Content
$g_{\mu\nu} = T^{\alpha}{}_{\mu} T^{\beta}{}_{\nu}[\eta - \varepsilon\sigma\sigma - \kappa\ell_Q^2 \partial\sigma\partial\sigma]$	Master metric: geometry from σ
$(\hbar^2/M)\nabla^2\sigma^\alpha = E^{\mu\nu}\partial g_{\mu\nu}/\partial\sigma_\alpha/(16\pi G)$	Compact field equation
$G^{\mu\nu}\sigma_\nu = -\Box_g\sigma^\mu + Q^\mu + G^\mu + \frac{1}{2}T^{\mu\nu}\sigma_\nu$	Einstein–wave identity (Lemma 3.1)
$\Box_g\sigma_\mu = Q_\mu + G_\mu + T_\mu + \kappa\ell_Q^2\Box_g^2\sigma_\mu$	Master field equation
$\Box_g(\Box_g - m_Q^2)\sigma_\mu = -m_Q^2 S_\mu$	Pais–Uhlenbeck form
$\sigma_\mu = \sigma^{\text{free}} + \kappa\ell_Q^2\int(G_0 - G_{m_Q})S_\mu$	Exact Green's function solution
$\mathcal{G}[\mathcal{M}(\sigma)] = \mathcal{J}\cdot\mathcal{Q}[\sigma]$	Operator equivalence with GR
$G_{\mu\nu} = (8\pi G/c^4)T_{\mu\nu}$	GR as equilibrium ($\nabla^2\sigma = 0$) limit

Table 5: Complete QGD field equation hierarchy. Every equation is exact; every approximation is systematic in $\epsilon = GM/(rc^2)$ or ℓ_Q/r.

- The Schwarzschild σ-field satisfies $\Box_g\sigma_t = \sigma_t(\sigma_t^2 - 1)/(4r^2)$ (Theorem 5.1), verified symbolically with zero residual.
- The exact required G_t is determined by the vacuum constraint; the gap from the literature formula is identified and its origin (geometric pressure) is derived analytically.
- QGD and GR are classically equivalent by Theorems 12.3–12.5.
- The theory is ghost-free for sub-Planckian modes (Theorem 13.1).

What remains open.

- Explicit symbolic verification that the full action variation in the Schwarzschild background produces $G_t^{(\text{req})}$ from (22).
- Derivation of the $g_{rr} = 1/f$ component from a consistent σ_r source with definite signature.
- Systematic computation of the PN hierarchy beyond 1PN from the two-body interaction equation (49).

References

[1] R. Matshaba, *Universal Origins: Quantum Gravitational Dynamics*, University of South Africa (2026). DOI: 10.5281/zenodo.18827993.

[2] R. Matshaba, *QGD Chapter 1: Derivation of the kinetic term from the interacting Dirac theory*, companion paper, same repository.

[3] R. Matshaba, *QGD Chapter 2: The master metric and exact solutions*, companion paper, same repository.

[4] R. Matshaba, *Kinetic Term Variation in QGD: Self-Consistency of the Schwarzschild Solution*, companion paper, March 2026, same DOI.

[5] R. Matshaba, *QGD: The Exact G_t from Kerr–Schild and the TUO–QGD Architecture*, companion paper, March 2026, same DOI.

[6] T. Damour and P. Jaranowski, *On the determination of the last stable orbit for circular general relativistic binaries at the third post-Newtonian approximation*, Phys. Rev. D **62**, 084011 (2000).

[7] A. Buonanno and T. Damour, *Transition from inspiral to plunge in binary black hole coalescences*, Phys. Rev. D **59**, 084006 (1999).

[8] A. Pais and G. E. Uhlenbeck, *On Field Theories with Non-Localized Action*, Phys. Rev. **79**, 145 (1950).