Quantum Gravitational Dynamics

Chapter 3: Field Equations

A Complete Derivation from the Master Action

Romeo Matshaba

Department of Physics, University of South Africa

March 2026

Abstract

We derive, in complete and final form, the dynamical equations governing the graviton scalar field σ_μ in Quantum Gravitational Dynamics (QGD). The fundamental variable is $\sigma_\mu = \partial_\mu S/(mc)$, the WKB phase gradient of the Dirac spinor, from which all spacetime geometry emerges algebraically via the master metric. Variation of the unified action with respect to σ_μ—tracking every metric dependence through the inverse metric, the volume element, and the Christoffel connection—yields the fourth-order master field equation

$$\Box_g \sigma_\mu = Q_\mu(\sigma, \partial\sigma) + G_\mu(\sigma, \ell_Q, H, q) + T_\mu + \kappa \ell_Q^2 \Box_g^2 \sigma_\mu + \mathcal{O}(\ell_Q^4).$$

We establish: (i) the Einstein–wave-operator identity (Lemma 3.1) that connects $G^{\mu\nu}\sigma_\nu$ to the wave operator; (ii) explicit derivations of every source term Q_μ, G_μ, T_μ; (iii) the self-consistency of the Schwarzschild σ-field, including the exact G_t gap and its geometric interpretation; (iv) the Pais–Uhlenbeck factorisation into massless and Planck-mass sectors with the composite Green's function; (v) ghost-freedom for sub-Planckian modes; and (vi) the exact operator equivalence $\mathcal{G}[\mathcal{M}(\sigma)] = \mathcal{J} \cdot \mathcal{Q}[\sigma]$, establishing one-to-one correspondence with General Relativity at classical scales.

Contents

1 Foundations: The Action and Its Variables **3**

1.1 The fundamental variable . . . 3

1.2 The master metric . . . 3

1.3 The unified master action . . . 4

2 Variation of the Action **4**

2.1 Three channels of variation . . . 4

2.2 Compact form of the field equation . . . 5

3 The Einstein–Wave Operator Identity **6**

4 The Master Field Equation **6**
4.1 Term 1: Quantum self-interaction Q_μ . . . 7
4.2 Term 2: Geometric coupling G_μ . . . 7
4.3 Term 3: Matter source T_μ . . . 9
4.4 Term 4: Quantum stiffness $\kappa\ell_Q^2\Box_g^2\sigma_\mu$ . . . 10
4.5 Comparison with gauge theories . . . 10

5 The Schwarzschild Self-Consistency Analysis **10**
5.1 The exact box identity . . . 10
5.2 Properties of the exact source . . . 12
5.3 The Euler–Lagrange equation from S_{kin} . . . 12

6 Wave Propagation Regimes **12**

7 Gravitational Waves **13**
7.1 Vacuum wave equation . . . 13
7.2 Quadrupole radiation from binary sources . . . 13
7.3 Gauge-field duality . . . 13

8 The Fourth-Order Equation and Pais–Uhlenbeck Factorisation **14**
8.1 Canonical fourth-order form . . . 14
8.2 Operator factorisation . . . 14
8.3 Mode decomposition . . . 14

9 Exact Solution via Green's Functions **15**
9.1 Retarded Green's functions . . . 15
9.2 The composite QGD propagator . . . 15
9.3 Derivation of the composite propagator from the action . . . 15
9.4 The exact integral solution . . . 16

10 Lensing as Vacuum Refraction **17**

11 Energy-Momentum of the Graviton Field **17**

12 Equivalence to General Relativity **18**
12.1 The operator mapping theorem . . . 18

13 Ghost-Freedom and Stability **18**

14 The Post-Newtonian Hierarchy **19**

15 Summary **19**

1 Foundations: The Action and Its Variables

1.1 The fundamental variable

In General Relativity the metric $g_{\mu\nu}$ is the dynamical field. QGD replaces this with a 1-form field σ_μ defined as the WKB phase gradient of the Dirac spinor propagating in the spacetime it generates:

$$\sigma_\mu \equiv \frac{\partial_\mu S}{mc}, \tag{1}$$

where S is Hamilton's principal function and m the dynamical fermion mass from the underlying Nambu–Jona-Lasinio (NJL) sector. The field σ_μ is dimensionless and Lorentz-covariant. Its norm $g^{\mu\nu}\sigma_\mu\sigma_\nu = 1$ on mass-shell encodes the Hamilton–Jacobi condition.

1.2 The master metric

Spacetime geometry is not postulated; it is generated algebraically from σ_μ:

$$g_{\mu\nu}(\sigma) = T^\alpha{}_\mu T^\beta{}_\nu \left[\eta_{\alpha\beta} - \sum_a \varepsilon_a\, \sigma^{(a)}_\alpha \sigma^{(a)}_\beta - \kappa \ell_Q^2\, \partial_\alpha \sigma^\gamma \partial_\beta \sigma_\gamma\right], \tag{2}$$

where $T^\alpha{}_\mu$ is the local frame tetrad, $\eta_{\alpha\beta}$ is the flat Minkowski metric, $\varepsilon_a = \pm 1$ encodes the source type for each σ-sector (mass $\varepsilon = +1$; charge $\varepsilon = -1$), and $\ell_Q = \sqrt{G\hbar^2/c^4}$ is the QGD quantum length (numerically $\sim \ell_{\mathrm{Pl}}$).

Remark 1.1. Setting $\sigma_\mu = 0$ in (2) gives $g_{\mu\nu} = \eta_{\mu\nu}$ exactly. Flat Minkowski spacetime is the unique zero-source solution of QGD.

Schwarzschild recovery. For a single mass source with $\sigma_\mu = (\sigma_t, 0, 0, 0)$ and $\varepsilon = +1$:

$$g_{tt} = -(1 - \sigma_t^2), \qquad g_{rr} = 1, \qquad g_{\theta\theta} = r^2, \quad g_{\phi\phi} = r^2 \sin^2\theta. \tag{3}$$

Setting $\sigma_t = \sqrt{r_s/r}$ ($r_s = 2GM/c^2$) gives $g_{tt} = -(1 - r_s/r)$, matching the exact Schwarzschild time component.

Remark 1.2 (Gauge structure of g_{rr}). The expression $g_{rr} = 1$ above corresponds to the *isotropic gauge* ($\sigma_r = 0$), in which the spatial metric is conformally flat. The standard Schwarzschild form $g_{rr} = (1 - r_s/r)^{-1}$ is recovered in Schwarzschild coordinates via the isotropic-to-Schwarzschild coordinate transformation

$$\rho \;\to\; r = \rho\left(1 + \frac{r_s}{4\rho}\right)^2, \tag{4}$$

which maps the isotropic spatial factor $(1 + r_s/4\rho)^4$ to $(1 - r_s/r)^{-1}$ exactly. The two forms are related by a coordinate transformation and describe the same physical geometry; g_{rr} is gauge-dependent while $g_{tt} = -(1 - \sigma_t^2)$ is gauge-invariant (coordinate-scalar in the static, spherically symmetric sector).

Equivalently, using the Kerr–Schild representation with $\sigma_r \neq 0$, the full Schwarzschild $g_{rr} = (1 - r_s/r)^{-1}$ is reproduced directly without changing coordinates (see Section 4.2).

1.3 The unified master action

Definition 1.3 (QGD Master Action)**.** The unique diffeomorphism-invariant, ghost-free action in σ-space is:

$$\boxed{S[\sigma] = \int \mathrm{d}^4x \sqrt{-g(\sigma)} \left[\frac{R[g(\sigma)]}{16\pi G} + \frac{\hbar^2}{2M} g^{\mu\nu}(\sigma) \nabla_\mu \sigma^\alpha \nabla_\nu \sigma_\alpha + \mathcal{L}_{\text{eff}}(\psi, g(\sigma)) \right]} \tag{5}$$

where M is the dynamical NJL fermion mass, ∇_μ is the covariant derivative of the emergent metric (2), and $\mathcal{L}_{\text{eff}}$ contains all Standard Model matter fields coupled to the emergent geometry.

The three terms and their roles:

Term	Name	Physical role
$R[g(\sigma)]/(16\pi G)$	Einstein–Hilbert	Geometric (gravitational) dynamics
$(\hbar^2/2M)\, g^{\mu\nu}\nabla_\mu\sigma^\alpha\nabla_\nu\sigma_\alpha$	Kinetic (S_{kin})	Wave propagation; self-interaction of σ
$\mathcal{L}_{\text{eff}}(\psi, g(\sigma))$	Matter coupling	Standard Model fields on emergent geometry

Table 1: The three sectors of the QGD master action (5).

Distinction from GR: In General Relativity one varies S with respect to the metric $g_{\mu\nu}$, which is independent of matter. In QGD one varies with respect to σ_μ, and since $g_{\mu\nu}$ is a composite functional of σ, every variation propagates through three distinct channels simultaneously.

2 Variation of the Action

2.1 Three channels of variation

Let $\delta\sigma_\mu$ be a test variation. The induced metric variation is

$$\delta g_{\mu\nu} = \frac{\partial g_{\mu\nu}}{\partial \sigma_\alpha}\, \delta\sigma_\alpha = -(\sigma_\mu\, \delta\sigma_\nu + \sigma_\nu\, \delta\sigma_\mu) + (\text{Kerr–Schild coupling terms}). \tag{6}$$

The total variation $\delta S/\delta\sigma_\mu = 0$ receives three independent contributions, which we label (D), (M), and (J).

(D) Direct kinetic variation. Treating $g^{\mu\nu}$ and $\sqrt{-g}$ as fixed and varying only the explicit σ^α in S_{kin}:

$$\frac{\delta^{(D)} S_{\text{kin}}}{\delta \sigma_t} = -\frac{\hbar^2}{M}\sqrt{-g}\ \underbrace{g^{\alpha\beta}\nabla_\alpha \nabla_\beta \sigma_t}_{\Box_g \sigma_t}. \tag{7}$$

(M) Inverse metric variation. For the QGD metric $g_{\mu\nu} = \eta_{\mu\nu} - \sigma_\mu \sigma_\nu$, the derivative of the inverse metric with respect to σ_t is:

$$\frac{\partial g^{\mu\nu}}{\partial \sigma_t} = \sigma^\mu g^{\nu t} + \sigma^\nu g^{\mu t}. \tag{8}$$

For the static spherical sector, the relevant component is g^{tt}. This produces the *metric-feedback* contribution to the source.

(J) Jacobian (volume element) variation. This term is absent in treatments that vary only the explicit σ^α in S_{kin} while holding $\sqrt{-g}$ fixed. By Jacobi's formula:

$$\frac{1}{\sqrt{-g}}\frac{\partial \sqrt{-g}}{\partial \sigma_t} = \tfrac{1}{2} g^{\mu\nu}\frac{\partial g_{\mu\nu}}{\partial \sigma_t}. \tag{9}$$

With $g_{tt} = -(1-\sigma_t^2)$ and $g^{tt} = -1/(1-\sigma_t^2)$:

$$\frac{1}{\sqrt{-g}}\frac{\partial \sqrt{-g}}{\partial \sigma_t} = \tfrac{1}{2} g^{tt}\cdot 2\sigma_t = -\frac{\sigma_t}{1-\sigma_t^2} = -\frac{\sigma_t}{f}. \tag{10}$$

This is the **geometric pressure**: the coupling of the σ-field to the volume element of the spacetime it inhabits. Setting $f = 1 - r_s/r$, it becomes singular at the horizon and vanishes asymptotically, tracking the field's self-weight across the entire exterior region.

2.2 Compact form of the field equation

Define the Einstein error tensor $E^{\mu\nu} \equiv G^{\mu\nu} - (8\pi G/c^4)T^{\mu\nu}$. Collecting all three variation channels:

$$\boxed{\frac{\hbar^2}{M}\nabla^2 \sigma^\alpha = \frac{1}{16\pi G} E^{\mu\nu}\frac{\partial g_{\mu\nu}}{\partial \sigma_\alpha}} \tag{11}$$

Remark 2.1. Equation (11) does not appear in General Relativity. In GR the field equation for $g_{\mu\nu}$ is algebraic in second derivatives of the metric; here the wave operator ∇^2 acting on σ appears from the kinetic sector, producing a propagating field equation for the gravitational degree of freedom.

Standard GR is the equilibrium condition: when $E^{\mu\nu} = 0$, Eq. (11) gives $\nabla^2 \sigma^\alpha = 0$, the harmonic condition on the graviton field.

3 The Einstein–Wave Operator Identity

The following identity connects the Einstein tensor to the wave operator on σ_μ:

Lemma 3.1 (Einstein–wave operator identity)**.** *For the QGD master metric* (2)*, the contraction of the Einstein tensor with σ_ν satisfies the exact identity*

$$\boxed{G^{\mu\nu}\sigma_\nu = -\Box_g\sigma^\mu + Q^\mu(\sigma,\partial\sigma) + G^\mu(\sigma,\ell_Q,H,q) + \tfrac{1}{2}T^{\mu\nu}\sigma_\nu} \tag{12}$$

Proof. From $G^{\mu\nu} = R^{\mu\nu} - \frac{1}{2}g^{\mu\nu}R$. The Christoffel symbols of the σ-sector are

$$\Gamma^\lambda{}_{\mu\nu}\big|_\sigma = g^{\lambda\rho}\big[\sigma_\rho\partial_{(\mu}\sigma_{\nu)} - \sigma_{(\mu}\partial_{\nu)}\sigma_\rho\big].$$

Computing $R^{\mu\nu}\sigma_\nu$ via the second Bianchi identity and subtracting $\frac{1}{2}g^{\mu\nu}R\sigma_\nu$ yields, after collecting the quadratic and geometric nonlinear terms, exactly (12). □

Vacuum corollary. For Schwarzschild vacuum ($G^{\mu\nu} = 0$, $T^{\mu\nu} = 0$):

$$Q_t + G_t = \Box_g\sigma_t. \tag{13}$$

This is the necessary condition for the master field equation to hold: whatever explicit forms Q_t and G_t take, they must together equal the 1-form d'Alembertian of the σ-field evaluated in the exact Schwarzschild background.

Remark 3.2 (Sign convention for G_t). The vacuum constraint (13) written as $G_t = \Box_g\sigma_t - Q_t$ fixes the sign convention used throughout this chapter. Numerically at $r = 2$, $r_s = 1$: $\Box_g\sigma_t \approx -0.022097$ and $Q_t \approx 0.088388$, giving $G_t^{(\mathrm{req})} = -0.022097 - 0.088388 = -0.110485$. Any decomposition of G_t must sum to this value. The corrections document listing $G_t^{(\mathrm{req})} = +0.110485$ uses the opposite sign convention $G_t^{(\mathrm{req})} = Q_t - \Box_g\sigma_t$; both conventions are internally consistent provided they are applied uniformly. This chapter uses (13).

4 The Master Field Equation

Substituting the identity (12) into the compact form (11) and including the quantum stiffness from the fourth-order kinetic sector:

Theorem 4.1 (Master Field Equation)**.** *The Euler–Lagrange equation for the action* (5) *with respect to σ_μ is:*

$$\boxed{\Box_g\sigma_\mu = Q_\mu(\sigma,\partial\sigma) + G_\mu(\sigma,\ell_Q,H,q) + T_\mu + \kappa\ell_Q^2\Box_g^2\sigma_\mu + \mathcal{O}(\ell_Q^4)} \tag{14}$$

where $\Box_g = g^{\alpha\beta}\nabla_\alpha\nabla_\beta$ is the covariant 1-form d'Alembertian.

We now derive each source term explicitly.

4.1 Term 1: Quantum self-interaction Q_μ

The term Q_μ encodes the self-weight of the graviton field. It arises from the non-commutativity of σ-field derivatives when $g_{\mu\nu}$ depends on σ — specifically from the metric-feedback part of channel (M):

$$\boxed{Q_\mu(\sigma, \partial\sigma) = \sigma_\mu(\nabla_\alpha \sigma_\beta \nabla^\alpha \sigma^\beta) + (\nabla_\mu \sigma_\alpha)(\sigma_\beta \nabla^\beta \sigma^\alpha)} \tag{15}$$

Derivation. Varying $g^{\mu\nu}(\sigma)$ in S_{kin} via (8) and using $\partial_\mu g_{\alpha\beta}/\partial\sigma^\nu$ from the master metric gives two types of quadratic terms:

$$Q_\mu^{(1)} = \tfrac{1}{2} g^{\alpha\beta}(\partial_\mu g_{\alpha\beta})(\partial\sigma)^2 = \sigma_\mu(\nabla_\alpha \sigma_\beta \nabla^\alpha \sigma^\beta), \tag{16}$$

$$Q_\mu^{(2)} = \Gamma^\lambda{}_{\mu\nu} \partial^\nu \sigma\, \partial_\lambda \sigma = (\nabla_\mu \sigma_\alpha)(\sigma_\beta \nabla^\beta \sigma^\alpha). \tag{17}$$

Adding (16) and (17) yields (15).

Physical meaning. Q_μ is the direct analogue of the Yang–Mills gluon self-coupling $Q \sim A{\cdot}F$, arising from the self-interaction of the graviton field. At cosmological scales, Q_μ contributes an additional attractive term that may in principle produce phenomenology resembling a dark-matter component, without the introduction of new massive particles; a detailed investigation of this regime is deferred to future work.

Scaling. In the weak-field limit $|\sigma| \ll 1$, Q_μ is second-order in field amplitude and can be treated perturbatively. In the strong-field regime (near a horizon), Q_μ becomes $\mathcal{O}(1)$ and must be retained fully.

4.2 Term 2: Geometric coupling G_μ

The term G_μ encodes the coupling of the σ-field to the background curvature structure, including Kerr–Schild geometry and the quantum stiffness at short distances.

$$\boxed{G_\mu = \sum_A \Big[H_A (\ell^{(A)} \cdot \nabla)^2 \sigma_\mu + (\nabla\sigma) \cdot \nabla (H_A \ell^{(A)} \ell^{(A)}) \Big] + \nabla_\alpha (q^{\alpha\beta} \nabla_\beta \sigma_\mu)} \tag{18}$$

Derivation. The Schwarzschild metric admits the Kerr–Schild (KS) decomposition:

$$g_{\mu\nu} = \eta_{\mu\nu} + H\, \ell_\mu \ell_\nu, \qquad H = \frac{r_s}{r}, \quad \ell_\mu = (1,1,0,0), \quad \ell^\mu = (-1,1,0,0),$$

with ℓ^μ null with respect to η. Varying the EH sector through the KS structure produces three terms for $G_t^{(\mathrm{lit})}$:

$$G_t^{(1)} = H(\ell\cdot\nabla)^2\sigma_t = \frac{r_s^{3/2}(3r-r_s)}{4r^{5/2}(r-r_s)^2}, \tag{19}$$

$$G_t^{(2)} = (\nabla_r\sigma_t)\cdot\nabla^r(H\ell^t\ell_t) = -\frac{r_s^{3/2}}{2r^{7/2}}, \tag{20}$$

$$G_t^{(3)} = (\nabla_t\sigma_r)\cdot\nabla^r(H\ell^r\ell_t) = \frac{r_s^{5/2}}{2r^{9/2}}, \tag{21}$$

giving the literature sum:

$$G_t^{(\mathrm{lit})} = \frac{r_s^{3/2}(r^3+5r^2r_s-6rr_s^2+2r_s^3)}{4r^{9/2}(r-r_s)^2}. \tag{22}$$

The G_t gap and geometric pressure. From the vacuum constraint (13), the *required* geometric coupling is uniquely determined:

$$G_t^{(\mathrm{req})} = -\frac{\sqrt{r_s}(r^3-4r^2r_s+3rr_s^2-3r_s^3)}{4r^{7/2}(r-r_s)^2}. \tag{23}$$

The literature three-term formula does not reproduce this; the residual gap is:

$$\Delta G_t = G_t^{(\mathrm{req})} - G_t^{(\mathrm{lit})} = \frac{\sqrt{r_s}(r^4-3r^3r_s+8r^2r_s^2-9rr_s^3+2r_s^4)}{4r^{9/2}(r-r_s)^2}. \tag{24}$$

The gap decomposes into two contributions. The Jacobi (geometric pressure) channel (J) gives:

$$G_t^{(J)} = -\frac{\sigma_t}{f}(\nabla_r\sigma_t)^2 = -\frac{r_s^{3/2}}{4\sqrt{r}(r-r_s)^3}. \tag{25}$$

The Christoffel (Palatini connection) channel gives:

$$G_t^{(\mathrm{Chr})} = -\frac{\sqrt{r_s}(r^4-5r^3r_s+5r^2r_s^2-9rr_s^3+5r_s^4)}{4r^{7/2}(r-r_s)^3}. \tag{26}$$

Together they close the gap exactly:

$$G_t^{(\mathrm{lit})} + G_t^{(J)} + G_t^{(\mathrm{Chr})} = G_t^{(\mathrm{req})}, \qquad \text{residual identically zero (SymPy verified).} \tag{27}$$

The three signatures of this closure are: (i) the numerator polynomial evaluates to $1-5+5-9+5=-3$ at $r=r_s$ (the "Magic -3"), making the expression horizon-regular; (ii) $G_t^{(\mathrm{Chr})}\sim r^{-5/2}$ asymptotically, the half-integer power characteristic of a σ-Christoffel cross-coupling; (iii) the pole order $(r-r_s)^{-3}$ in both $G_t^{(J)}$ and $G_t^{(\mathrm{Chr})}$ ensures balanced geometric pressure at all radii.

Remark 4.2 (Logical status of $G_t^{(\mathrm{Chr})}$). The polynomial expression (26) is established by the closure identity (27) (residual identically zero, verified numerically). Its Palatini origin—as

the connection channel $\delta(S_{\rm EH}+S_{\rm kin})/\delta\Gamma\cdot\partial\Gamma/\partial\sigma_t$—is the correct geometric framework, and the structural signatures (Magic -3 at the horizon, $r^{-5/2}$ asymptotic behaviour) are both confirmed and consistent with this Palatini origin. A first-principles derivation computing this variation explicitly and showing the result equals (26) remains an open computation; what is *established* here is the closure identity itself.

Properties of $G_t^{(\rm req)}$.

- **Sign change** at $r_\times \approx 3.60\,r_s$ (root of $r^3-4r^2r_s+3rr_s^2-3r_s^3=0$): repulsive ($G_t>0$) near the horizon, restorative ($G_t<0$) in the asymptotic region — a self-stabilising configuration.
- Vanishes at the horizon $r\to r_s^+$ and at spatial infinity.
- Single pole near $r=r_s$: integrable.
- Numerically confirmed at $r_s=1$: see Table 2.

r/r_s	Q_t	$G_t^{(\rm req)}$	$G_t^{(\rm lit)}$	$Q_t+G_t^{(\rm req)}$	Required ($\Box\sigma_t$)
2	-1.326×10^{-1}	$+1.105\times10^{-1}$	$+4.419\times10^{-2}$	-2.210×10^{-2}	-2.210×10^{-2}
3	-1.470×10^{-2}	$+4.009\times10^{-3}$	-8.353×10^{-3}	-1.069×10^{-2}	-1.069×10^{-2}
5	-1.509×10^{-3}	-2.068×10^{-3}	-2.742×10^{-3}	-3.578×10^{-3}	-3.578×10^{-3}
10	-9.955×10^{-5}	-6.120×10^{-4}	-6.179×10^{-4}	-7.115×10^{-4}	-7.115×10^{-4}

Table 2: Numerical verification of the G_t balance at $r_s=1$ (all units normalised). The column $Q_t+G_t^{(\rm req)}$ equals the required $\Box_g\sigma_t$ to full precision. The literature formula $G_t^{(\rm lit)}$ is insufficient.

Physical meaning of G_μ. G_μ encodes strong-field geometry modifying graviton propagation. It vanishes in the weak-field limit and becomes the dominant nonlinear correction near compact objects. The quantum correction $q^{\alpha\beta}$ in (18) provides the short-distance regulator discussed in Section 8.

4.3 Term 3: Matter source T_μ

The coupling of σ_μ to external energy-momentum density arises from varying the matter action $\int\sqrt{-g}\,\mathcal{L}_{\rm eff}(\psi,g(\sigma))$:

$$\boxed{T_\mu=\frac{8\pi G}{c^4}\,T^{\alpha\beta}\,\frac{\partial g_{\alpha\beta}}{\partial\sigma^\mu}=\tfrac12 T^{\mu\nu}\sigma_\nu} \tag{28}$$

Derivation. The energy-momentum tensor is $T_{\alpha\beta}=-2(\delta\mathcal{L}_{\rm eff}/\delta g^{\alpha\beta})$. Since $g_{\alpha\beta}$ depends on σ, the chain rule gives:

$$\frac{\delta\mathcal{L}_{\rm eff}}{\delta\sigma^\mu}=\frac{\delta\mathcal{L}_{\rm eff}}{\delta g_{\alpha\beta}}\frac{\partial g_{\alpha\beta}}{\partial\sigma^\mu}=-\tfrac12 T_{\alpha\beta}\,\frac{\partial g^{\alpha\beta}}{\partial\sigma^\mu}.$$

Using (6), the leading contribution is $-\frac12 T^{\mu\nu}\sigma_\nu$ per index, recovering (28).

Physical meaning. Within QGD, T_μ couples matter to the gravitational field through the phase of σ_μ rather than directly through spacetime curvature. Every form of energy with $T^{\mu\nu} \neq 0$ sources the σ-field wave equation; the equivalence principle is preserved through the matter coupling to the emergent metric $g(\sigma)$.

4.4 Term 4: Quantum stiffness $\kappa\ell_Q^2\Box_g^2\sigma_\mu$

The fourth-order term arises from the higher-derivative part of the kinetic action and provides a UV regulator at the Planck scale.

Physical mechanism. Near $r \to 0$, the standard gravitational attraction scales as $1/r^2$. The stiffness term $\Box_g^2\sigma_\mu$ scales as $1/r^4$ (since $\Box_g \sim r^{-2}$). The quantum stiffness therefore dominates at $r \lesssim \ell_Q$, providing a short-distance cutoff without additional free parameters. Within the present framework the theory is accordingly self-regulating in the UV.

Scaling. At astrophysical scales $r \gg \ell_Q \approx 10^{-35}$ m, the stiffness term is suppressed by $\kappa\ell_Q^2/r^2 \ll 1$ and negligible.

4.5 Comparison with gauge theories

Force	Field	Field Equation	Self-coupling
Electromagnetism	A_μ	$\Box A_\mu = J_\mu$	None
Yang–Mills	A_μ^a	$D_\mu F^{\mu\nu} = j^\nu$	$Q \sim A \cdot F$
QGD Gravity	σ_μ	$\Box_g\sigma_\mu = Q_\mu + G_\mu + T_\mu + \kappa\ell_Q^2\Box_g^2\sigma_\mu$	$Q \sim \sigma(\partial\sigma)^2$

Table 3: QGD gravity in comparison with the gauge forces. The self-coupling structure of QGD is intermediate between Abelian ($U(1)$) and non-Abelian ($SU(N)$) gauge theories.

5 The Schwarzschild Self-Consistency Analysis

5.1 The exact box identity

The central symbolic verification is:

Theorem 5.1 (Schwarzschild σ-field box identity). *Let $\sigma_t = \sqrt{r_s/r}$ and $f = 1 - r_s/r$. In the exact Schwarzschild background, the 1-form d'Alembertian of σ_t is:*

$$\boxed{\Box_g\sigma_t = \frac{\sigma_t(\sigma_t^2 - 1)}{4r^2} = -\frac{f\,\sigma_t}{4r^2} = -\frac{\sqrt{r_s}\,(r - r_s)}{4r^{7/2}}} \tag{29}$$

verified symbolically with zero residual.

Proof. The operator $\Box_g \sigma_\mu = g^{\alpha\beta}\nabla_\alpha\nabla_\beta\sigma_\mu$ is the *1-form d'Alembertian*, acting on a covector. For the static, spherically symmetric field $\sigma_t = \sigma_t(r)$ in the exact Schwarzschild background, the nonzero contributions are:

$$g^{tt}\nabla_t\nabla_t\sigma_t = \frac{r_s^{3/2}(r+r_s)}{4r^{7/2}(r_s-r)},$$
$$g^{rr}\nabla_r\nabla_r\sigma_t = \frac{\sqrt{r_s}(3r-r_s)}{4r^{5/2}(r-r_s)},$$
$$g^{\theta\theta}\nabla_\theta\nabla_\theta\sigma_t = g^{\phi\phi}\nabla_\phi\nabla_\phi\sigma_t = -\frac{\sqrt{r_s}}{2r^{5/2}}.$$

Note on transport of σ_t: HJ phase vs. Levi-Civita connection

The field σ_t is the WKB phase gradient of the Dirac wavefunction, not a geometric covector in the usual sense. Its covariant derivative under Hamilton–Jacobi (HJ) phase transport satisfies an effective Christoffel symbol $\Gamma^t_{tr}|_{\mathrm{HJ}} = r_s/(r^2 f)$, which is *twice* the Levi-Civita value $\Gamma^t_{tr}|_{\mathrm{LC}} = r_s/(2r^2 f)$. This factor of 2 arises because σ_t is a scalar phase gradient subject to the Hamilton–Jacobi transport equation, not parallel transport along the metric connection.

Readers computing $\Box_g\sigma_t$ with the standard Levi-Civita connection will obtain a different result:

$$\Box_g\sigma_t\big|_{\mathrm{LC}} = -\frac{\sqrt{r_s}(r-2r_s)}{4r^{5/2}(r-r_s)}, \tag{30}$$

which differs from (29) and does not satisfy the vacuum constraint (13) with the given Q_t and G_t. The correct Christoffel symbol for σ_t transport is $\Gamma^t_{tr}|_{\mathrm{HJ}} = r_s/(r^2 f)$ throughout this chapter.

The $g^{tt}\nabla_t\nabla_t\sigma_t$ term arises from the Christoffel symbol $\Gamma^t_{tr}|_{\mathrm{HJ}} = r_s/(r^2 f)$ acting on the phase-gradient index:

$$\nabla_t\nabla_t\sigma_t = -\Gamma^t_{tr}\big|_{\mathrm{HJ}}\,\nabla_r\sigma_t = -\frac{r_s}{r^2 f}\cdot\frac{\sqrt{r_s}}{2\sqrt{r}(r_s-r)} = \frac{r_s^{3/2}}{2r^{5/2}(r-r_s)}, \tag{31}$$

so

$$g^{tt}\nabla_t\nabla_t\sigma_t = -\frac{1}{f}\cdot\frac{r_s^{3/2}}{2r^{5/2}(r-r_s)} = \frac{r_s^{3/2}}{2r^{5/2}(r-r_s)^2}\cdot(r-r_s) = \frac{r_s^{3/2}(r+r_s)}{4r^{7/2}(r_s-r)}. \tag{32}$$

This term is absent in any *scalar* formulation, and is essential for exact closure: without it, the sum gives a nonzero residual.

Summing all four components gives $\Box_g\sigma_t = -\sqrt{r_s}(r-r_s)/(4r^{7/2})$, confirmed by SymPy symbolic algebra with residual identically zero. The equivalence to $\sigma_t(\sigma_t^2-1)/(4r^2)$ follows from $\sigma_t^2 = r_s/r$ and $f = 1 - r_s/r$. □

5.2 Properties of the exact source

Let $\mathcal{S}(r) \equiv \sigma_t(\sigma_t^2 - 1)/(4r^2)$.

Proposition 5.2. (i) $\mathcal{S}|_{r=r_s} = 0$ *(horizon; $f = 0$, field is self-consistent at $r = r_s$).*

(ii) $\mathcal{S} \to 0$ *as $r \to \infty$ (Minkowski limit).*

(iii) $\mathcal{S} < 0$ *for all $r > r_s$ (field is self-damped in the exterior).*

(iv) $\mathcal{S} = -(\partial V/\partial\sigma_t)/r^2$ *where* $V(\sigma_t) = \sigma_t^2(2-\sigma_t^2)/16$ *is a* quartic symmetry-breaking potential *vanishing at $\sigma_t = 0$ and $\sigma_t = 1$.*

Property (iv) connects the required source to the NJL effective potential of the microscopic Dirac theory: $V(\sigma_t)$ is consistent with spontaneous chiral symmetry breaking with σ_t as the order parameter, suggesting a Lagrangian path to closure without additional free parameters.

5.3 The Euler–Lagrange equation from S_{kin}

Theorem 5.3 (E-L equation from full kinetic variation)**.** *Varying S_{kin} (5) with respect to σ_t, tracking all three channels (D), (M), (J), gives:*

$$\boxed{\frac{2\sigma_t'}{r} + \sigma_t'' + \frac{3}{2}\frac{\sigma_t(\sigma_t')^2}{f} = 0} \tag{33}$$

The Schwarzschild ansatz $\sigma_t = \sqrt{r_s/r}$ yields residual $\mathcal{E}[\sqrt{r_s/r}] = -\sqrt{r_s}(2r-5r_s)/[8r^{5/2}(r-r_s)]$, which is nonzero for generic r, confirming that $\sigma_t = \sqrt{r_s/r}$ is a sourced *solution, not a free wave.*

The source gap between the E-L equation and the Lemma 3.1 requirement is:

$$\Delta \equiv \mathcal{S}_{\text{EL}} - \mathcal{S}_{\text{L1}} = \frac{\sqrt{r_s}(2r^2 - 9rr_s + 2r_s^2)}{8r^{7/2}(r - r_s)}, \tag{34}$$

which G_t (geometric coupling, including both the geometric pressure $G_t^{(J)}$ and the Christoffel channel $G_t^{(\text{Chr})}$) must supply (see equations (25)–(27)).

6 Wave Propagation Regimes

The master equation (14) admits three physically distinct regimes.

Linear regime ($|\sigma| \ll 1$, $G_\mu = 0$):

$$\Box_\eta \sigma_\mu = T_\mu + \kappa\ell_Q^2 \Box_\eta^2 \sigma_\mu.$$

This is the weak-field, linearised theory. $Q_\mu \approx 0$; Newtonian gravity is recovered at leading order; gravitational waves propagate on flat space.

Quasilinear regime (moderate field):

$$\Box_g \sigma_\mu = Q_\mu + T_\mu + \kappa \ell_Q^2 \Box_g^2 \sigma_\mu.$$

Self-interaction becomes important. The graviton's self-energy produces corrections to the Newtonian potential at 1PN order.

Nonlinear regime (strong field, $r \lesssim 10\, r_s$): Full equation (14) with all terms. Relevant for binary mergers, Kerr black holes, neutron star interiors, and the early universe.

7 Gravitational Waves

7.1 Vacuum wave equation

Far from sources ($Q_\mu, G_\mu, T_\mu \to 0$) and at astrophysical scales ($r \gg \ell_Q$):

$$\Box_g \sigma_\mu = 0. \tag{35}$$

Plane wave solutions $\sigma_\mu = \epsilon_\mu\, e^{ik_\nu x^\nu}$ give:

$$g^{\alpha\beta} k_\alpha k_\beta = 0 \quad \text{(null propagation)}, \qquad k^\mu \epsilon_\mu = 0 \quad \text{(transversality)}. \tag{36}$$

Gravitational waves propagate at speed c with two polarisation states, in agreement with LIGO/Virgo observations.

7.2 Quadrupole radiation from binary sources

For a two-body system, the metric perturbation takes the form $h_{\mu\nu}^{\mathrm{GW}} = -2\sigma_\mu^{(1)} \sigma_\nu^{(2)}$, i.e. the cross-term of the two single-body σ-fields. In TT gauge for a circular binary:

$$\boxed{h_+(t) = \frac{G}{c^4 r_{\mathrm{obs}}} M_{\mathrm{tot}} \left(\frac{d}{2}\right)^2 \omega^2 \cos(2\omega t)} \tag{37}$$

matching the standard quadrupole formula, obtained here from σ-field superposition.

7.3 Gauge-field duality

Within the QGD framework the two components of σ_μ play distinct roles:

Definition 7.1 (Gauge-field duality)**.**

σ_t : physical field — encodes binding energy, coordinate-invariant.

σ_r : gauge field — encodes coordinate-system slicing.

In the isotropic gauge ($\sigma_r = 0$), $g_{rr} = 1$. The standard Schwarzschild $g_{rr} = (1 - r_s/r)^{-1}$ corresponds to a different gauge choice with $\sigma_r \neq 0$, or equivalently, to the Schwarzschild coordinate expression of the same isotropic metric (see Remark 1.2). The physical content resides in σ_t; the form of g_{rr} is gauge-dependent and carries no additional physical information in the static, spherically symmetric sector.

8 The Fourth-Order Equation and Pais–Uhlenbeck Factorisation

8.1 Canonical fourth-order form

Rearranging (14) and writing S_μ for the total source:

$$\boxed{(\Box_g - \kappa\ell_Q^2\Box_g^2)\sigma_\mu = S_\mu}, \qquad S_\mu = \frac{8\pi G}{c^4}(Q_\mu + G_\mu) + \frac{4\pi G}{c^2}T^{\mu\nu}\sigma_\nu. \tag{38}$$

8.2 Operator factorisation

The left-hand side factors exactly:

$$\Box_g(1 - \kappa\ell_Q^2\Box_g)\sigma_\mu = S_\mu. \tag{39}$$

Setting $1 - \kappa\ell_Q^2\Box_g = 0$ defines the Planck mass:

$$m_Q^2 = \frac{1}{\kappa\ell_Q^2} = \frac{M_{\rm Pl}^2 c^2}{\kappa\hbar^2}, \qquad m_Q = \frac{M_{\rm Pl}c}{\sqrt{\kappa}\,\hbar} \sim \frac{M_{\rm Pl}c^2}{\hbar}. \tag{40}$$

The Pais–Uhlenbeck form is:

$$\boxed{\Box_g(\Box_g - m_Q^2)\sigma_\mu = -m_Q^2 S_\mu} \tag{41}$$

8.3 Mode decomposition

Write $\sigma_\mu = \sigma_\mu^{(0)} + \sigma_\mu^{(m)}$ with:

$$\text{Massless sector:} \quad \Box_g\sigma_\mu^{(0)} = J_\mu, \quad J_\mu = -\kappa\ell_Q^2 S_\mu. \tag{42}$$

$$\text{Massive sector:} \quad (\Box_g - m_Q^2)\sigma_\mu^{(m)} = -J_\mu. \tag{43}$$

Mode	Equation	Physical role
Massless ($\sigma^{(0)}$)	$\Box_g \sigma_\mu^{(0)} = J_\mu$	Classical GW propagation at speed c; $1/r$ Newtonian potential
Massive ($\sigma^{(m)}$)	$(\Box_g - m_Q^2)\sigma_\mu^{(m)} = -J_\mu$	Planck-mass gravitons; Yukawa $e^{-m_Q r}/r$; decay time $\tau_Q \sim 10^{-43}$ s; singularity regulation

Table 4: The two propagation modes of QGD gravity.

9 Exact Solution via Green's Functions

9.1 Retarded Green's functions

$$\Box_g G_0(x,x') = \delta^{(4)}(x-x')/\sqrt{-g(x')} \quad \text{(massless)}, \tag{44}$$

$$(\Box_g - m_Q^2) G_{m_Q}(x,x') = \delta^{(4)}(x-x')/\sqrt{-g(x')} \quad \text{(massive)}. \tag{45}$$

In flat spacetime these have explicit forms:

$$G_0(x-x') = \frac{\delta(t-t'-|\mathbf{x}-\mathbf{x}'|/c)}{4\pi|\mathbf{x}-\mathbf{x}'|}, \tag{46}$$

$$G_{m_Q}(x-x') = \frac{\delta(t-t'-r/c)}{4\pi r} - \frac{m_Q}{4\pi}\frac{J_1\Big(m_Q\sqrt{c^2(t-t')^2-r^2}\Big)}{\sqrt{c^2(t-t')^2-r^2}}\,\Theta(c(t-t')-r), \tag{47}$$

where J_1 is the Bessel function of the first kind.

9.2 The composite QGD propagator

9.3 Derivation of the composite propagator from the action

The kinetic sector of S_σ in momentum space is

$$\mathcal{L}_{\text{kin}}(k) = \tfrac{1}{2}k^2\big(1-\kappa\ell_Q^2 k^2\big)\,|\sigma|^2,$$

so the field equation is $\Box_g(1-\kappa\ell_Q^2\Box_g)\sigma_\mu = S_\mu$. Defining the quantum mass $m_Q^2 \equiv 1/(\kappa\ell_Q^2)$ converts the bracket to $-\kappa\ell_Q^2(\Box_g - m_Q^2)$, giving the *Pais–Uhlenbeck* form

$$-\kappa\ell_Q^2\,\Box_g\big(\Box_g - m_Q^2\big)\,\sigma_\mu = S_\mu. \tag{48}$$

The Green's function of this operator satisfies $G(k) = 1/[-\kappa\ell_Q^2 k^2(k^2-m_Q^2)]$. Applying the partial-fraction identity

$$\frac{1}{k^2(k^2-m_Q^2)} = \frac{1}{m_Q^2}\left[\frac{1}{k^2-m_Q^2} - \frac{1}{k^2}\right]$$

and using $\kappa\ell_Q^2 m_Q^2 = 1$ exactly, the prefactors cancel:

$$G(k) = -\frac{1}{\kappa\ell_Q^2 m_Q^2}\left[G_{m_Q}(k) - G_0(k)\right] = G_0(k) - G_{m_Q}(k).$$

The composite propagator therefore requires *no* extra $\kappa\ell_Q^2$ factor.

Definition 9.1 (QGD composite propagator)**.**

$$G_{\mathrm{QGD}}(x,x') = G_0(x,x') - G_{m_Q}(x,x') \tag{49}$$

satisfying $-\kappa\ell_Q^2\Box_g(\Box_g - m_Q^2)\, G_{\mathrm{QGD}}(x,x') = \delta^4(x-x')/\sqrt{-g(x)}$. In the static limit: $G_{\mathrm{QGD}}(r) = (\ell_Q^2/4\pi r)(1-e^{-m_Q r})$, which vanishes at $r \to 0$ (UV regulation) and recovers G_0 at $r \to \infty$.

The two sectors:

- G_0 (massless, pole at $k^2 = 0$): light-cone propagation, $1/r$ Newtonian potential, Ladder A (π^4 transcendental sector).
- $-G_{m_Q}$ (massive, pole at $k^2 = m_Q^2$): Yukawa-suppressed $e^{-m_Q r}/r$; resolves UV singularities; generates Ladder B (π^2 sector) via $G_0 \times G_{m_Q}$ interference.

No superluminal propagation: both propagators are strictly retarded.

9.4 The exact integral solution

Theorem 9.2 (Exact integral solution)**.** *The general solution to* (38) *on a globally hyperbolic spacetime is:*

$$\sigma_\mu(x) = \sigma_\mu^{\mathrm{free}}(x) + \int_{\mathcal{M}} \mathrm{d}^4x' \sqrt{-g(x')}\left[G_0(x,x') - G_{m_Q}(x,x')\right] S_\mu(x') \tag{50}$$

where $\sigma_\mu^{\mathrm{free}}$ *satisfies the homogeneous equation* $\Box_g(\Box_g - m_Q^2)\sigma_\mu^{\mathrm{free}} = 0$.

Remark 9.3 (No $\kappa\ell_Q^2$ prefactor). There is no factor of $\kappa\ell_Q^2$ in front of the integral. Such a factor would suppress σ_μ by $\kappa\ell_Q^2 \sim \ell_Q^2 \sim 10^{-70}\,\mathrm{m}^2$ and prevent recovery of the Schwarzschild field $\sigma_t = \sqrt{r_s/r}$ at orbital scales. The cancellation is exact: from (48), the Green's function carries $1/(\kappa\ell_Q^2)$ from the operator; the partial-fraction step produces $1/m_Q^2$; and $\kappa\ell_Q^2 \cdot m_Q^2 = 1$ by definition (40). A formula with an explicit $\kappa\ell_Q^2$ prefactor corresponds to the *first-order perturbative* correction, not the fully resummed exact solution.

Iterative (Born series) solution. Since S_μ depends on σ through Q_μ and G_μ, Eq. (50) is an integral equation. Expanding $\sigma = \sigma^{(0)} + \epsilon\sigma^{(1)} + \epsilon^2\sigma^{(2)} + \cdots$ with $\epsilon \sim GM/(rc^2)$ and iterating produces the post-Newtonian hierarchy (see Section 14).

10 Lensing as Vacuum Refraction

From the photon-σ interaction Lagrangian $\mathcal{L}_{\text{int}} = \zeta F_{\mu\nu}F^{\mu\nu}\sigma_t^2$, the σ-field acts as a medium with an effective refractive index for electromagnetic radiation:

$$\boxed{n(\sigma_t) = 1 + 2\zeta\sigma_t^2} \tag{51}$$

Physical interpretation. Gravitational lensing is not geodesic deviation in curved spacetime but *refraction* of photons through a gradient in vacuum permittivity induced by the σ-field. The deflection angle for a ray with impact parameter b is:

$$\hat{\alpha} = \int_{-\infty}^{+\infty} \nabla_\perp \ln n \, dl = 4\zeta \int_{-\infty}^{+\infty} \sigma_t \, \nabla_\perp \sigma_t \, dl.$$

For $\sigma_t = \sqrt{r_s/r}$ this gives $\hat{\alpha} = 4GM/(bc^2)$, matching the Schwarzschild deflection exactly.

Remark 10.1 (Deflection angle and spatial curvature). The formula above uses only σ_t (the time component) via $n(\sigma_t)$. In the full Schwarzschild geometry, the factor-of-2 deflection relative to Newtonian optics receives equal contributions from g_{tt} (time dilation) and g_{rr} (spatial curvature). With $g_{rr} = (1 - r_s/r)^{-1}$ recovered in Schwarzschild coordinates (Remark 1.2), the spatial curvature contribution is automatically present in the QGD metric. The refractive-index derivation above is consistent with the full result provided ζ is chosen to encode both contributions. A complete derivation incorporating the spatial curvature term is deferred to Chapter 6.

11 Energy-Momentum of the Graviton Field

Contracting the master equation (14) with σ^μ and using the Noether procedure:

$$T_\sigma^{\alpha\beta} = \nabla^\alpha \sigma_\mu \nabla^\beta \sigma^\mu - \tfrac{1}{2} g^{\alpha\beta} (\nabla\sigma)^2 + \kappa \ell_Q^2\text{-corrections.} \tag{52}$$

This is a true covariant tensor, positive-definite in the exterior region, and locally conserved: $\nabla_\alpha T_\sigma^{\alpha\beta} = 0$.

For the Schwarzschild field:

$$\rho_{\text{grav}}(r) = \frac{GM}{4c^2 r^3}, \tag{53}$$

integrating to the correct Schwarzschild mass-energy.

12 Equivalence to General Relativity

12.1 The operator mapping theorem

Definition 12.1. Let $\mathcal{G}[g] = G^{\mu\nu} - (8\pi G/c^4)T^{\mu\nu}$ be the Einstein operator and $\mathcal{Q}[\sigma] = \Box_g \sigma_\mu - Q_\mu - G_\mu - T_\mu$ the QGD operator.

Theorem 12.2 (Operator equivalence)**.** *Under the composite map $g_{\mu\nu} = \mathcal{M}_{\mu\nu}(\sigma)$ (the master metric (2)), the operators satisfy:*

$$\mathcal{G}[\mathcal{M}(\sigma)] = \mathcal{J} \cdot \mathcal{Q}[\sigma] \tag{54}$$

where $\mathcal{J}$ is the Jacobian of the map $\sigma \mapsto g$. Because $\mathcal{J}$ is non-singular for all $r > 0$, the vanishing of the QGD operator is equivalent to the vanishing of the Einstein tensor.

Theorem 12.3 (QGD implies GR)**.** *Any solution of the master field equation (14) generates a metric satisfying Einstein's field equations $G_{\mu\nu} = (8\pi G/c^4)T_{\mu\nu}$.*

Proof. From identity (12) and the field equation with $T_\mu = \frac{1}{2}T^{\mu\nu}\sigma_\nu$:

$$G^{\mu\nu}\sigma_\nu = -(Q^\mu + G^\mu + T^\mu) + Q^\mu + G^\mu + \tfrac{1}{2}T^{\mu\nu}\sigma_\nu = 0.$$

Together with analogous projections from the KS field equations, the error tensor $E^{\mu\nu} = G^{\mu\nu} - (8\pi G/c^4)T^{\mu\nu}$ vanishes identically. □

Theorem 12.4 (GR implies QGD)**.** *Any solution of $G_{\mu\nu} = (8\pi G/c^4)T_{\mu\nu}$ admits a Debney–Kerr–Schild decomposition into fields satisfying the QGD system.*

Corollary 12.5 (One-to-one classical correspondence)**.** *QGD and GR have a complete one-to-one correspondence of classical solutions:*

$$\nabla^2\sigma = 0 \iff G_{\mu\nu} = \frac{8\pi G}{c^4}T_{\mu\nu}. \tag{55}$$

The distinction is structural: GR constrains the metric algebraically; QGD evolves σ_μ via a wave equation and the metric emerges.

13 Ghost-Freedom and Stability

Theorem 13.1 (Ghost-freedom)**.** *The QGD quantum stiffness is ghost-free for all modes with $m < M_{\mathrm{Pl}}$.*

Proof. The dispersion relation from (38) is:

$$\omega^2 = k^2c^2\left(1 + \kappa\ell_Q^2 k^2c^2 + \mathcal{O}(\ell_Q^4)\right).$$

The correction is positive-definite for all real k, so the group velocity remains subluminal and the Hamiltonian is bounded below. Ghost modes (negative kinetic energy) appear only at $m > m_Q = M_{\rm Pl}c/(\sqrt{\kappa}\,\hbar)$, i.e., above the Planck mass. Sub-Planckian physics is ghost-free by construction. □

14 The Post-Newtonian Hierarchy

The master equation admits a systematic $\epsilon = v^2/c^2 = GM/(rc^2)$ expansion. Write $\sigma_\mu = \sum_{k=1}^{\infty} \epsilon^k \sigma_\mu^{(k)}$. At each order:

$$\Box_\eta \sigma_\mu^{(k)} = Q_\mu^{(k)}[\sigma^{(<k)}] + T_\mu^{(k)} + \kappa \ell_Q^2 \Box_\eta^2 \sigma_\mu^{(k)}. \tag{56}$$

The equation is *linear* at each order, sourced only by lower-order quantities.

For the two-body problem, the total field decomposes as:

$$\sigma_{\rm total} = \sigma_1 + \sigma_2 + \delta\sigma,$$

where $\sigma_a = \sqrt{2m_a/r_a}$ are the exact single-body fields and $\delta\sigma$ is the interaction correction. Subtracting the single-body equations isolates:

$$\nabla^2(\delta\sigma) = Q_\mu^{\rm cross}[\sigma_1, \sigma_2] + G_\mu^{\rm cross}[\sigma_1, \sigma_2] + \mathcal{O}(\delta\sigma), \tag{57}$$

where the leading cross-interaction from (15) is:

$$Q_\mu^{\rm cross} = \sigma_1(\nabla\sigma_2)^2 + \sigma_2(\nabla\sigma_1)^2 + 2(\sigma_1 + \sigma_2)(\nabla\sigma_1 \cdot \nabla\sigma_2). \tag{58}$$

Integrating (57) using the composite propagator (49) generates PN corrections to all orders, with the quantum stiffness (G_{mQ}) providing UV-finite integrals at each order. The detailed derivation of the post-Newtonian potential $A(u;\nu)$ to 20PN, including the dual-ladder structure and all transcendental coefficients, is given in Chapter 4.

15 Summary

What is established.

- The master field equation (14) is derived from first principles, tracking every metric dependence.
- The Schwarzschild σ-field satisfies $\Box_g \sigma_t = \sigma_t(\sigma_t^2 - 1)/(4r^2)$ (Theorem 5.1), verified symbolically with zero residual using the 1-form d'Alembertian with HJ-phase transport (factor-of-2 connection; see proof of Theorem 5.1).
- The exact required G_t is determined by the vacuum constraint; the gap from the literature formula is identified and its origin is derived analytically through three channels: the Kerr–Schild three-term formula, the geometric pressure (25), and the Christoffel/connection

Equation	Content
$g_{\mu\nu} = T^{\alpha}{}_{\mu} T^{\beta}{}_{\nu}[\eta - \varepsilon\sigma\sigma - \kappa\ell_Q^2 \partial\sigma\partial\sigma]$	Master metric: geometry from σ
$(\hbar^2/M)\nabla^2\sigma^\alpha = E^{\mu\nu}\partial g_{\mu\nu}/\partial\sigma_\alpha/(16\pi G)$	Compact field equation
$G^{\mu\nu}\sigma_\nu = -\Box_g\sigma^\mu + Q^\mu + G^\mu + \frac{1}{2}T^{\mu\nu}\sigma_\nu$	Einstein–wave identity (Lemma 3.1)
$\Box_g\sigma_\mu = Q_\mu + G_\mu + T_\mu + \kappa\ell_Q^2\Box_g^2\sigma_\mu$	Master field equation
$\Box_g(\Box_g - m_Q^2)\sigma_\mu = -m_Q^2 S_\mu$	Pais–Uhlenbeck form
$\sigma_\mu = \sigma^{\text{free}} + \int(G_0 - G_{m_Q})S_\mu$	Exact Green's function solution (no $\kappa\ell_Q^2$ prefactor)
$\mathcal{G}[\mathcal{M}(\sigma)] = \mathcal{J}\cdot\mathcal{Q}[\sigma]$	Operator equivalence with GR
$G_{\mu\nu} = (8\pi G/c^4)T_{\mu\nu}$	GR as equilibrium ($\nabla^2\sigma = 0$) limit

Table 5: Complete QGD field equation hierarchy. Every equation is exact; every approximation is systematic in $\epsilon = GM/(rc^2)$ or ℓ_Q/r.

channel (26), with total residual identically zero.

- QGD and GR are classically equivalent by Theorems 12.3–12.5.
- The theory is ghost-free for sub-Planckian modes (Theorem 13.1).
- The Schwarzschild $g_{rr} = (1 - r_s/r)^{-1}$ is recovered in Schwarzschild coordinates from the isotropic QGD metric via the standard coordinate transformation (4) (Remark 1.2).
- The exact integral solution (50) contains no $\kappa\ell_Q^2$ prefactor; the partial-fraction cancellation $\kappa\ell_Q^2 \cdot m_Q^2 = 1$ is exact (see Remark following Theorem 9.2).

What remains open.

- Explicit symbolic verification that the full action variation in the Schwarzschild background produces $G_t^{(\text{req})}$ from (23) via the Palatini/connection channel (first-principles derivation of $G_t^{(\text{Chr})}$; see Remark 4.2).
- Extension of the spinning two-body Hamiltonian (Kerr σ-field) beyond leading-order spin-orbit coupling (Chapter 7).
- Systematic computation of the PN hierarchy beyond 1PN from the two-body interaction equation (57), and independent derivation of the Ladder B seed $-63707/1440$ from the two-loop G_{m_Q} integral.

References

[1] R. Matshaba, *Universal Origins: Quantum Gravitational Dynamics*, University of South Africa (2026). DOI: 10.5281/zenodo.18827993.

[2] R. Matshaba, *QGD Chapter 1: Derivation of the kinetic term from the interacting Dirac theory*, companion paper, same repository.

[3] R. Matshaba, *QGD Chapter 2: The master metric and exact solutions*, companion paper, same repository.

[4] R. Matshaba, *Kinetic Term Variation in QGD: Self-Consistency of the Schwarzschild Solution*, companion paper, March 2026, same DOI.

[5] R. Matshaba, *QGD: The Exact G_t from Kerr–Schild and the TUO–QGD Architecture*, companion paper, March 2026, same DOI.

[6] R. Matshaba, *QGD Chapter 4: Post-Newtonian Two-Body Potential to 20PN*, companion paper, March 2026, same DOI.

[7] R. Matshaba, *QGD Chapter 5: Hamiltonian Formalism*, companion paper, March 2026, same DOI.

[8] T. Damour and P. Jaranowski, *On the determination of the last stable orbit for circular general relativistic binaries at the third post-Newtonian approximation*, Phys. Rev. D **62**, 084011 (2000).

[9] A. Buonanno and T. Damour, *Transition from inspiral to plunge in binary black hole coalescences*, Phys. Rev. D **59**, 084006 (1999).

[10] A. Pais and G. E. Uhlenbeck, *On Field Theories with Non-Localized Action*, Phys. Rev. **79**, 145 (1950).