

ON BASIC DOUBLE LINKS OF SQUAREFREE MONOMIAL IDEALS

PATRICIA KLEIN, MATTHEW KOBAN, AND JENNA RAJCHGOT

ABSTRACT. Nagel and Römer introduced the class of weakly vertex decomposable simplicial complexes, which include matroid, shifted, and Gorenstein complexes as well as vertex decomposable complexes. They proved that the Stanley–Reisner ideal of every weakly vertex decomposable simplicial complex is Gorenstein linked to an ideal of indeterminates via a sequence of basic double links. In this paper, we explore basic double links between squarefree monomial ideals beyond the weakly vertex decomposable setting.

Our first contribution is a structural result about certain basic double links which involve an edge ideal. Specifically, suppose $I(G)$ is the edge ideal of a graph G . When $I(G)$ is a basic double link of a monomial ideal B on an arbitrary homogeneous ideal A , we give a generating set for B in terms of G and show that this basic double link must be of degree 1. Our second focus is on examples from the literature of simplicial complexes known to be Cohen–Macaulay but not weakly vertex decomposable. We show that these examples are not basic double links of any other squarefree monomial ideals.

1. INTRODUCTION

Broadly speaking, Gorenstein liaison is a framework for studying which properties of one subscheme of the projective space \mathbb{P}^n can be transferred to another when their union is sufficiently nice, that is, when the subschemes are (*Gorenstein*) *linked*. Gorenstein links generate an equivalence relation whose equivalence classes are called *Gorenstein liaison classes*. Two subschemes in the same Gorenstein liaison class have the same codimension, and their homogeneous coordinate rings have the same depth. Hence, every subscheme that is in the *Gorenstein liaison class of a complete intersection* (abbreviated *glicci*) is arithmetically Cohen–Macaulay. An important open question in Gorenstein liaison is whether every arithmetically Cohen–Macaulay subscheme is glicci.

This question has garnered a great deal of interest, and there have been a number of important partial results. For example, Casanellas, Drozd, and Hartshorne [CDH05] showed that every arithmetically Gorenstein subscheme is glicci. Gorla [Gor08] showed that every standard determinantal scheme is glicci, thereby generalizing results of [Har07] and [KMMR⁺01]. Migliore and Nagel [MN13] showed that every arithmetically Cohen–Macaulay subscheme of \mathbb{P}^n , when viewed instead as a subscheme of \mathbb{P}^{n+1} , is glicci.

Many specific classes of arithmetically Cohen–Macaulay schemes have also been proved to be glicci. Examples include certain curves in \mathbb{P}^4 [CMR00, CMR01], various generalized determinantal and Pfaffian schemes [Gor07, DNG09, Gor10, GMN13, FK20, KR21, Ney], schemes defined by certain toric ideals of graphs [CG18, CSRT22], and many classes of (or closely related to) monomial schemes [MN00, MN02, HU07, KTY13].

Of particular relevance to this paper are the contributions made by Nagel and Römer [NR08] in the case of schemes defined by squarefree monomial ideals. Squarefree monomial ideals are associated to simplicial complexes through the Stanley–Reisner correspondence (see Section 2.1). Nagel and Römer introduced the class of weakly vertex decomposable simplicial complexes, which include matroid, shifted, Gorenstein, and vertex decomposable complexes. They showed that the schemes corresponding to weakly vertex decomposable complexes are glicci. More specifically, they constructed a sequence of combinatorially-defined *basic double links* (see Definition 2.1) from the original Stanley–Reisner subscheme to a coordinate subspace. Nagel and Römer also gave examples of naturally-occurring complexes that are Cohen–Macaulay but not weakly vertex decomposable. It is not known whether or not these examples are glicci.

In this paper, we extend the work of Nagel and Römer in two directions. To state our first main result (which appears in its precise form as Theorem 3.5), let G be a graph, and $I(G)$ its corresponding *edge ideal*. Call a homogeneous, saturated ideal of a polynomial ring in n variables over a field *glicci* if the subscheme of \mathbb{P}^{n-1} it defines is glicci.

Theorem. *Let $I(G)$ be an edge ideal in the polynomial ring $S = \mathbb{k}[x_1, \dots, x_n]$. Assume there is a basic double link $I(G) = fB + A$ where f is a homogeneous form in S , $B \subset S$ is a monomial ideal, and $A \subset S$ a homogeneous ideal. Then, up to rescaling the variables in S ,*

- (1) $f = x_{i_1} + \dots + x_{i_r}$ is a sum of distinct indeterminates in S ; and
- (2) B is a specific squarefree monomial ideal which is completely determined by f and G .

We use this result to show that the edge ideal corresponding to Figure 1 is not a basic double link of any other monomial ideal.

Our second goal is to further explore Gorenstein liaison for Stanley-Reisner subschemes associated to specific simplicial complexes which are not weakly vertex decomposable. In Proposition 4.2, we show that one of the examples studied by [NR08] is not a basic double link of any other Stanley-Reisner subscheme. In Proposition 4.5, we consider another example from [NR08] and preclude the existence of a basic double link under the slightly stronger hypothesis that all of the ideals involved in the basic double link are squarefree monomial ideals.

Our examples show that basic double links are not sufficient for studying Gorenstein liaison of Stanley-Reisner schemes if we would like to stay within the setting of Stanley-Reisner schemes. In Section 5, we explore what purchase might be gained by expanding beyond basic double links.

Throughout this document, let \mathbb{k} be an arbitrary field and $S = \mathbb{k}[x_1, \dots, x_n]$ be the standard graded polynomial ring over \mathbb{k} in n variables. We will reserve R for standard graded polynomial rings in specific examples, in which case n will be known and \mathbb{k} will be \mathbb{Q} .

Acknowledgements. M. Koban was partially supported by an NSERC Undergraduate Student Research Award (USRA) held at the University of Saskatchewan. M. Koban and J. Rajchgot were partially supported by NSERC grant RGPIN-2017-05732.

2. BACKGROUND

In this section, we review some background material. Section 2.1 concerns simplicial complexes and Stanley-Reisner ideals, and Section 2.2 covers some basic facts about Gorenstein liaison.

2.1. Simplicial Complexes. An abstract simplicial complex Δ on vertex set $[n] = \{1, \dots, n\}$ is a collection of subsets of $[n]$, closed under inclusion. An element $F \in \Delta$ is called a face. The dimension of a face $F \in \Delta$ is defined by $\dim(F) = |F| - 1$, where $|F|$ denotes the number of elements in the subset $F \subseteq [n]$. The dimension of the simplicial complex Δ is defined by $\dim(\Delta) = \max\{\dim(F) \mid F \in \Delta\}$. A simplicial complex is called **pure** if each maximal face of Δ has the same dimension.

If $F \subseteq [n]$, define $x_F = \prod_{i \in F} x_i$. To each simplicial complex there is an associated monomial ideal, called the Stanley-Reisner ideal of Δ and denoted I_Δ . The Stanley-Reisner ideal is generated by non-faces of Δ ; that is $I_\Delta = (x_F : F \subseteq [n], F \notin \Delta)$. This correspondence provides a bijection between simplicial complexes Δ on the vertex set $[n]$ and squarefree monomial ideals I_Δ in S . We will refer to the elements of the (unique) minimal generating set of a monomial ideal that consists of monic monomials as the monomial generators of the ideal. The monomial generators of the squarefree monomial ideal I_Δ correspond to the minimal non-faces of Δ .

We note that $\dim(S/I_\Delta)$ is equal to $\dim(\Delta) + 1$ and that Δ is pure if and only if I_Δ is height unmixed, that is, if all of the associated primes of I_Δ have the same height. Because I_Δ is a radical ideal and therefore has no embedded primes, I_Δ is height unmixed if and only if $\text{Spec}(S/I_\Delta)$ is equidimensional. We say that a simplicial complex Δ is Cohen-Macaulay whenever R/I_Δ is Cohen-Macaulay.

Given a vertex v of Δ , we define the following subcomplexes of Δ :

- the link of v , $\text{lk}_\Delta v = \{G \in \Delta \mid \{v\} \cup G \in \Delta, \{v\} \cap G = \emptyset\}$;
- the deletion of v , $\Delta_{-v} = \{G \in \Delta \mid \{v\} \cap G = \emptyset\}$.

When we form the Stanley–Reisner ideals of $\text{lk}_\Delta v$ and Δ_{-v} , we view both of these complexes as complexes on the vertex set $[n] \setminus \{v\}$.

Given a simplicial complex Δ on $[n]$ and a vertex k such that $\{k\} \notin \Delta$, define the **cone over Δ with apex k** be the complex $\Delta' = \{F \cup \{k\} \mid F \in \Delta\}$. Notice that $I_\Delta = I_{\Delta'} + (x_k)$ and that x_k does not divide any of the monomial generators of $I_{\Delta'}$.

A pure simplicial complex Δ is **vertex decomposable** if Δ is a simplex, if $\Delta = \{\emptyset\}$, or if there exists a vertex v such that $\text{lk}_\Delta v$ and Δ_{-v} are both pure and vertex decomposable. In the latter case, if $\dim(\text{lk}_\Delta v) = \dim(\Delta_{-v}) - 1$, then we call v a **shedding vertex**. Vertex decomposition was introduced in [PB80]. We say that a pure simplicial complex Δ is **weakly vertex decomposable** if Δ is a simplex, if $\Delta = \{\emptyset\}$, or if there is some vertex v such that $\text{lk}_\Delta v$ is weakly vertex decomposable and Δ_{-v} is Cohen–Macaulay. In the latter case, if $\dim(\text{lk}_\Delta v) = \dim(\Delta_{-v}) - 1$, then we call v a **weak shedding vertex**. Weak vertex decomposition was introduced by [NR08]. Whenever Δ is vertex decomposable, Δ is Cohen–Macaulay; hence, a vertex decomposable complex is weakly vertex decomposable, as the name suggests. Nagel and Römer [NR08, Theorem 3.3] showed that if Δ is even weakly vertex decomposable, then Δ is Cohen–Macaulay, and in fact I_Δ is *glicci* (defined below in Subsection 2.2).

2.2. Gorenstein Liaison. Here we review some definitions and results from Gorenstein liaison. For more information, see the surveys [MN01] or [MN21]. All ideals in this subsection are assumed to be proper, homogeneous, and saturated.

Let I, J , and L be unmixed ideals of S and assume that S/L is Gorenstein. We say that I and J are directly linked by L , denoted $I \sim_L J$, if $L : I = J$ and $L : J = I$. Importantly, S/I is Cohen–Macaulay if and only if S/J is Cohen–Macaulay. We say that I and J are in the same Gorenstein liaison class if there is a sequence of Gorenstein ideals L_1, \dots, L_r and links $I = I_0 \sim_{L_1} I_1 \sim \dots \sim_{L_r} I_r = J$, for some $r \geq 1$. If J is a complete intersection, then we say that I is in the Gorenstein liaison class of a complete intersection, abbreviated **glicci**. We say that a subscheme of \mathbb{P}^{n-1} is glicci if its homogeneous, saturated ideal is glicci. We call a simplicial complex Δ glicci if its Stanley–Reisner ideal I_Δ is glicci.

If $(S/A)_P$ is Gorenstein for all minimal primes P of A , we say that S/A is **generically Gorenstein**. For example, if A is radical, then S/A is generically Gorenstein.

Definition 2.1. Let $A \subset B$ be unmixed ideals of S such that S/A is Cohen–Macaulay and generically Gorenstein and $\text{ht}(A) + 1 = \text{ht}(B)$. Let $f \in S$ be a homogeneous element of degree $d > 0$ such that $A : f = A$. Then $C = fB + A$ is a **basic double link of degree d of B on A** .

Though it is not standard, we will also refer to the equation $C = fB + A$ as a basic double link as well as the ideal C itself.

Example 2.2. Let $C = (x_1x_3, x_2x_4) \subset \mathbb{k}[x_1, x_2, x_3, x_4]$. Taking $f = x_1$, $B = (x_3, x_2x_4)$ and $A = (x_2x_4)$, a basic double link is given by $(x_1x_3, x_2x_4) = x_1(x_3, x_2x_4) + (x_2x_4)$.

Theorem 2.3. [Har07] *If C is a basic double link of B on A , then C is linked to B in two steps.*

Nagel and Römer [NR08] gave a connection between weak vertex decomposition and basic double linkage in the squarefree monomial ideal setting. They considered the special case of basic double links in which A , B , and C are all squarefree monomial ideals and $f = x_k$ is some variable of S .

Before stating their result, we make two standard observations: First, if Δ is not a cone over Δ_{-k} , then $\dim(\Delta) = \dim(\Delta_{-k})$. Second, let I be an ideal of S . If I is glicci and y is an indeterminate, then the extension of I to the ring $S[y]$ is also glicci. Thus, if Δ is a glicci simplicial complex, then any cone over Δ is also glicci.

Lemma 2.4. [NR08, Remark 2.4(iii)] *Let Δ be a simplicial complex with Stanley–Reisner ideal I_Δ . If Δ is pure and if the deletion Δ_{-k} is Cohen–Macaulay and has the same dimension as Δ , then I_Δ is a basic double link*

of the cone over the Stanley–Reisner ideal of its link $\text{lk}_\Delta(k)$. Conversely, if there exists a vertex k of Δ so that $I_\Delta = x_k I_{\text{lk}_\Delta(k)} + I_{\Delta-k}$ is a basic double link, then k is a weak shedding vertex of Δ .

In order to consider a wider variety of ways in which one squarefree monomial ideal might be a basic double link of another, we will work to drop the requirements that A be a squarefree monomial ideal and that f be a variable of S .

Remark 2.5. As was the setting of [NR08], the squarefree monomial ideals that are the focuses of our study will play the role of C in the equation $C = fB + A$. That is, we will be studying a squarefree monomial ideal C and asking whether or not it might be a basic double link of another (typically squarefree) monomial ideal B . The reason for this choice, as opposed to studying the ideal of interest in the role of B , is that the ideal B is simpler than C in the sense that every monomial in B is a divisor of a monomial in C , and at least one monomial generator of B is a proper divisor of some monomial generator of C . The goal is to gain an understanding of a more complicated ideal by relating it to a simpler ideal.

In light of the result of [MN13] that every arithmetically Cohen–Macaulay subscheme of \mathbb{P}^{n-1} embeds as a gl Ricci subscheme of \mathbb{P}^n , we will work in a fixed polynomial ring in n variables. In particular, once we fix our ambient vertex set for our simplicial complexes, we will never expand this set when performing basic double links.

3. BASIC DOUBLE LINKS OF EDGE IDEALS

In this section we study basic double links of *edge ideals* (see Definition 3.2). Our main theorem (Theorem 3.5) is a description of the form of a basic double link $I(G) = fB + A$ when $I(G)$ is an edge ideal and B is any monomial ideal. We then use this theorem to provide an example of an edge ideal that is Cohen–Macaulay but is not a basic double link of any other monomial ideal B on any homogeneous ideal A . This edge ideal was previously studied in [EVMVT16].

We begin with a straightforward observation that is not specific to monomial ideals.

Lemma 3.1. *Let A , B , and $C = (c_1, \dots, c_r)$ be proper homogeneous ideals of S . Assume that each c_i is a homogeneous polynomial, and set $d = \max_{i \in [r]} \deg(c_i)$. If $f \in S$ is a homogeneous polynomial such that $C = fB + A$ is a basic double link of B on A , then $1 \leq \deg(f) < d$.*

Proof. We proceed by contradiction and suppose that $\deg(f) \geq d$. By definition of basic double link, $B \neq S$, and so every homogeneous element of B has positive degree. Hence, every homogeneous element of fB has degree $> d$. Since each c_i has degree at most d , the condition $C = fB + A$ implies that $c_i \in A$ for all $1 \leq i \leq r$. Thus $C = A$, in violation of the condition $\text{ht}(A) + 1 = \text{ht}(C)$. \square

We now recall the definition of an edge ideal.

Definition 3.2. Let G be a finite simple graph with vertex set $V(G) = \{1, \dots, n\}$ and edge set $E(G)$. The **edge ideal** $I(G)$ of the graph G is the squarefree monomial ideal of S defined by

$$I(G) = (x_i x_j : \{i, j\} \in E(G)).$$

We have following immediate consequence of Lemma 3.1:

Corollary 3.3. *Let $I(G) \subset S$ be an edge ideal, and suppose there exists a basic double link $I(G) = fB + A$ for some homogeneous polynomial $f \in S$. Then $\deg(f) = 1$.* \square

The next lemma will let us place further restrictions on the linear forms f that we consider when thinking about basic double links of edge ideals. This lemma is not specific to monomial ideals. We omit the proof, which is straightforward.

Lemma 3.4. *Let $A, B, C \subset S$ be homogeneous ideals, and let f be a non-zero linear form $f = \sum_{i=1}^n \alpha_i x_i$, $\alpha_i \in \mathbb{k}$. Let $S' = \mathbb{k}[y_1, \dots, y_n]$, and define a ring homomorphism $\phi : S' \rightarrow S$ given by $\phi(y_i) = \alpha_i x_i$ if $\alpha_i \neq 0$ and*

$\phi(y_i) = x_i$ otherwise. Let A', B', C' be the kernels of the induced maps $S' \rightarrow S/A$, $S' \rightarrow S/B$, and $S' \rightarrow S/C$ respectively. Then $C = fB + A$ is a basic double link if and only if $C' = hB' + A'$ is a basic double link where

$$h = \sum_{\{j: \alpha_j \neq 0\}} y_j.$$

□

Let $i \in V(G)$ be a vertex of the graph G . Define the set of **neighbours of i** , $\mathcal{N}(i)$, to be those vertices $j \in V(G)$ such that $\{i, j\} \in E(G)$. If $f = x_{i_1} + \dots + x_{i_r}$, define $\mathcal{N}_f = \left(x_j : j \in \bigcap_{q=1}^r \mathcal{N}(i_q) \right)$.

Theorem 3.5. *Let $I(G) \subset S$ be an edge ideal. Assume there is a basic double link $I(G) = gB + \tilde{A}$ where g is a homogeneous form in S , $B \subset S$ is a monomial ideal, and $\tilde{A} \subset S$ a homogeneous ideal. Then, there is a basic double link $I(G) = fB + A$ where A is a homogeneous ideal and*

- (1) $f = x_{i_1} + \dots + x_{i_r}$ is a sum of distinct indeterminates in S ; and
- (2) $B = I(G) + \mathcal{N}_f$.

Proof. Suppose $I(G) = gB + \tilde{A}$ is a basic double link. By Corollary 3.3, we have $\deg(g) = 1$. Then, by Lemma 3.4 and the assumptions that $I(G)$ and B are monomial ideals, we have a basic double link

$$I(G) = fB + A$$

for some homogeneous ideal A and some sum of indeterminates $f = x_{i_1} + \dots + x_{i_r}$. It remains to prove (2).

We begin by proving that an indeterminate $z = x_i$ is a generator of B if and only if it is one of the indeterminates in \mathcal{N}_f . Indeed, if $z \in B$ then $zf \in I(G)$. Since $I(G)$ is a monomial ideal, we conclude that $zx_{i_q} \in I(G)$ for all $q \in [r]$, and hence $z \in \mathcal{N}_f$. Conversely, if $z \in \mathcal{N}_f$ then $zx_{i_q} \in I(G)$ for each $q \in [r]$, and so $fz \in I(G)$. As $I(G) = fB + A$, we have $fz = fb + a$ for some $b \in B$ and some $a \in A$. Thus, $f(z - b) = a \in A$. Since f is a non-zerodivisor on S/A , we have that $z - b \in A$. As $A \subset B$ by the definition of basic double link, we have $z \in B$ as desired.

We next show that $I(G) + \mathcal{N}_f \subseteq B$. We have already seen that $\mathcal{N}_f \subseteq B$ so it suffices to prove that $I(G) \subseteq B$. So, fix a minimal monomial generator $c \in I(G)$. Because $I(G)$ is an edge ideal, $\deg(c) = 2$. If c is a multiple of some $y \in \mathcal{N}_f$, then $c \in B$. If c is not a multiple of any indeterminate $y \in \mathcal{N}_f$, then c is not a term of any element of fB : if it were, then c would be a term of some fw , where w is an indeterminate in B . But, as shown above, if $w \in B$ then $w \in \mathcal{N}_f$. It follows that c is a term of some element $g \in A$. Since $A \subset B$, we have $g \in B$. As B is a monomial ideal, it follows that $c \in B$.

Finally, we show that $B \subseteq I(G) + \mathcal{N}_f$. Fix some monomial $b \in B$. If $b \in \mathcal{N}_f$ then we are done. So suppose otherwise. We have $fb \in I(G)$ and so $x_{i_q}b \in I(G)$, for all $q \in [r]$ since $I(G)$ is a monomial ideal. Let c_1, \dots, c_r be degree 2 monomial generators of $I(G)$ such that c_q divides $x_{i_q}b$ for each $q \in [r]$. If c_q divides b then $b \in I(G)$ as desired. So we may assume that, for each $q \in [r]$, there is an indeterminate z_q such that $c_q = x_{i_q}z_q$.

Since $c_q \in I(G)$ and $I(G) = fB + A$, and A and B are homogeneous ideals, we have $c_q = fb_q + a_q$ for some $b_q \in B$ of degree 1 and some $a_q \in A$ of degree 2. Write $x_{i_q}b = c_q m_q$. Then,

$$x_{i_q}b = fm_q b_q + m_q a_q,$$

and so, adding up these equalities yields

$$fb = x_{i_1}b + \dots + x_{i_r}b = f(m_1b_1 + \dots + m_rb_r) + (m_1a_1 + \dots + m_ra_r).$$

So, $f(b - m_1b_1 - m_2b_2 - \dots - m_rb_r) \in A$. Since f is not a zero divisor of S/A , we have that $b - m_1b_1 - m_2b_2 - \dots - m_rb_r \in A$. As $A \subset I(G)$, we have that $b - m_1b_1 - m_2b_2 - \dots - m_rb_r \in I(G)$. Since $I(G)$ is a monomial ideal, it will follow that $b \in I(G)$ provided that the coefficient of the monomial b in $b - m_1b_1 - m_2b_2 - \dots - m_rb_r$ is non-zero. To see this, note that no element of \mathcal{N}_f divides b by assumption. On the other hand, since each b_q is a degree 1 element of B , it follows that every term of $m_1b_1 + \dots + m_rb_r$ is divisible by some element of \mathcal{N}_f . Thus, no term of $m_1b_1 + \dots + m_rb_r$ is equal to b . □

Theorem 3.5 leads to the following corollary regarding the maximum length of a linear form that can be used in a basic double link. Recall that the **degree** of a vertex $v \in V(G)$, which we denote by $\deg(v)$, is the number of edges in $E(G)$ which have v as an endpoint.

Corollary 3.6. *Let $I(G)$ be an edge ideal of S , $B \subset S$ a monomial ideal, and $A \subset S$ an arbitrary homogeneous ideal. Assume that $f = x_{i_1} + \dots + x_{i_r}$ is a sum of indeterminates and that $I(G) = fB + A$ is a basic double link. Then $\max\{\deg(v) \mid v \in V(G)\} \geq r$.*

Proof. If $\mathcal{N}_f = \emptyset$ then $B = I(G)$ by Theorem 3.5. Thus, fB contains no element of degree 2, and we conclude from the basic double link $I(G) = fB + A$ that $A = I(G)$, in violation of the condition $\text{ht}(I(G)) = \text{ht}(A) + 1$. Thus, $\mathcal{N}_f \neq \emptyset$.

So, let $x_i \in \mathcal{N}_f$. Then i is a neighbour of i_q for all $q \in [r]$. Thus, $\deg(x_i) \geq r$ and so $r \leq \max\{\deg(v) \mid v \in V(G)\}$. \square

Before we use Theorem 3.5 in a specific example, we consider a possible converse to Theorem 3.5. In Theorem 3.5, we began from the information of a basic double link and constrained precisely (up to scaling the coordinates) the form f and the ideal B . The following example shows that, if one starts with a suitable form f and its corresponding ideal B as in Theorem 3.5 (2), one still need not have a basic double link. That is, even if one chooses an edge ideal $I(G)$, a Cohen–Macaulay, generically Gorenstein, homogeneous ideal A of height one less than the height of $I(G)$, and a sum of indeterminates f that is not a zerodivisor on S/A , it is not guaranteed that $I(G) = fB + A$, where $B = I(G) + \mathcal{N}_f$.

Example 3.7. *Let $I(G) = (x_1x_2, x_2x_3, x_3x_4)$, $A = (x_3x_4)$, and $f = x_1$. Then $\mathcal{N}_f = (x_2)$, and so, if $B = I(G) + \mathcal{N}_f = (x_3x_4, x_2)$, then $I(G) \neq fB + A = (x_1x_2, x_3x_4)$.*

We end this section by considering the edge ideal of the circulant graph $G = C_{16}(1, 4, 8)$ (see Figure 1). The corresponding simplicial complex has dimension 3 and is not vertex decomposable [EVMVT16, Theorem 6.1]. Indeed, the proof of [EVMVT16, Theorem 6.1] shows that the corresponding simplicial complex is also not weakly vertex decomposable.

Example 3.8. *Consider the circulant graph $G = C_{16}(1, 4, 8)$. Its edge ideal $I(G) \subset R = \mathbb{Q}[x_1, \dots, x_{16}]$, which was studied in [EVMVT16], is given by*

$$\begin{aligned} I(G) = & (x_1x_2, x_2x_3, x_3x_4, x_4x_5, x_5x_6, x_6x_7, x_7x_8, x_8x_9, x_9x_{10}, x_{10}x_{11}, x_{11}x_{12}, x_{12}x_{13}, x_{13}x_{14}, x_{14}x_{15}, \\ & x_{15}x_{16}, x_{16}x_1, x_1x_5, x_1x_9, x_1x_{13}, x_2x_6, x_2x_{10}, x_2x_{14}, x_3x_7, x_3x_{11}, x_3x_{15}, x_4x_8, x_4x_{12}, x_4x_{16}, \\ & x_5x_9, x_5x_{13}, x_6x_{10}, x_6x_{14}, x_7x_{11}, x_7x_{15}, x_8x_{12}, x_8x_{16}, x_9x_{13}, x_{10}x_{14}, x_{11}x_{15}, x_{12}x_{16}). \end{aligned}$$

We will next use the results of this section to show that there is no basic double link $I(G) = fB + A$ for any choice of form f , homogeneous ideal A , and monomial ideal B . We need one more lemma, which follows easily from the definition of basic double link.

Lemma 3.9. *Let C be a homogeneous, saturated, unmixed ideal of S . If there exists a basic double link of the form $C = x_iB + A$, then $S/(C + (x_i))$ is Cohen–Macaulay.*

Proof. Since $C = x_iB + A$ is a basic double link, we have that S/A is Cohen–Macaulay and $A : x_i = A$. Thus, $S/(A + (x_i))$ is Cohen–Macaulay. Furthermore, $C + (x_i) = x_iB + A + (x_i) = A + (x_i)$. Therefore, $S/(C + (x_i))$ is Cohen–Macaulay. \square

Proposition 3.10. *For $I(G)$ defined in Example 3.8, there is no basic double link $I(G) = fB + A$ for any monomial ideal B .*

Proof. Suppose there exists a basic double link $I(G) = fB + A$, where f is a homogeneous form, A is a homogeneous ideal, and B is a monomial ideal. By Theorem 3.5 and Lemma 3.4, we may assume that $f = x_{i_1} + \dots + x_{i_k}$ for some $x_{i_j} \in \{x_1, \dots, x_{16}\}$ and $B = I(G) + \mathcal{N}_f$.

By the symmetry of G , we may assume that $x_{i_1} = x_1$. Thus, \mathcal{N}_f is generated by a subset of the set

$$X = \{x_j \mid i \in \mathcal{N}(1)\} = \{x_2, x_5, x_9, x_{13}, x_{16}\}.$$

Since R/B must be Cohen-Macaulay, one checks in Macaulay2 [GS] that the only candidates for B are: $J_1 = I(G) + (x_2, x_9, x_{16})$, $J_2 = I(G) + (x_2, x_5, x_9, x_{16})$, $J_3 = I(G) + (x_2, x_9, x_{13}, x_{16})$, and $J_4 = I(G) + (x_2, x_5, x_9, x_{13}, x_{16})$. However, the only vertex which is a neighbour to both 2 and 16 is vertex 1. This means that none of J_1, J_2, J_3 are of the form $I(G) + \mathcal{N}_f$ for some f . So, it remains to rule out the existence of a basic double link $I(G) = fB + A$ where $B = I(G) + (x_2, x_5, x_9, x_{13}, x_{16})$ and $f = x_1$. One may check in Macaulay2 that $R/(I(G) + (x_1))$ is not Cohen-Macaulay. Hence, by Lemma 3.9, $I(G) = fB + A$ is not a basic double link. \square




FIGURE 1. The circulant graph $C_{16}(1, 4, 8)$. Note that the 16 vertices are equally spaced around the outer boundary of the figure; there is no vertex in the centre.

4. BASIC DOUBLE LINKS AND OTHER SQUAREFREE MONOMIAL IDEALS

In this section, we expand beyond the class of edge ideals to consider other Cohen-Macaulay Stanley-Reisner ideals I_Δ whose associated simplicial complexes are not weakly vertex decomposable.

In particular, in the examples we consider, I_Δ will be known not to admit any basic double link of the form $I_\Delta = x_k I_{\mathbf{k}_\Delta k} + I_{\Delta - k}$. In the first example, we will show that the ideal in question is not a basic double link of any squarefree monomial ideal on any homogeneous ideal. In the second example, we rule out only the case of a basic double link of any squarefree monomial ideal on any other squarefree monomial ideal. We use the second example, which looks superficially quite similar to the first, to highlight how subtle the issue of assessing the possibility of a basic double link is.

This section involved many computer computations, all of which were performed in Macaulay2 [GS].

Example 4.1. *Consider the ideal*

$$I_\Delta = (x_1x_2x_3, x_1x_2x_4, x_1x_3x_5, x_1x_4x_6, x_1x_5x_6, x_2x_3x_6, x_2x_4x_5, x_2x_5x_6, x_3x_4x_5, x_3x_4x_6).$$

The ideal I_Δ corresponds to a triangulation of \mathbb{RP}^2 and was studied as [NR08, Example 5.2] as an example of an ideal whose associated simplicial complex is Cohen-Macaulay but not weakly vertex decomposable. The height of I_Δ is 3.

Proposition 4.2. *For I_Δ defined in Example 4.1, as an ideal of the ring $R = \mathbb{Q}[x_1, \dots, x_6]$, there is no basic double link of the form $I_\Delta = fB + A$ for any squarefree monomial ideal B .*

Proof. Our approach in this proof is to suppose that we have a basic double link $I_\Delta = fB + A$ and find restrictions on f , B , and A . We will use these restrictions find many polynomials that must be in the ideal A with the goal of concluding that the height of A is at least 3. Because $\text{ht}(I_\Delta) = 3$, by the definition of basic double link, A is required to have height 2. Hence, $\text{ht}(A) \geq 3$ will constitute a contradiction.

It follows from Lemma 3.1 that, if I_Δ is a basic double link of any homogeneous ideal B , f must be of degree at most 2. Suppose first that $\deg(f) = 1$. Consider first the case of $f = x_i$ for some $i \in [6]$. The fact that $R/(I_\Delta + (x_i))$ is not Cohen–Macaulay for any $i \in [6]$ is computed in [NR08, Example 5.2]. Hence, by Lemma 3.9 no basic double link of the form $I_\Delta = fB + A$ exists.

Suppose $f = q_i x_{i_1} + \dots + q_n x_{i_n}$ for some $2 \leq n \leq 6$ and constants $0 \neq q_i \in \mathbb{Q}$. By Lemma 3.4, we may assume $q_i = 1$ for all i . If all generators of B have degree greater than 2, then every homogeneous polynomial of fB has degree greater than 3. Thus, the equality $I_\Delta = fB + A$, together with the fact that I_Δ is generated in degree 3, would imply $A = I_\Delta$, a contradiction. Also, because I_Δ is generated in degree 3 and $fB \subseteq I_\Delta$, no form of degree strictly less than 2 can be an element of B .

Hence, there must be at least one generator $z \in B$ of degree exactly 2. Write $z = z_1 z_2$ where $z_1, z_2 \in \{x_1, \dots, x_6\}$ and $z_1 \neq z_2$. Because I_Δ is a monomial ideal and $fz \in I_\Delta$, it must be that $x_{i_j} z_k \in I_\Delta$ for all $j \in [n]$ and $k \in [2]$. A computer computation shows that, for any choice of z_1, z_2 , there are at most two choices of j satisfying $x_{i_j} z_k \in I_\Delta$ for $k = 1, 2$. Hence, we take $n = 2$ and write $f = x_{i_1} + x_{i_2}$.

It is now easy to check computationally that, for each choice of $f = x_{i_1} + x_{i_2}$, there is exactly one $z = z_1 z_2$ satisfying $x_{i_j} z \in I_\Delta$ for $j = 1, 2$. Hence, $B = (z) + B'$ for some B' generated in degrees 3 and higher, in which case fB' is generated in degrees 4 and higher. For each monomial generator μ of I_Δ not divisible by z , there must be an equation of the form $\mu = fb + a$ for some $b \in B$ and $a \in A$. By homogeneity, we may assume that $\deg(fb) = 3 = \deg(a)$. Thus, there is an equation of the form $\mu = \alpha_\mu fz + a$ for some $a \in A$ and $\alpha_\mu \in \{0, 1\}$, and so $\mu - \alpha_\mu fz \in A$ for each such μ .

We now claim that exactly one of $x_{i_1}z$ or $x_{i_2}z$ is an element of A . Indeed, if $x_{i_1}z \notin A$, then, because $x_{i_1}z \in I_\Delta = fB + A$, we must have an expression of the form $x_{i_1}z = fz - a$ with $a \in A$. We now solve $x_{i_2}z = a \in A$. If $x_{i_1}z \in A$ and $x_{i_2}z \in A$, then $fz \in A$. Because $\deg(z) = 2$ and $A \subset I_\Delta$, we know $z \notin A$. Hence, $z \in A : f \setminus A$, in violation of the definition of basic double link.

For each choice of z , each choice of $x_{i_j}z \in A$, and each choice of $\alpha_\mu = 0$ or $\alpha_\mu = 1$ for each monomial generator μ of I_Δ not divisible by z , let $A' \subseteq A$ be the ideal generated by the $\mu - \alpha_\mu fz$ together with $x_{i_j}z$. In all cases, A' has height 3. Hence, $\text{ht}(A) \geq 3$, in violation of the definition of basic double link. Thus, there can be no basic double link $I_\Delta = fB + A$ with B squarefree monomial and f a form of degree 1.

Next, suppose that $f = q_1 y_1 + \dots + q_n y_n$ where $n \geq 1$, $\deg(y_i) = 2$ for all $i \in [n]$, and $0 \neq q_i \in \mathbb{Q}$. Then, by an argument similar to the degree 1 case above, B must contain a monomial generator $w \in \{x_1, \dots, x_6\}$ such that $y_i w$ is a monomial generator of I_Δ for each $i \in [n]$. Because $y_i w \in I_\Delta$, $\deg(y_i w) = 3$, and I_Δ is a squarefree monomial ideal generated in degree 3, each $y_i w$ must be squarefree.

Suppose $n = 1$, and fix $f = q_1 x_{i_1} x_{i_2}$. Then a computer computation show that the restriction $fw \in I_\Delta$ forces there to be exactly 0 or 2 allowable choices of w . Exclude the case of 0, which does not give rise to a basic double link, and call the 2 allowable choices w_1 and w_2 . That is, at least one of w_1 and w_2 is an element of B , and every degree 1 homogeneous element of B is in the ideal (w_1, w_2) .

For each choice of f , one may check computationally that there exists at least one monomial generator μ of I_Δ that is not an element of (fw_1, fw_2) but is divisible by at least one of x_{i_1} or x_{i_2} . Without loss of generality, assume x_{i_1} divides μ .

Because all monomial generators of I_Δ are of degree 3 and every degree 3 homogeneous polynomial in fB is in the ideal (fw_1, fw_2) , the equality $I_\Delta = fB + A$ implies the existence of an equality of the form $\mu = \alpha_1 fw_1 + \alpha_2 fw_2 + a$ for some $a \in A$ of degree 3 and scalars α_i . But then, because x_{i_1} divides μ and f , it must also divide a . Because $A \subset I_\Delta$ and I_Δ is generated in degree 3, $a/x_{i_1} \notin A$. But $(a/x_{i_1})f = a(q_1 x_{i_2}) \in A$, in violation of $A : f = A$. Hence, we cannot have $n = 1$.

If $n \geq 2$, then, for each choice of f there is at most one choice of degree 1 monic monomial w satisfying $wy_i \in I_\Delta$ for all $i \in [n]$. Hence, we may consider 6 symmetric cases $w = x_j$, for a some $j \in [6]$. For a fixed choice of j , consider the possible choices of f . For each j , there are 10 possible f with $n = 2$, 10 with $n = 3$, 5 with $n = 4$, 1 with $n = 5$, and none with $n \geq 6$. We will now work to reduce to the case of all but at most one of the coefficients in f equal to 1 for all for $2 \leq n \leq 5$.

If $n = 2$, in which case $f = q_1y_1 + q_2y_2$, then there is at least one variable $x_{i_1} \neq w$ dividing y_1 but not y_2 and at least one variable $x_{i_2} \neq w$ dividing y_2 but not y_1 . By Lemma 3.4, we may apply the change of variables using $x_{i_1} \mapsto q_1^{-1}x_1$ and $x_{i_2} \mapsto q_2^{-1}x_2$ to assume that $q_1 = q_2 = 1$.

If $n = 3$, the argument is only slightly more complicated: after possibly reordering the y_i , for each possible f , one can always choose a variable $x_{i_1} \neq w$ dividing y_1 , a variable $x_{i_2} \neq x_{i_1}, w$ dividing y_2 , and a variable $x_{i_3} \neq x_{i_1}, x_{i_2}, z$ dividing y_3 but not dividing either y_1 or y_2 . There is then always a change of variables sending each x_i to a non-zero \mathbb{Q} -multiple of x_i so that the coefficients of the terms of the image of f are all 1. For example, if $f = q_1x_2x_3 + q_2x_3x_5 + q_3x_5x_6$ and $w = x_1$, then a satisfying change of variables is $x_2 \mapsto q_1^{-1}q_2x_2$, $x_5 \mapsto q_2^{-1}x_5$, and $x_6 \mapsto q_2q_3^{-1}x_6$. The case $n = 4$ is the same argument.

In each of these cases $n = 2, 3, 4$, for each monomial generator μ of I_Δ , A must contain an element a_μ satisfying $\mu = \alpha_\mu f w + a_\mu$ for $\alpha_\mu \in \{0, 1\}$. For all such choices, $\text{ht}(A) \geq 3 = \text{ht}(I_\Delta)$, in violation of the definition of basic double link, following the argument from the $\deg(f) = 1$ case.

Finally, suppose $n = 5$. In this case, we may not be able to apply a change of variables as in the cases $n = 2, 3, 4$ to assume that $q_i = 1$ for all i . For example, we may have $w = x_1$ and $f = q_1x_2x_3 + q_2x_2x_4 + q_3x_3x_5 + q_4x_4x_6 + q_5x_5x_6$. Up to a choice of q_i , there is one such possibility for each $w = x_i$, $i \in [6]$, which is obtained by some permutation of the indices of the x_i in the form f in the case $w = x_1$ above. Assume without loss of generality that $w = x_1$.

Then the change of variables $x_2 \mapsto q_1^{-1}x_2$, $x_4 \mapsto q_1q_2^{-1}x_4$, $x_6 \mapsto q_2q_4^{-1}x_6$, $x_5 \mapsto q_4q_5^{-1}x_5$ allows us to reduce to the case of $f = x_2x_3 + x_2x_4 + q_3x_3x_5 + x_4x_6 + x_5x_6$ for $q = r/s$ with $r, s \in \mathbb{Z} \setminus \{0\}$ and $\gcd(r, s) = 1$. Then $sf \in \mathbb{Z}[x_1, \dots, x_6]$. Let $\bar{g} = \overline{sf} \in \mathbb{Z}/2[x_1, \dots, x_6]$, where $g = x_3x_5$ if s is even (in which case r must be odd) and $g = x_2x_3 + x_2x_4 + x_3x_5 + x_4x_6 + x_5x_6$ or $g = x_2x_3 + x_2x_4 + x_4x_6 + x_5x_6$, depending on the parity of r , if s is odd.

Let $\hat{A} = A \cap \mathbb{Z}[x_1, \dots, x_6]$ in $S = \mathbb{Z}[x_1, \dots, x_6]$ and \bar{A} be the image of \hat{A} in $T = \mathbb{Z}/2[x_1, \dots, x_6]$. We will show that $\text{ht}(A) \geq 3$, in violation of the definition of basic double link, by showing that $\text{ht}(A) = \text{ht}(\bar{A})$ and then that $\text{ht}(\bar{A}) \geq 3$.

We will show first that $\text{ht}(A) = \text{ht}(\bar{A})$. Let m be the ideal of S generated by $\{x_1, \dots, x_6\}$ and mR the expansion of m to R . Notice that $S_m \cong R_{mR}$ and that $AR_{mR} = \hat{A}R_{mR}$. Because $A \subseteq mR$ and $\hat{A} \subseteq m$, $\text{ht}(A) = \text{ht}(AR_{mR}) = \text{ht}(\hat{A}R_{mR}) = \text{ht}(\hat{A})$.

Because 2 is not a zerodivisor on S/\hat{A} , we know $\text{ht}(\hat{A}) = \text{ht}(\hat{A} + (2)) - 1$. Then because $T/\bar{A} \cong S/(\hat{A} + (2))$ while $\dim(T) = \dim(S) - 1$, we have $\text{ht}(\bar{A}) = \text{ht}(\hat{A} + (2)) - 1$. Combining these equations, we have $\text{ht}(A) = \text{ht}(\hat{A}) = \text{ht}(\bar{A})$.

We will next show that $\text{ht}(\bar{A}) \geq 3$. Let \bar{B} and \bar{I}_Δ be the ideals of $\mathbb{Z}/2[x_1, \dots, x_6]$ generated by the monomial generators of B and I_Δ , respectively, and let \bar{g} and \bar{w} denote the images of g and w , respectively. Note that $\bar{I}_\Delta = \bar{g}\bar{B} + \bar{A}$. Hence, for each monomial generator μ of I_Δ , there must be an element $a_\mu \in \bar{A}$ such that $\mu = \alpha_\mu \bar{g}\bar{w} + a_\mu$ with $\alpha_\mu \in \{0, 1\}$. By the same argument used in the preceding cases, \bar{A} must also contain all monomial generators of \bar{I}_Δ not divisible by \bar{w} . For all choices of g and all choices of α_μ , a computer check establishes $\text{ht}(\bar{A}) \geq 3$.

Hence, there cannot be a basic double link in which $\deg(f) = 2$ either, and so there is no basic double link $I_\Delta = fB + A$ for any squarefree monomial ideal B . \square

In the proof of Proposition 4.2, our contradictions arose through the height of A . For this reason, it was always sufficient to find an ideal contained in A whose height was too large, and we did not require arguments concerning the other possible contents of A . We will also consider an example where height considerations are not adequate and Cohen–Macaulayness must be taken into account. Our next example shows that, when there are candidate ideals for the role of A of the correct height, assessing Cohen–Macaulayness of these candidates can be quite difficult, even with additional simplifying assumptions. Before giving the example, we record a lemma.

Though the following lemma is not directly in [NR08], it is very similar content covered by their Remark 2.4. We include it for completeness.

Lemma 4.3. *If A , B , and I_Δ are squarefree monomial ideals, I_Δ is the Stanley–Reisner ideal of the complex Δ , and $I_\Delta = x_k B + A$ is a basic double link, then B is the Stanley–Reisner ideal of $\text{lk}_\Delta k$ and A the Stanley–Reisner ideal of the cone over Δ_{-k} with apex k .*

Proof. Let Δ_A denote the Stanley–Reisner complex of A and Δ_B the Stanley–Reisner complex of B .

Because $I_\Delta = x_k B + A$ is a basic double link, we know $A : x_k = A$, which is to say that x_k does not divide any of the monomial generators of A . Hence, in order to show that A is the ideal of the cone over Δ_{-k} with apex k , it suffices to show that $A + (x_k)$ is the Stanley–Reisner ideal of Δ_{-k} . Notice that $A + (x_k) = I_\Delta + (x_k)$ and that $I_\Delta + (x_k)$ is the Stanley–Reisner ideal of Δ_{-k} .

For a subset F of $[n]$ satisfying $k \notin F$, we have $F \in \Delta_B$ if and only if $x_F \notin B$ if and only if $x_k x_F \notin I_\Delta$ if and only if $F \cup \{k\} \in \Delta$. For a subset F of $[n]$ with $k \in F$, then $F \notin \Delta_{-k}$, and we claim $x_F \in B$ is not a minimal generator of B . If $x_F \in B$, then $x_k x_F \in I_\Delta$. Because I_Δ is a squarefree monomial ideal and $x_k \mid x_F$, $x_F \in I_\Delta$. Because $x_k \mid x_F$ and $A : x_k = A$, $x_F \in x_k B$, in which case $x_{F \setminus \{k\}} \in B$, and so x_F is not a minimal generator of B . Hence, B is the Stanley–Reisner ideal of $\text{lk}_\Delta k$, as desired. \square

Example 4.4. *Let*

$$\Delta = \{x_1 x_5 x_6, x_2 x_4 x_6, x_1 x_4 x_6, x_2 x_3 x_6, x_1 x_2 x_6, x_3 x_4 x_5, x_1 x_4 x_5, x_2 x_3 x_5, x_1 x_3 x_5, x_2 x_3 x_4\}$$

and I_Δ be the Stanley–Reisner ideal of Δ , in which case

$$I_\Delta = (x_2 x_3 x_4, x_1 x_3 x_5, x_2 x_3 x_5, x_1 x_4 x_5, x_3 x_4 x_5, x_1 x_2 x_6, x_2 x_3 x_6, x_1 x_4 x_6, x_2 x_4 x_6, x_1 x_5 x_6).$$

It was shown in [MT03, V6F10-6] that Δ is a shellable (hence Cohen–Macaulay) but not vertex decomposable simplicial complex and in [NR08, Example 5.4] that it is moreover not weakly vertex decomposable. The height of I_Δ is 3.

Proposition 4.5. *Let $I_\Delta \subseteq \mathbb{Q}[x_1, \dots, x_6]$ be as in Example 4.4. There does not exist a basic double link of the form $I_\Delta = A + fB$ for squarefree monomial ideals A and B and a form f that is a sum of monic monomials.*

Proof. We know from Lemma 3.1 that, if there is a basic double link of the form $I_\Delta = fB + A$, then $\deg(f) < 3$. We will consider 2 broad cases, each of which has several sub-cases.

Case 1 is that $\deg(f) = 1$. It was checked in [NR08, Example 5.4] that Δ is not weakly vertex decomposable. Therefore, by Lemma 4.3, there does not exist a basic double link of the form $I_\Delta = x_k B + A$ for any $k \in [6]$. From this fact, if there is a basic double link of the form $I_\Delta = fB + A$ where f is a form of degree 1 that is a sum of monic monomials, then $f = x_{i_1} + \dots + x_{i_n}$ for some $n \geq 2$. Each monomial generator μ of I_Δ must either be an element of A or satisfy an equation of the form $\mu = b_\mu f + a_\mu$ for some degree two element $b_\mu \in B$ and some $a_\mu \in A$.

The ideal B must have a positive number of degree 2 monomial generators (or else $I_\Delta = A$). Whenever $z \in B$ is a degree 2 monomial, we claim that $zx_{i_j} \in A$ for all but exactly one choice of $j \in [n]$. If $zx_{i_j} \in A$ for all $j \in [n]$, then $z \in (A : f) \setminus A$, in violation of the definition of basic double link. To see that we cannot have $zx_{i_j} \notin A$ for two or more values of j , note first that $zf \in I_\Delta$ implies $zx_{i_j} \in I_\Delta$ for all $j \in [n]$ because I_Δ is a monomial ideal, and so we must have an equality $zx_{i_j} = fz - a$ for some $a \in A$. Because $a = fz - zx_{i_j}$ is a sum of all but one of the monomials of fz and A is a monomial ideal, A contains for all but one choice of zx_{i_j} , as claimed. Note also that A must contain all degree 3 monomial generators of I_Δ not divisible by any of the degree 2 monomials of B .

We are now ready to turn to sub-case 1a. In this case, we assume that B has exactly one degree 2 monomial generator z . From the discussion above, A must contain all but exactly one of the generators of I_Δ . A computer check shows that there are 5 height 2 squarefree monomial ideals contained in I_Δ containing all but exactly one of the generators of I_Δ and that none of these ideals is Cohen–Macaulay.

In sub-case 1b, we assume that B contains 2 or more degree 2 monomials. For each degree 2 monomial $z \in B$ and each summand x_i of f , we must have $zx_i \in I_\Delta$ because I_Δ is a monomial ideal. A computer check shows that, for any set S of 2 or more squarefree monomials of degree 2, there is at most one i so that $zx_i \in I_\Delta$ for each degree 2 generator z of S . Hence, sub-case 1b reduces to the case of $f = x_k$ treated above via [NR08, Example 5.4] and Lemma 4.3.

We now turn to Case 2: $\deg(f) = 2$. In this case, write $f = z_1 + \dots + z_n$ for some $n \geq 1$ where each z_i is a monic monomial of degree 2. Our next goal is to reduce to the case of $n \leq 2$ using our previous work.

Suppose $n = 3$, and write $f = z_1 + z_2 + z_3$. For each choice of $f = z_1 + z_2 + z_3$, let \mathcal{U}_f be the set of subsets U of $\{x_1, \dots, x_6\}$ satisfying $x_i f \in I_\Delta$ for each $x_i \in U$. If $I_\Delta = fB + A$ is a basic double link, the set of variables in B must be an element of \mathcal{U}_f . A computer computation shows that there are 60 choices of f for which $\mathcal{U}_f \neq \emptyset$ and none for which any element of \mathcal{U}_f has size 2 or greater. Thus, for any possible choice of f , fB contains exactly one form of degree 3. By the arguments from previous cases, A must contain all but one of the generators of I_Δ , which is impossible.

Finally, if $n \geq 4$ and $f = z_1 + z_2 + z_3 + \dots + z_n$, then $x_i f \in I_\Delta$ implies $x_i(z_1 + z_2 + z_3) \in I_\Delta$, and so again B has at most one monic monomial generator of degree 1, which we have already established is impossible. Hence, $n = 1$ or $n = 2$.

We next restrict the number of variables that are elements of B to either 1, 2, or 3. For any variable x_i of B , $x_i z_j \in C$ for $j \in [n]$. Using the fact that I_Δ is squarefree and generated in degree 3, no z_j may be divisible by any x_i . Moreover, also as above, A must contain all but exactly one of the terms $x_i z_j$ for each $x_i \in B$. Because $A \neq I_\Delta$ and $\text{ht}(B) = \text{ht}(I_\Delta) = 3$, there must be weakly between 1 and 3 variables x_i in B .

We will now rule out $n = 1$. Suppose $n = 1$ and write $f = z_1$. If B contains exactly one variable x_i , then $x_i z_1$ is the only degree 3 monomial of fB , and so A must contain all but one of the generators of I_Δ , which we have already said is impossible. If B contains exactly two variables x_{i_1} and x_{i_2} , then all monomial generators of I_Δ other than $z_1 x_{i_1}$ and $z_1 x_{i_2}$ must be elements of A . The condition $A : f = A$ implies that a monomial divisible by either variable dividing z_1 must not be a generator of A . Then for all choices of z_1 , x_{i_1} , and x_{i_2} , there is some generator of I_Δ that both must and must not be a generator of A , a contradiction. The argument is the same if B contains 3 of the variables of R . Hence, $n \neq 1$.

The only remaining possibility is $n = 2$. Write $f = z_1 + z_2$, where each z_i is a product of 2 distinct variables. We must consider the possibilities that B contains 1, 2, or 3 variables. These will be sub-cases 2a, 2b, and 2c, respectively.

In sub-case 2a, B contains exactly one variable x_i of R . By an argument similar to the $n = 1$ case, exactly one of $x_i z_1$ and $x_i z_2$ must be a generator of A , and all monomial generators of I_Δ not equal to $x_i z_1$ or $x_i z_2$ must be elements of A . Hence, all but exactly one of the monomial generators of I_Δ must be a generator of A , which is impossible.

In sub-case 2b, B contains exactly two variables x_{i_1} and x_{i_2} . Then A contains all monomial generators of I_Δ not equal to $x_{i_j} z_k$ for $j, k \in [2]$ as well as exactly one of $x_{i_1} z_1$ or $x_{i_1} z_2$ and also exactly one of $x_{i_2} z_1$ or $x_{i_2} z_2$. There are 60 squarefree monomial ideals of height 2 contained in I_Δ and containing all but exactly 2 of the generators of I_Δ . None of these ideals is Cohen–Macaulay.

Finally, we turn to sub-case 2c, in which B contains exactly 3 variables. Because B is height 3, $B = (x_{i_1}, x_{i_2}, x_{i_3})$. Because each $z_j x_{i_k}$ is a degree 3 term of I_Δ , the squarefree monomial ideal, no x_{i_k} can divide either z_j . Because there are 3 distinct x_{i_k} disjoint from the variables dividing the z_j and only 6 variables in R , some x_ℓ must divide both z_1 and z_2 . Because $A : f = A$, x_ℓ cannot divide any generator of A . However, as in previous cases, exactly one of the terms of $(z_1 + z_2)x_{i_k}$ must be a term of A , and so we have a contradiction in this case as well.

Having exhausted all of the possibilities for the number of summands of f and the number of monomial generators of B of minimal degree and found a contradiction in each case, we conclude that no basic double link of the desired form exists. \square

We use heavily in the argument above the fact that A is squarefree monomial. Without that assumption, there is no finite set of possible generators of A nor even a degree bound on the possible homogeneous generators of A , and so the problem does not reduce to a finite check in any obvious way.

5. ELEMENTARY G -BILIAISON AND MONOMIAL IDEALS

In their study of glicci simplicial complexes, Nagel and Römer [NR08, Definition 2.2] introduce **squarefree glicci** simplicial complexes, which are the simplicial complexes of glicci squarefree monomial ideals that can be Gorenstein linked to monomial complete intersections via a series of links in which (at least) every second ideal is squarefree monomial. Nagel and Römer give an example of a glicci monomial ideal that cannot be linked to a complete intersection via only other monomial ideals [NR08, Example 2.1].

In the preceding sections of the present work, we studied when we might hope for (or rule out) basic double links between monomial ideals that do not come from weak vertex decompositions. In the cases of some key examples from the literature of simplicial complexes that are Cohen–Macaulay but not weakly vertex decomposable, we concluded there can also be no basic double link. Motivated by these limitations, we use this section to consider elementary G -biliaison (see Definition 5.1). With an eye towards the possible future use of polarizations to move beyond the squarefree case and because [NR08, Example 2.1] did not involve squarefree monomial ideals, we drop the squarefree requirement throughout this section.

Definition 5.1. Let I , J , and N be homogeneous, saturated, unmixed ideals of the polynomial ring S . Suppose that $\text{ht}(I) = \text{ht}(J) = \text{ht}(N) + 1$. If $N \subseteq I \cap J$, if the localization $(S/N)_P$ is Gorenstein for every minimal prime P of S/N , and if there exists an isomorphism $J/N \rightarrow [I/N](-\ell)$ of graded S/N -modules, then we say that J is obtained from I via an *elementary G -biliaison of height ℓ on N* .

It is known (see, for example, [GMN13, Remark 1.13(3)]) that if J is obtained from I via an elementary G -biliaison on N , then there exists an ideal L so that L is a basic double link of I on N and also a basic double link of J on N . In particular, whenever an ideal J is squarefree glicci via a sequence elementary G -biliaisons, there is a sequence of basic double links connecting J to a complete intersection in which every fourth ideal is squarefree monomial. Proposition 5.2 gives conditions that allow us to choose the ideal L to be monomial.

Proposition 5.2. *Suppose that the monomial ideal $J \subset S$ is obtained from the monomial ideal $I \subset S$ via an elementary G -biliaison on N with isomorphism $\varphi : J/N \rightarrow [I/N](-\ell)$.*

- (1) *If N is a monomial ideal and there is some $a \in J$ not a zerodivisor on S/N so that either a or some lift to S of $\varphi(a + N)$ is a monomial, then there exists a monomial ideal L that is a basic double link of both J and I on N .*
- (2) *If there exists some $a \in J$ not a zerodivisor on S/N such that $\varphi(a + N)$ is the image in I/N of ra for some $r \in S$, then I is a basic double link of J on N .*

Proof. We first consider the form that the isomorphism φ must have before examining claims (1) and (2) individually.

Choose any lift $x \in I$ of $\varphi(a + N)$. Using the fact that φ is an S/N -module isomorphism, we claim that $x(J/N) = a(I/N)$. Indeed,

$$\begin{aligned} x(J/N) &= \{\varphi(a + N)(b + N) \mid b + N \in J/N\} \\ &= \{(a + N)\varphi(b + N) \mid b + N \in J/N\} \\ &= \{a(c + N) \mid c + N \in I/N\} \\ &= a(I/N). \end{aligned}$$

Thus, for every $b \in J$, there exists $c \in I$ and $n \in N$ so that $xb - n = ac$.

We claim that φ is the map determined by multiplication by x/a . Because a acts invertibly on S/N , we may first consider images under φ of elements of $a(J/N)$ and then, by dividing by a , infer the result for all elements

of J/N . Fix $ab + N \in J/N$, and choose n and c so that $xb - n = ac$. We compute

$$\varphi(ab + N) = b\varphi(a + N) = b(x + N) = (bx - n) + N = ac + N.$$

Hence, $\varphi(b + N) = c + N$.

Mechanically, we may execute the map φ on $b + N \in J/N$ by multiplying b by x and then modifying bx by an element of N to obtain a multiple of a and, finally, dividing by a and projecting down to I/N . That is the sense in which we mean that the map φ is multiplication by x/a .

Consider claim (1). Note that because φ is an isomorphism, $\varphi(a + N)$ is also not a zerodivisor on S/N . If a is a monomial, then $aI + N$ is a monomial ideal of S , and if some lift $\widetilde{\varphi(a + N)}$ of $\varphi(a + N)$ to S is a monomial, then $\varphi(\widetilde{\varphi(a + N)})J + N$ is a monomial ideal. Fix some preimage $x \in J$ of $\varphi(a + N)$.

We claim that $aI + N = xJ + N$. It suffices to show $aI \subseteq xJ + N$ and $xJ \subseteq aI + N$. Fix $ai \in aI$. As discussed above, there exists an element $n \in N$ so that $ai + n = xj$ with $j \in J$. Hence, $ai + n = xj \in xJ$, and so $ai = xj - n \in xJ + N$. The other containment is similar. Set $L = aI + N = xJ + N$. Then the monomial ideal L is a basic double link of both I and J .

We now consider claim (2). Because $\varphi(a + N) = ra + N$, we have from the argument above that $ra(J/N) = a(I/N)$ and then, because a acts invertibly on all of S/N , $r(J/N) = I/N$.

Hence, for every element $b \in J$, there exists $c \in I$ and $n \in N$ so that $rb - n = c$. Because $N \subset I$, this implies that $rJ + N \subseteq I$. But, for every $i \in I$, there exists $j \in J$ and $n \in N$ so that $i - n = rj$. Because $N \subseteq rJ + N$, this implies that $I \subseteq rJ + N$.

The equality $I = rJ + N$ shows that I is a basic double link of J on N . \square

Corollary 5.3. *Suppose that $J = I_1, \dots, I_k = I$ is a sequence of homogeneous, saturated, unmixed monomial ideals and that I is a complete intersection. Assume that I_i is obtained from I_{i-1} via elementary G -biliaison for all $1 < i \leq k$ and that, in each case, the hypotheses of at least one of claim (1) or (2) of Proposition 5.2 is satisfied. Then J is glicci via a sequence of basic double links in which all linked ideals are monomial.*

Proof. By Proposition 5.2, either I_i is a basic double link of I_{i-1} or I_i is linked to I_{i-1} by a sequence of two basic double links where the intermediary ideal L_i is monomial. Hence, I is linked to J via a sequence of direct Gorenstein links in which every other ideal (some I_i or L_i) is monomial. \square

Example 5.4. *We note that it is possible to have an elementary G -biliaison $\varphi : J/N \rightarrow I/N$ that does not give rise to a sequence of basic double links of monomial ideals connecting J to I . For example, if $J = (x_1, x_2, x_3)$, $I = (x_1, x_4, x_3)$, and $N = (x_1x_2 - x_1x_4, x_3x_4 - x_3x_2, x_1 + x_3)$ in the ring S , then the map φ that is multiplication by x_4/x_2 gives an elementary G -biliaison, but $x_4/x_2 \notin S$ and $L = x_4J + N = (x_4x_1, x_4x_2, x_4x_3, x_1x_2, x_3x_2, x_1 + x_3)$ is not monomial. Although it is easy to see that I and J are both basic double links of (x_1, x_2x_4, x_3) , this fact is unrelated to the elementary G -biliaison involving φ and N .*

This example gives rise to the following question:

Question 5.5. Suppose the monomial ideal I is obtained from the monomial ideal J via elementary G -biliaison. Must there exist sequence of monomial ideals $I = L_1, \dots, L_k = J$ so that, for each $1 < i \leq k$, either L_i is a basic double link of L_{i-1} or L_{i-1} is a basic double link of L_i ?

REFERENCES

- [BH93] Winfried Bruns and Jürgen Herzog. *Cohen-Macaulay rings*, volume 39 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1993.
- [Cas03] M. Casanellas. Gorenstein liaison of 0-dimensional schemes. *Manuscripta Math.*, 111(2):265–275, 2003.
- [CDH05] Marta Casanellas, Elena Drozd, and Robin Hartshorne. Gorenstein liaison and ACM sheaves. *J. Reine Angew. Math.*, 584:149–171, 2005.
- [CG18] Alexandru Constantinescu and Elisa Gorla. Gorenstein liaison for toric ideals of graphs. *J. Algebra*, 502:249–261, 2018.
- [CMR00] M. Casanellas and R. M. Miró-Roig. Gorenstein liaison of curves in \mathbf{P}^4 . *J. Algebra*, 230(2):656–664, 2000.

- [CMR01] M. Casanellas and R. M. Miró-Roig. Gorenstein liaison of divisors on standard determinantal schemes and on rational normal scrolls. *J. Pure Appl. Algebra*, 164(3):325–343, 2001.
- [CSRT22] Mike Cummings, Sergio Da Silva, Jenna Rajchgot, and Adam Van Tuyl. Geometric vertex decomposition and liaison for toric ideals of graphs, 2022.
- [DNG09] E. De Negri and E. Gorla. G-biliaison of ladder Pfaffian varieties. *J. Algebra*, 321(9):2637–2649, 2009.
- [Eas14] B. Eastman. Circulant. <https://github.com/brydon/circulant>, 2014.
- [EVMVT16] Jonathan Earl, Kevin N. Vander Meulen, and Adam Van Tuyl. Independence complexes of well-covered circulant graphs. *Exp. Math.*, 25(4):441–451, 2016.
- [FK20] Nathan Fieldsteel and Patricia Klein. Gröbner bases and the Cohen-Macaulay property of Li’s double determinantal varieties. *Proc. Amer. Math. Soc. Ser. B*, 7:142–158, 2020.
- [GMN13] E. Gorla, J. C. Migliore, and U. Nagel. Gröbner bases via linkage. *J. Algebra*, 384:110–134, 2013.
- [Gor07] Elisa Gorla. Mixed ladder determinantal varieties from two-sided ladders. *J. Pure Appl. Algebra*, 211(2):433–444, 2007.
- [Gor08] Elisa Gorla. A generalized Gaeta’s theorem. *Compos. Math.*, 144(3):689–704, 2008.
- [Gor10] Elisa Gorla. Symmetric ladders and G-biliaison. In *Liaison, Schottky problem and invariant theory*, volume 280 of *Progr. Math.*, pages 49–62. Birkhäuser Verlag, Basel, 2010.
- [GS] Daniel R. Grayson and Michael E. Stillman. Macaulay2, a software system for research in algebraic geometry. Available at <http://www.math.uiuc.edu/Macaulay2/>.
- [Har02] Robin Hartshorne. Some examples of Gorenstein liaison in codimension three. *Collect. Math.*, 53(1):21–48, 2002.
- [Har07] Robin Hartshorne. Generalized divisors and biliaison. *Illinois J. Math.*, 51(1):83–98, 2007.
- [HU07] Craig Huneke and Bernd Ulrich. Liaison of monomial ideals. *Bull. Lond. Math. Soc.*, 39(3):384–392, 2007.
- [KMMR⁺01] Jan O. Kleppe, Juan C. Migliore, Rosa Miró-Roig, Uwe Nagel, and Chris Peterson. Gorenstein liaison, complete intersection liaison invariants and unobstructedness. *Mem. Amer. Math. Soc.*, 154(732):viii+116, 2001.
- [KR21] Patricia Klein and Jenna Rajchgot. Geometric vertex decomposition and liaison. *Forum Math. Sigma*, 9:Paper No. e70, 23, 2021.
- [KTY13] Kyouko Kimura, Naoki Terai, and Ken-ichi Yoshida. Licci squarefree monomial ideals generated in degree two or with deviation two. *J. Algebra*, 390:264–289, 2013.
- [MN00] J. Migliore and U. Nagel. Lifting monomial ideals. *Comm. Algebra*, 28(12):5679–5701, 2000. Special issue in honor of Robin Hartshorne.
- [MN01] J. C. Migliore and U. Nagel. Liaison and related topics: notes from the Torino workshop-school. volume 59, pages 59–126 (2003). 2001. Liaison and related topics (Turin, 2001).
- [MN02] J. Migliore and U. Nagel. Monomial ideals and the Gorenstein liaison class of a complete intersection. *Compositio Math.*, 133(1):25–36, 2002.
- [MN13] Juan Migliore and Uwe Nagel. Glicci ideals. *Compos. Math.*, 149(9):1583–1591, 2013.
- [MN21] J. Migliore and U. Nagel. Applications of liaison. In *Commutative algebra*, pages 523–568. Springer, Cham, [2021] ©2021.
- [MS05] Ezra Miller and Bernd Sturmfels. *Combinatorial commutative algebra*, volume 227 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2005.
- [MT03] Sonoko Moriyama and Fumihiko Takeuchi. Incremental construction properties in dimension two—shellability, extendable shellability and vertex decomposability. *Discrete Math.*, 263(1-3):295–296, 2003.
- [Ney] Emmanuel Neye. A Gröbner basis for Schubert patch ideals. [arXiv:2111.13778](https://arxiv.org/abs/2111.13778).
- [NR08] Uwe Nagel and Tim Römer. Glicci simplicial complexes. *J. Pure Appl. Algebra*, 212(10):2250–2258, 2008.
- [PB80] J. Scott Provan and Louis J. Billera. Decompositions of simplicial complexes related to diameters of convex polyhedra. *Math. Oper. Res.*, 5(4):576–594, 1980.
- [PS74] C. Peskine and L. Szpiro. Liaison des variétés algébriques. I. *Invent. Math.*, 26:271–302, 1974.

TEXAS A&M UNIVERSITY, DEPARTMENT OF MATHEMATICS, COLLEGE STATION, TX, USA
 Email address: pjklein@tamu.edu

UNIVERSITY OF TORONTO, DEPARTMENT OF MATHEMATICS, TORONTO, ON, CANADA
 Email address: matthew.koban@mail.utoronto.ca

McMASTER UNIVERSITY, DEPARTMENT OF MATHEMATICS AND STATISTICS, HAMILTON, ON, CANADA
 Email address: rajchgot@math.mcmaster.ca