

SNSN-323-63  
November 6, 2016

# Measurement of the differential cross section for t-channel single-top-quark production at 13 TeV

MATTHIAS KOMM<sup>1</sup>, ON BEHALF OF THE CMS COLLABORATION

*Centre for Cosmology, Particle Physics and Phenomenology  
Université catholique de Louvain, B-1348 Louvain-la-Neuve, BELGIUM*

PRESENTED AT

9<sup>th</sup> International Workshop on Top Quark Physics  
Olomouc, Czech Republic, September 19–23, 2016

---

<sup>1</sup>Work supported by Fonds de la Recherche Scientifique (FNRS).

- 1 Introduction
- 2 Event selection
- 3 Signal extraction




Figure 1: Distributions of (left) the transverse W boson mass and (right) the BDT discriminant after requiring  $m_T(W) > 50$  GeV.

- 3.1 Boosted decision tree
- 3.2 Maximum likelihood fitting
- 3.3 Unfolding
- 4 Results
- 5 Conclusion

## ACKNOWLEDGEMENTS




Figure 2: .

## References

- [1] F. A. Mesmer, Proc. Wien. Acad. Sci. **13**, 1564, 1593 (1762).




Figure 3: Normalized differential t-channel single-top-quark cross section as a function of the parton-level top quark (left) transverse momentum and (right) rapidity.