

PHIL 4110: Problem Set 3

Due on February 26, 2024

Professor Lockhart

Matthew Rogers

Problem 1

Solution

Part One

$$s(v) = 1, \quad \mathfrak{M}, s \models \exists x(A(f(z), c) \supset \forall y(A(y, x) \vee A(f(y), x)))$$

Let $\varphi_1 \equiv A(f(z), c) \supset \forall y(A(y, x) \vee A(f(y), x))$

$\mathfrak{M}, s \models \exists x\varphi_1$ iff $\mathfrak{M}, s[m_1/x] \models \varphi_1$ for some $m_1 \in |\mathfrak{M}|$

$\mathfrak{M}, s[m_1/x] \models A(f(z), c) \supset \forall y(A(y, x) \vee A(f(y), x))$ for some $m_1 \in |\mathfrak{M}|$

This is equivalent to the expression $\mathfrak{M}, s[m_1/x] \not\models A(f(z), c)$ or $\mathfrak{M}, s[m_1/x] \models \forall y(A(y, x) \vee A(f(y), x))$ for some $m_1 \in |\mathfrak{M}|$.

Evaluating $\mathfrak{M}, s[m_1/x] \not\models A(f(z), c)$ for some $m_1 \in |\mathfrak{M}|$:

$$\begin{aligned} &= \langle \text{Val}_{s[m_1/x]}^{\mathfrak{M}}(f(z)), c \rangle \notin A^{\mathfrak{M}} \\ &= \langle f^{\mathfrak{M}}(\text{Val}_{s[m_1/x]}^{\mathfrak{M}}(z)), \text{Val}_{s[m_1/x]}^{\mathfrak{M}}(c) \rangle \notin A^{\mathfrak{M}} \\ &= \langle f^{\mathfrak{M}}(1), c^{\mathfrak{M}} \rangle \notin A^{\mathfrak{M}} \\ &= \langle 2, 3 \rangle \notin A^{\mathfrak{M}} = \text{False, since } \langle 2, 3 \rangle \text{ is in } A^{\mathfrak{M}} \end{aligned}$$

Evaluating $\mathfrak{M}, s[m_1/x] \models \forall y(A(y, x) \vee A(f(y), x))$ for some $m_1 \in |\mathfrak{M}|$:

Let $\varphi_2 \equiv A(y, x) \vee A(f(y), x)$

$\mathfrak{M}, s[m_1/x] \models \forall y\varphi_2$ iff $\mathfrak{M}, s[m_1/x][m_2/y] \models \varphi_2$ for some m_1 for any m_2 , such that $m_1, m_2 \in |\mathfrak{M}|$

This is just the expression $\mathfrak{M}, s[m_1/x][m_2/y] \models A(y, x)$ or $\mathfrak{M}, s[m_1/x][m_2/y] \models A(f(y), x)$, for some m_1 for any m_2 .

Evaluating $\mathfrak{M}, s[m_1/x][m_2/y] \models A(y, x)$, for some m_1 for any m_2 :

$$\begin{aligned} &= \langle \text{Val}_{s[m_1/x][m_2/y]}^{\mathfrak{M}}(y), \text{Val}_{s[m_1/x][m_2/y]}^{\mathfrak{M}}(x) \rangle \in A^{\mathfrak{M}} \\ &= \langle m_2, m_1 \rangle \in A^{\mathfrak{M}} \text{ for some } m_1 \text{ for any } m_2, \text{ such that } m_1, m_2 \in |\mathfrak{M}| \\ &\text{We must check for all possible values of } m_2 \text{ in the domain:} \\ &= \langle 1, m_1 \rangle \in A^{\mathfrak{M}} \text{ and } \langle 2, m_1 \rangle \in A^{\mathfrak{M}} \text{ and } \langle 3, m_1 \rangle \in A^{\mathfrak{M}} \end{aligned}$$

Evaluating $\mathfrak{M}, s[m_1/x][m_2/y] \models A(f(y), x)$, for some m_1 for any m_2 .

$$\begin{aligned} &= \langle \text{Val}_{s[m_1/x][m_2/y]}^{\mathfrak{M}}(f(y)), \text{Val}_{s[m_1/x][m_2/y]}^{\mathfrak{M}}(x) \rangle \in A^{\mathfrak{M}} \\ &= \langle f^{\mathfrak{M}}(\text{Val}_{s[m_1/x][m_2/y]}^{\mathfrak{M}}(y)), m_1 \rangle \in A^{\mathfrak{M}} \\ &= \langle f^{\mathfrak{M}}(m_2), m_1 \rangle \in A^{\mathfrak{M}} \text{ for some } m_1 \text{ for any } m_2, \text{ such that } m_1, m_2 \in |\mathfrak{M}| \\ &\text{We must check for all possible values of } m_2 \text{ in the domain:} \\ &= \langle f^{\mathfrak{M}}(1), m_1 \rangle \in A^{\mathfrak{M}} \text{ and } \langle f^{\mathfrak{M}}(2), m_1 \rangle \in A^{\mathfrak{M}} \text{ and } \langle f^{\mathfrak{M}}(3), m_1 \rangle \in A^{\mathfrak{M}} \\ &= \langle 2, m_1 \rangle \in A^{\mathfrak{M}} \text{ and } \langle 3, m_1 \rangle \in A^{\mathfrak{M}} \text{ and } \langle 2, m_1 \rangle \in A^{\mathfrak{M}} \\ &= \langle 2, m_1 \rangle \in A^{\mathfrak{M}} \text{ and } \langle 3, m_1 \rangle \in A^{\mathfrak{M}} \end{aligned}$$

$\mathfrak{M}, s[m_1/x][m_2/y] \models \varphi_2$ becomes the expression:

$$(\langle 1, m_1 \rangle \in A^{\mathfrak{M}} \text{ and } \langle 2, m_1 \rangle \in A^{\mathfrak{M}} \text{ and } \langle 3, m_1 \rangle \in A^{\mathfrak{M}}) \text{ or } (\langle 2, m_1 \rangle \in A^{\mathfrak{M}} \text{ and } \langle 3, m_1 \rangle \in A^{\mathfrak{M}})$$

for some $m_1 \in |\mathfrak{M}|$

Let $m_1 = 3$, then the second half of the 'or' operation is satisfied, since $\langle 2, 3 \rangle, \langle 3, 3 \rangle \in A^{\mathfrak{M}}$.

Therefore, $\mathfrak{M}, s[m_1/x][m_2/y] \models \varphi_2$ and $\mathfrak{M}, s[m_1/x] \models \forall y(A(y, x) \vee A(f(y), x))$

Recall that $\mathfrak{M}, s \models \exists x \varphi_1$ iff $(\mathfrak{M}, s[m_1/x] \not\models A(f(z), c) \text{ or } \mathfrak{M}, s[m_1/x] \models \forall y(A(y, x) \vee A(f(y), x)))$ for some $m_1 \in |\mathfrak{M}|$.

We have shown the second half of the 'or' operation, therefore $\mathfrak{M}, s \models \exists x \varphi_1$ and \mathfrak{M}, s satisfies the original formula.

Part Two

Use the original variable assignment and structure but have $A^{\mathfrak{M}} = \{\langle 1, 3 \rangle \langle 2, 2 \rangle \langle 3, 3 \rangle\}$.

Problem 2

Solution

Part One

$$\varphi \equiv \psi \supset \chi$$

By definition, $\mathfrak{M}, s_1 \models \varphi$ iff $\mathfrak{M}, s_1 \not\models \psi$ or $\mathfrak{M}, s_1 \models \chi$.

By the IH we have $\mathfrak{M}, s_2 \not\models \psi$ or $\mathfrak{M}, s_2 \models \chi$. Then, $\mathfrak{M}, s_2 \models \varphi$.

Part Two

$$\varphi \equiv \forall x \psi$$