

SUPPORTING INFORMATION

Spatial variation in predator diel activity patterns – feral cats avoid red foxes in time, not space

MW Rees, BA Wintle, JH Pascoe, M Le Pla, EK Birnbaum & BA. Hradsky

Table S1: Summary of the number of camera-trap deployments, unique survey sites and 'independent' counts of invasive predator detections across Ecological Vegetation Class groups within the Glenelg and Otway regions, south-west Victoria, Australia.

Vegetation	Region	Sites	Deployments	Fox counts	Cat counts
Dry Forest	Glenelg	25	69	347	9
	Otways	111	314	341	158
Heathland	Glenelg	40	119	265	59
	Otways	3	9	8	6
Heathy Woodland	Glenelg	154	424	574	96
	Otways	82	256	160	66
Herb-rich Woodland	Glenelg	59	373	863	198
	Otways	2	6	3	2
Lowland Forest	Glenelg	383	1046	1900	290
	Otways	52	163	190	35
Swampy Scrub	Glenelg	4	10	19	8
	Otways	36	98	64	88
Wet Forest	Otways	281	780	715	1187
Total		1232	3667	5449	2202

Table S2: Generalised additive model summaries for invasive predator spatiotemporal activity in south-west Victoria, Australia.

Species	Model	EDF	dev.expl	r.sq
fox	1_spatial	772.6469	0.3514	0.1170
fox	2_vegetation_type	766.1054	0.3498	0.1155
fox	3_habitat_type	768.2692	0.3487	0.1139
cat	1_spatial	562.9236	0.2314	0.0772
cat	2_vegetation_type	566.7562	0.2300	0.0769
cat	3_fox_by_habitat_type	585.8580	0.2317	0.0779

Note:

EDF - estimated degrees of freedom of all model terms.

dev.expl - proportion of the null deviance explained by the model.

r.sq - adjusted r-squared value.




Figure S1: Marginal effect of time (model 1) on predators across both study regions in south-west Victoria, Australia.
White crosses depict unique camera-trap sites




Figure S2: Marginal effect of space on predators across south-west Victoria, Australia (model 1). White crosses depict unique camera-trap sites

Glenelg region: Red fox

Space-time interaction effect (excluding marginal effects)




Figure S3: Interaction effect of space-time on feral cat *Felis catus* activity across each hour of the day (0 - 23) in the Glenelg region, Australia (model 1). White crosses depict unique camera-trap sites.

Glenelg region: Feral cat

Space-time interaction effect (excluding marginal effects)




Figure S4: Interaction effect of space-time on feral cat *Felis catus* activity across each hour of the day (0 - 23) in the Glenelg region, Australia (model 1). White crosses depict unique camera-trap sites.

Otway Ranges: Feral cat

Space-time interaction effect (excluding marginal effects)




Figure S5: Interaction effect of space-time on feral cat *Felis catus* activity across each hour of the day (0 - 23) in the Otway Ranges, Australia (model 1). White crosses depict unique camera-trap sites.

Otway Ranges: Red fox




Figure S6: Overall spatial activity of red foxes *Vulpes vulpes* for each hour of the day (0 - 23) in the Glenelg region, Australia (model 1). White crosses depict unique camera-trap sites

Glenelg region: Feral cat




Figure S7: Overall spatial activity of feral cats *Felis catus* for each hour of the day (0 - 23) in the Glenelg region, Australia (model 1). White crosses depict unique camera-trap sites

Otway Ranges: Red fox




Figure S8: Overall spatial activity of red foxes *Vulpes vulpes* for each hour of the day (0 - 23) in the Otway Ranges, Australia (model 1). White crosses depict unique camera-trap sites

Otway Ranges: Feral cat




Figure S9: Overall spatial activity of feral cats *Felis catus* for each hour of the day (0 - 23) in the Otway Ranges, Australia (model 1). White crosses depict unique camera-trap sites