


Predicate Logic

Part 9

1




Predicates

Just the facts...

2

Predicates

- A predicate is a statement about one or more variables
- It is stated as a fact – being true for the data provided




Spring 2020 Sacramento State - Cook - CSc 28 3

3

Predicates

- Predicates express *properties*
- These can apply to a single entity or *relations* which may hold on more than one individual



Spring 2020 Sacramento State - Cook - CSc 28 4

4

Predicate Notation

- It follows the same basic syntax as function calls in Java (and most programming languages)
- However, type case is important:
 - constants start with lower case letters
 - predicates start with upper case letters

Spring 2020 Sacramento State - Cook - CSc 28 5

5

Single variable predicate

- Predicates can have one variable (at a minimum)
- The following sentence states one that the cat named Pattycakes has the "sleepy" property

"Pattycakes The Cat is sleepy"

Spring 2020 Sacramento State - Cook - CSc 28 6

6

Single variable predicate

- Alternatively, we can write that property in predicate form
- "Sleepy" predicate for "Pattycakes" is true
- Note the uppercase and lowercase!

Sleepy(pattycakes)

Spring 2020

Sacramento State - Cook - CSc 28

7

7

Two Variable Predicate

- Predicates can have multiple variables (unlimited actually... well within reason)
- The following is a classic example of a two-variable relationship

$x < y$

Spring 2020

Sacramento State - Cook - CSc 28

8

8

Two Variable Predicate

- The LessThan predicate is true for x, y

LessThan(x, y)

Spring 2020

Sacramento State - Cook - CSc 28

9

9

Predicates Summary

- 1-place predicates assign properties to individuals:
 - ___ is a cat
 - ___ is sleepy
- 2-place assign relations to a pair
 - ___ is sleeping on ___
 - ___ is the capitol of ___

Spring 2020

Sacramento State - Cook - CSc 28

10

10

Predicates Summary

- 3-place predicates assign relations to triples
 - ___ wants ___ to ___
 - Cat named ___ likes to ___ on ___
- Etc...

Spring 2020

Sacramento State - Cook - CSc 28

11

11



Quantified Statements

More Symbols

12

Quantified Statements

- Sometimes we want to say that *every element in the universe* has some property
- Let's say the universe is the people in this Zoom "room" & we want to say "*everyone in the room is awake*"



Spring 2020

Sacramento State - Cook - CSc 28

13

13

Propositional Approach #1

- We can write out a descriptive sentence
- Shortcomings:
 - it is monolithic and inflexible
 - not "mathematical" enough

Everyone in this room is awake.

Spring 2020

Sacramento State - Cook - CSc 28

14

14

Propositional Approach #2

- We can also write out that sentence using a long list of predicates
- So, we list them all or make a pattern
- Shortcomings: cumbersome & verbose

P(moe) and P(larry) and P(curly) and ...

Spring 2020

Sacramento State - Cook - CSc 28

15

15

Limitations of Propositional Logic

- While propositional logic, which we covered, can express a great deal of complex logical expressions, it ultimately is insufficient for all arguments
- Why? The premises (and conclusions) in propositional logic have no internal structure.

Spring 2020


Sacramento State - Cook - CSc 28

16

16

Limitations of Propositional Logic

- For example, it cannot show the validity of Socrates Argument
- This argument states:
*"All humans are mortal.
Socrates is a human.
Therefore, he is mortal."*



Spring 2020

Sacramento State - Cook - CSc 28

17

17

Socrates Argument

- The following is the propositional logic form of the Socrates Argument
- Can we prove the conclusion?

**All humans are mortal
Socrates is a human

Therefore, Socrates is mortal**

Spring 2020

Sacramento State - Cook - CSc 28

18

18

The Socrates Argument

- The following is the argument in normal form
- A problem arises since the validity of this argument comes from the internal structure which propositional logic cannot "see"

H	
S	
—	
M	

Spring 2020


Sacramento State - Cook - CSc 28

19

19

Solution

- It's time to break apart the logic and see the internal structure
- So, ***we are splitting the atom!***



Spring 2020


Sacramento State - Cook - CSc 28

20

20

Why Go Nuclear?

1. Expose the internal structure of those "atomic" sentences
2. Create new terminology to describe the semantics
3. Introduce laws to use and manage them



Spring 2020

Sacramento State - Cook - CSc 28

21

21

New Notation: For All

- The "For-All" symbol states every element x in the universe makes $P(x)$ true
- So, it is true if and only if the ***every element*** x in the universe has P as true

$\forall x P(x)$

Spring 2020

Sacramento State - Cook - CSc 28

22

22

New Notation: Exists

- The "Exists" States at ***least one element*** x in the universe makes $P(x)$ true
- True if just a single P is true

$\exists x P(x)$

Spring 2020

Sacramento State - Cook - CSc 28

23

23

Example: Pineapple Pizza

- Let's create the quantified statement for ***"Someone doesn't like pineapple pizza!"***
- Let's create a predicate $P(x)$ means ***"x likes pineapple pizza"***
- What does someone mean? At least one person?

Spring 2020

Sacramento State - Cook - CSc 28

24

24

Pineapple Pizza: Try #1

- How about the following expression?
- It's **not true** if at least **one** person likes pineapple
- This means *"nobody likes pineapple pizza"*

$$\neg (\exists x P(x))$$

Spring 2020

Sacramento State - Cook - CSc 28

25

25

Pineapple Pizza: Try #2

- We can also negate the predicate
- Means, for at least one person, they dislike pineapple pizza
- "Someone doesn't like pineapple pizza!"*


$$\exists x \neg P(x)$$

Spring 2020

Sacramento State - Cook - CSc 28

26

26




Quantifier Equivalence

Quantifier Conversion

27

Equivalence

- Just like propositional logic, quantitative expressions have equivalencies
- They follow the same basic logic we have seen before



28

Example: Opposite Expression

- Example: "Everyone in the room is awake"
- Let's create the reverse of this expression (*that still says the same thing*)

<input type="radio"/>	
<input type="radio"/>	"Everyone in the room is awake."
<input type="radio"/>	
<input type="radio"/>	

Spring 2020

Sacramento State - Cook - CSc 28

29

29

Example: Opposite Expression

- So, let's just negate the predicate "is awake" into "is asleep"
- Does that work? **No.**

<input type="radio"/>	
<input type="radio"/>	"Everyone in the room is asleep ."
<input type="radio"/>	
<input type="radio"/>	

Spring 2020


Sacramento State - Cook - CSc 28

30

30

Example: Opposite Expression

- Now, let's negate the quantifier "everyone" (for-all) into "someone" (exists)
- The expression below works: **almost**



Spring 2020


Sacramento State - Cook - CSc 28

31

31

Example: Opposite Expression

- Well, what if we change the quantifier "everyone" (for-all) into "someone" (exists)
- The expression below works: **yes**



Spring 2020


Sacramento State - Cook - CSc 28

32

32

Exists and For-All

- The previous two laws allows us to extrapolate two additional laws
- Note: we simply push the negative and remove the double-negation



Spring 2020


Sacramento State - Cook - CSc 28

33

33

Equivalence – Moving Negation

- You can push a negation through a quantifier by toggling the quantifier
- Read the expressions below carefully...




Spring 2020

Sacramento State - Cook - CSc 28

34

34




Conjunction & Disjunction

Breaking Apart and Combining

Conjunction & Disjunction

- Both the Exists and For-All quantifiers can be broken apart (and combined)
- This can occur if the expression contains an AND or an OR



Spring 2020

Sacramento State - Cook - CSc 28

36

36

Exists Disjunction

- If the Exists quantifier is used on a disjunction, it can be broken into two Exists
- This only works with \vee

$$\exists x (P(x) \vee Q(x)) \equiv \exists x P(x) \vee \exists x Q(x)$$

Spring 2020

Sacramento State - Cook - CSC 28

37

37

For-All Conjunction

- If the For-All is used on a conjunction (and), it can be broken into two For-All
- This only works with \wedge

$$\forall x (P(x) \wedge Q(x)) \equiv \forall x P(x) \wedge \forall x Q(x)$$

Spring 2020

Sacramento State - Cook - CSC 28

38


38



Please Wait

CSC 28
will begin shortly
(open the chat window)

39



Bound & Free Variables

Some variables are not variable

40

Bound & Free Variables

- Not all variables used in a quantified expression is treated the same
- Each variable in an expression is either considered "bound" or "free"



Spring 2020

Sacramento State - Cook - CSC 28

41

41

Bound & Free Variables

- A variable is *free* if a value must be supplied to it before expression can be evaluated
- A variable is *bound* if it not free (usually a dummy variable) and contains values that are not needed to be evaluated

Spring 2020

Sacramento State - Cook - CSC 28

42

42

Example 1

- Which variables need we supply a value before the expression can be evaluated?
- Both x and c
- Without knowing both we cannot evaluate the expression (both are free)

$(x^2 < 4 * c)$

Spring 2020

Sacramento State - Cook - CSIS 28

43

43

Example 2

- Which variables need to be supplied before the expression can be evaluated?
- x: no, it is a dummy variable
- c: yes, once we give a value for c, we can evaluate the expression

$\forall x (x^2 < 4 * c)$

Spring 2020

Sacramento State - Cook - CSIS 28

44

44



Multiple Quantifiers

Many E's and A's doing headstands

45

Multiple Quantifiers

- A quantified statement may have more than one quantifier
- In fact, most of the time, statements will contain several



Spring 2020

Sacramento State - Cook - CSIS 28

46

46

Example

- $x > y$ is an expression with two variables
- The expression is true if an x is supplied which is greater than y

$\forall x \exists y P(x, y)$

Spring 2020

Sacramento State - Cook - CSIS 28

47

47

Example

- $\exists y (x > y)$ is an expression with one free variable
- Evaluates to true if x is supplied and there is a y greater than the supplied x

$\forall x \exists y P(x, y)$

Spring 2020

Sacramento State - Cook - CSIS 28

48

48

Example

- $\forall x \exists y (x > y)$ contains no free variables
- Evaluates to true if $\exists y (x > y)$ is true for every x in the universe.

$\forall_x \exists_y P(x, y)$

Spring 2020

Sacramento State - Cook - CSc 28

49

49

Example 2

- The following is an implication with two quantifiers as operands
- It states that whenever " $\forall x P(x)$ " is a true statement, then so is " $\forall x Q(x)$ ".

$\forall_x P(x) \rightarrow \forall_x Q(x)$

Spring 2020

Sacramento State - Cook - CSc 28

50

50



Converting English to Logic

Do it bit by bit

51

Difficult Example

- Let's create a quantified statement for the following logical statement
- *We will go slowly, since this is not easy*

Everyone who has a friend who has Covid will have to be quarantined

Spring 2020

Sacramento State - Cook - CSc 28

52

52

Difficult Example

- "Everyone" is a For-All relationship
- What is everyone referring to? People
- So, the abstract object is a person

\forall_x (if x has a friend with Covid, then x must be quarantined)

Spring 2020

Sacramento State - Cook - CSc 28

53

53

Difficult Example

- So, we can factor it out into the expression below – x is a person
- *Now*, let's look at the sub expression...

\forall_x (if x has a friend with Covid, then x must be quarantined)

Spring 2020

Sacramento State - Cook - CSc 28

54

54

Difficult Example

- The sentence "if *x* has a friend with Covid, *x* must be quarantined" is an implication!
- Let's look at the antecedent (hypothesis)

if *x* has a friend with Covid,
then *x* must be quarantined

Spring 2020

Sacramento State - Cook - CSc 28

55

55

Difficult Example

- How do we write the concept: "*x* has a friend with Covid"?
- They just need a single friend
- So, this is an Exists quantifier

$\exists y$ (*x* is friends with *y*, and *y* has Covid)

Spring 2020

Sacramento State - Cook - CSc 28

56

56

Difficult Example

- Now that we have a basic form of the final version, let's make some predicates
- We will use single letter names for brevity

$F(x, y)$ means "*x* and *y* are friends"
 $C(x)$ means "*x* has Covid"
 $Q(x)$ means "*x* must be quarantined"

Spring 2020

Sacramento State - Cook - CSc 28

57

57

Difficult Example

- This says: "There exists a *person y* where *y* is friends with *person x*, and *y* has Covid"
- Note: *x* is not bound in this expression

$\exists y (F(x, y) \wedge C(y))$

Spring 2020

Sacramento State - Cook - CSc 28

58

58

Difficult Example

- So, what happens if a friend has Covid?
- Then, they must be quarantined
- Note: implication is outside the exists

$\exists y (F(x, y) \wedge C(y)) \rightarrow Q(x)$

Spring 2020

Sacramento State - Cook - CSc 28

59

59

Difficult Example

- Now we can put it all together...
- The following is the quantified expression for our original statement


$\forall x (\exists y (F(x, y) \wedge C(y)) \rightarrow Q(x))$

Spring 2020

Sacramento State - Cook - CSc 28

60

60




Bounded Quantifiers

Hidden Implication
(for those who hate to type)

61

Bound Quantifiers

- Some quantifiers can be more than meets the eye
- For brevity, many predicate and propositional expressions are merged with the \forall and \exists



Spring 2020 Sacramento State - Cook - CSc 28 62

62

Shorthand Notation

- The following type of expression is quite common
- So much so that a shortcut notation is often employed

$$\forall_{\mathbf{x}} (R(\mathbf{x}) \rightarrow P(\mathbf{x}))$$

Spring 2020 Sacramento State - Cook - CSc 28 63

63

Shorthand Notation

- The membership sub-expression is moved to the quantifier's subscript
- This is equivalent to the last

$$\forall_{R(\mathbf{x})} P(\mathbf{x})$$

Spring 2020 Sacramento State - Cook - CSc 28 64

64

Likewise...

- The sub-expression before the implication can be anything
- In this example, $x > 5$ is moved to the subscript

$$\forall_{\mathbf{x}} (\mathbf{x} > 5 \rightarrow P(\mathbf{x})) \equiv \forall_{\mathbf{x} > 5} P(\mathbf{x})$$

Spring 2020 Sacramento State - Cook - CSc 28 65

65