

1. (1 point) Let

$$A = \begin{bmatrix} 8 & -3 & -4 \\ -7 & 5 & 4 \end{bmatrix}.$$

Define the linear transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ by $T(\vec{x}) = A\vec{x}$.

Find the images of $\vec{u} = \begin{bmatrix} -1 \\ 2 \\ 3 \end{bmatrix}$ and $\vec{v} = \begin{bmatrix} a \\ b \\ c \end{bmatrix}$ under T .

$$T(\vec{u}) = \begin{bmatrix} ______ \\ ______ \end{bmatrix}$$

$$T(\vec{v}) = \begin{bmatrix} ______ \\ ______ \end{bmatrix}$$

Correct Answers:

$$\begin{bmatrix} -26 \\ 29 \end{bmatrix}$$

$$\begin{bmatrix} 8*a-3*b-4*c \\ 5*b-7*a+4*c \end{bmatrix}$$

2. (1 point) Consider a linear transformation T from \mathbb{R}^3 to \mathbb{R}^3 for which

$$T\left(\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} -5 \\ 3 \\ -5 \end{bmatrix}, \quad T\left(\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 5 \\ -2 \\ 2 \end{bmatrix}, \quad T\left(\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

Find the matrix A of T .

$$A = \begin{bmatrix} ______ & ______ & ______ \\ ______ & ______ & ______ \\ ______ & ______ & ______ \end{bmatrix}.$$

Correct Answers:

$$\begin{bmatrix} -5 & 5 & -3 \\ 3 & -2 & 1 \\ -5 & 2 & 4 \end{bmatrix}$$

3. (1 point) Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^4$ be the linear transformation defined by

$$T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} 5x-9y \\ 3y-7x \\ x-2y \\ 6y-4x \end{bmatrix}.$$

Find its standard matrix A .

$$A = \begin{bmatrix} ______ & ______ \\ ______ & ______ \\ ______ & ______ \\ ______ & ______ \end{bmatrix}$$

Correct Answers:

$$\begin{bmatrix} 5 & -9 \\ -7 & 3 \\ 1 & -2 \\ -4 & 6 \end{bmatrix}$$

4. (1 point)

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the linear transformation that first reflects points through the x -axis and then reflects points through the line $y = -x$. Find the standard matrix A for T .

$$A = \begin{bmatrix} ______ & ______ \\ ______ & ______ \end{bmatrix}.$$

Correct Answers:

$$\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$

5. (1 point)

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the linear transformation that first rotates points clockwise through 120° ($2\pi/3$ radians) and then reflects points through the line $y = x$. Find the standard matrix A for T .

$$A = \begin{bmatrix} ______ & ______ \\ ______ & ______ \end{bmatrix}$$

Correct Answers:

$$\begin{bmatrix} -0.866025 & -0.5 \\ -0.5 & 0.866025 \end{bmatrix}$$

6. (1 point) To every linear transformation T from \mathbb{R}^2 to \mathbb{R}^2 , there is an associated 2×2 matrix. Match the following linear transformations with their associated matrix.

- ___1. Reflection about the line $y=x$
- ___2. Clockwise rotation by $\pi/2$ radians
- ___3. Reflection about the y -axis
- ___4. The projection onto the x -axis given by $T(x,y)=(x,0)$
- ___5. Counter-clockwise rotation by $\pi/2$ radians
- ___6. Reflection about the x -axis

A. $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$

B. $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$

C. $\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$

D. $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$

E. $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$

F. $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$

G. None of the above

Correct Answers:

- F
- E
- C
- D
- B
- A

7. (1 point) Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the linear transformation defined by

$$T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} -5x - 15y \\ 5x + 15y \end{bmatrix}.$$

Find a vector \vec{w} that is **not** in the range of T .

$$\vec{w} = \begin{bmatrix} \text{---} \\ \text{---} \end{bmatrix}.$$

Correct Answers:

• $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$

8. (1 point) Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be the linear transformation defined by

$$T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} -2x - 2y \\ -4x - 4y \\ 3x - 4y \end{bmatrix}.$$

Find a vector \vec{w} that is **not** in the range of T .

$$\vec{w} = \begin{bmatrix} \text{---} \\ \text{---} \\ \text{---} \end{bmatrix}.$$

Correct Answers:

• $\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$

9. (1 point) Let T be an linear transformation from \mathbb{R}^r to \mathbb{R}^s . Let A be the matrix associated to T .

Fill in the correct answer for each of the following situations.

- ___1. The row-echelon form of A has a column corresponding to a free variable.
- ___2. Every column in the row-echelon form of A is a pivot column.
- ___3. The row-echelon form of A has no column corresponding to a free variable.
- ___4. Two columns in the row-echelon form of A are not pivot columns.
 - A. T is not one-to-one
 - B. T is one-to-one
 - C. There is not enough information to tell.

Correct Answers:

- A
- B
- B
- A

10. (1 point) Let T be an linear transformation from \mathbb{R}^r to \mathbb{R}^s . Let A be the matrix associated to T .

Fill in the correct answer for each of the following situations.

- ___1. Every row in the row-echelon form of A has a pivot.
- ___2. Two rows in the row-echelon form of A do not have pivots.
- ___3. The row-echelon form of A has a pivot in every column.
- ___4. The row-echelon form of A has a row of zeros.
 - A. T is onto
 - B. T is not onto
 - C. There is not enough information to tell.

Correct Answers:

- A
- B
- C
- B